UNIVERSIDAD POLITÉCNICA DE MADRID

Escuela Técnica Superior de Ingeniería Agronómica, Alimentaria y de
Biosistemas

“Drought-stressed tomato plants trigger bottom-up effects on key mite pests”

Tesis doctoral

Miguel González Ximénez de Embún

Ingeniero Agrónomo; MSc Plant Science

Madrid, 2017

Universidad Politécnica de Madrid

Escuela Técnica Superior de Ingeniería Agronómica, Alimentaria y de
Biosistemas

Departamento de Biotecnología

Centro de Investigaciones Biológicas-CSIC

Departamento de Biología Medioambiental

Laboratorio de Interacción Planta-Insecto

Tesis doctoral:

**“Drought-stressed tomato plants trigger bottom-up effects on key
mite pests”**

Autor:

Miguel González Ximénez de Embún

Ingeniero Agrónomo, MSc Plant Science

Directores:

Pedro Castañera Domínguez

Doctor Ingeniero Agrónomo

Félix Ortego Alonso

Doctor en Ciencias Biológicas

Agradecimientos / Acknowledgments

Esta tesis doctoral ha sido realizada en el Laboratorio de Interacción Planta-Insecto del Departamento de Biología Medioambiental del Centro de Investigaciones Biológicas (C.S.I.C.) de Madrid, gracias a la concesión de la beca predoctoral de la Junta de Ampliación de Estudios (Jae-preDoc_2011_00631) del Consejo Superior de Investigaciones Científicas (C.S.I.C.). Para la realización del trabajo se contó con financiación del INIA (GENOMITE, Proposal No 618105 FACCE Era Net Plus-Food security, Agriculture, Climate Change) y del CSIC No 20146754.

Toca ahora recapitular toda la gente que ha formado parte de manera directa o indirecta de esta aventura que es el doctorado. Sé que, dado que son muchos años y muchas experiencias, alguno quedará sin mencionar, igualmente gracias, no me lo tengáis en cuenta.

En primer lugar, quiero dar las gracias a mis dos directores de tesis, Pedro y Felix, por darme la oportunidad de hacer la tesis con vosotros. Porque siempre habéis confiado en mí, realizando la tesis en un tema que nos era lejano a los tres y que hemos sabido llevar a buen puerto. Gracias por todo lo que he aprendido de vosotros tanto a nivel personal como de vuestra amplia experiencia en investigación. Por guiarme estos años, evitando que me dispersara y por vuestra paciencia, dado que reconozco que muchas veces soy un poco caótico.

A Matilde Barón que nos asesoró en el desarrollo de un protocolo de inducción de sequía.

A toda la gente del consorcio Genomite marco en el que se desarrolló esta tesis. En especial a María, Alain y Philipe del CBGP (el de Francia) por facilitarnos la población de *T. evansi*, sus consejos en la cría de ácaros y su acogida en Montpellier, pasamos buenos momentos recolectando ácaros. A Cristina y Vava, por sus consejos y los momentos vividos en Montpellier. A Jerry y Mike por coordinar el proyecto y por sus comentarios como evaluadores externos que han contribuido de manera significativa a mejorar la tesis.

Al laboratorio de Isabel Díaz del CBGP (el de la UPM) por su ayuda en los protocolos de inhibidores de proteasas y qPCR. A Isabel, ya son unos cuantos años que nos conocemos desde las clases de agrónomos, pasando por Chile, gracias por tu dedicación y tu ayuda todos estos años. A Ana y Blanca por su ayuda y por acogerme en el laboratorio como si fuera el mío. A Estrella, la que nunca para, muchas gracias por todo lo que me has enseñado, has sido una

gran maestra y una muy buena compañera, nos queda pendiente coincidir en un congreso. Gracias también por toda tu ayuda en la recta final de la tesis.

Al servicio de química de proteínas, a Emilia y a Javier, siempre dispuestos a ayudar, incluso cuando algún análisis corría prisa.

A la maravillosa gente del laboratorio, a la que daba gusto ver cada mañana, voy a echar de menos las conversaciones durante las comidas, los cafés del 201 ... A Esteban que me pasó el testigo de la cría de ácaros. Al otro Miguel, gracias por tu apoyo en todo lo referente a sequía y que me reafirmó en lo que estaba haciendo, gracias por tus comentarios que han mejorado esta tesis. A Gema y Pedro, por echarme una mano cuando lo necesitaba y alegrar los cafés. A la gente del 201, laboratorio en el que resistí los 5 años. A Matías, fuente de sabiduría en lo personal y laboral; Nuria, siempre con optimismo y con imaginación; Carolina, rodeada de sus animalillos dispuesta siempre a echar una mano; Gabriela, la dj del labo, siempre nos quedará Juan Luis Guerra bailado junto a la poyata; María, que me cedió el testigo, siempre riendo, habría que resucitar el consultorio; Ana, la alimentadora de canciones pegadizas, y conversaciones absurdas. A la del 202, Marisa, con la que hice mis primeros ensayos; Ángel, cualquier tema es bueno para una buena conversación; Elena, creo que hacer un gel no volverá a ser tan divertido; Ana, gracias por reírte de mis tonterías, es bueno tener apoyos; Alberto, que nos sacaba de concierto por Madrid. Al lejano 240, Cristian, por los momentos del congreso y todas las enseñanzas durante las comidas; Enric, siempre con una sonrisa, al final no coincidimos en una excursión a la sierra. A Aitor, mi único patito, que me obligó a ordenar un poco mis ideas. A Matoya, por toda la ayuda en los papeleos, por alimentarnos y porque siempre traías alegría a los cafés.

A Isi, mi “madre putativa” del CIB, gracias por todas las conversaciones y risas compartidas a última hora del día que hacían que saliera del laboratorio con una sonrisa.

To all the people from the IBED in Amsterdam. Thanks to Merijn for allowing me do the 3 months stay on his lab, for making the right questions that made grow as a scientist. Thank you for opening to me the doors of IBED, where I felt like at home. To Joris who introduced me to the *Aculops* world, I was a pleasure work with you. A Juan, que me enseñó los secretos de la HPLC y los volátiles, siempre con paciencia y tranquilidad. To Saioa, Alexandra, Nina, Pascaline, Tom, Bart, Carlos, Josinaldo ... thank you for those beers after work and those bike trips. To Asker, without his appearance I would have been a homeless in Amsterdam.

A la gente del “master del universo” (Pablo, Carlos, Guille, Alex, Marta, Bea, Pio, Barbara) que me sacasteis de mis lecturas del primer año y con los que compartí tantos buenos momentos. Y donde encontré a Óscar, otro que no para, la enciclopedia andante, y si es irreverente, mejor; Cristina, parece que los duendes del laboratorio nos ayudaron a llegar a aquí; Eugenio, la mente cartesiana, aunque no tanto ..., que han acabado siendo grandes amigos, compañeros en los sin sabores de la ciencia.

A toda la gente con la que he compartido momentos en este patio de vecinos que es el C.I.B. Empezaré por arriba, por las chicas de los rubenes a Julia, con la compartí ELISAS, nuestro primer artículo y muchas visitas en las que las conversaciones nos hacía retrasar el trabajo una y otra vez, y tan felices de ello. A Loreine también presente en esas conversaciones en las que recordaba mis tiempos en Chile. A Helga, con mayúsculas, mi paso por el CIB hubiera sido diferente sin ti, tantos cafés y cañas que nos hacían más llevaderas nuestras horas en el labo, tantos momentos compartidos en parques, manifestaciones, excursiones, viajes . . . gracias de verdad. A toda la gente de InvestigAcción (Héctor, Gonzalo, Roberto, ...), que, si no sirvió para cambiar muchas cosas, sirvió para que nos uniéramos, gracias por los buenos ratos compartidos recortando cerebros. A la gente de la primera, Elvira y Estefi, vuestro positivismo se contagia, Silvia, Mónica, Nohemí, Marta por acogernos en vuestros festejos Sheila, imagino que no volverás a tener una camiseta roja. A Emi y Tin, por abrirme el maravilloso mundo del teatro y el swing. Lucía, por ser tan personaja, Ana Pelirroja, Alessandra y tantos otros.

A toda la gente que ha estado a mi lado fuera de la ciencia. A mis amigos, es larga la lista de toda la gente que, durante esta importante etapa de mi vida, y en las anteriores, han estado acompañándome, desde el colegio, el instituto, la escuela, ... Dispuestos a una cerveza un café o una conversación que hicieran olvidar, o al menos relajar los vaivenes de la tesis. A Sam que has sido realmente la víctima de esta tesis, soportando mis cambios de humor que solo tú eras capaz de ver, que hubiera sido de mi sin ‘las vistillas’.

A mi familia, sobre todo a mis padres, pilar fundamental de todas mis decisiones, de todas mis acciones, sin ellos no sería nada. A mi hermana que siempre me ha mantenido los pies en la tierra. A mis abuelos los que no conocí y los que sí, Manolita, como tú decías “Ahora ya está”.

... que culpa tiene el tomate,
que esta tranquilo en la mata ...

Quilapayún

... el hombre no puede ver el viento,
pero las hojas de los árboles que se mueven
denuncian su presencia ...

La voz de mi barril

Joaquín Ximénez de Embún González-Arnau

Index

	Page
Resumen	v
Abstract.....	vii
Abbreviations.....	ix
Chapter 1 GENERAL INTRODUCTION.....	1
1.1 Tomato	5
1.2 Mite pests	5
1.3 Tomato defense response to mites	10
1.3.1 Induced defense: Phytohormone signalling	11
1.3.2 Defensive secondary metabolites and anti-nutritional proteins	13
1.4 Mite adaptation to plant defenses.	16
1.5 Tomato response to drought stress	18
1.6 Drought effect on the tomato-spider mite interaction	21
1.7 Objectives and general outline.	25
Chapter 2 DROUGHT-STRESSED TOMATO PLANTS TRIGGER BOTTOM-UP EFFECTS ON THE INVASIVE <i>Tetranychus evansi</i>.....	27
2.1 Introduction	29
2.2 Materials and Methods	31
2.2.1 Plant material and mite rearing	31
2.2.2 Drought stress regime and experimental design	31
2.2.3 Performance of <i>T. evansi</i> and leaf damage	33
2.2.4 Chemical and biochemical analysis of tomato plant material	33
2.2.5 Proline feeding stimulant test	37
2.2.6 Statistical analysis	38
2.3 Results	38
2.3.1 Effects of drought on tomato plant growth, stomatal conductance and photosynthetic efficiency	38
2.3.2 Effect of drought on <i>T. evansi</i> population growth and leaf damage	40
2.3.3 Effects of drought and <i>T. evansi</i> on plant nutritional composition and defense proteins	41
2.3.4 Enzymatic activities of <i>T. evansi</i> fed on drought-stressed and well-watered tomato plants	43
2.3.5 Proline feeding stimulant assay	45
2.4 Discussion	45

2.5 Conclusions	48
2.6 Supporting information	49
Chapter 3 DROUGHT STRESS IN TOMATO INCREASES THE PERFORMANCE OF ADAPTED AND NON-ADAPTED STRAINS OF <i>Tetranychus urticae</i>	51
3.1 Introduction	53
3.2 Materials and methods	54
3.2.1 Plant material and mite rearing	54
3.2.2 Drought stress regime	55
3.2.3 Bioassays to test drought stress effects on mite performance and plant nutritional composition	56
3.2.4 Bioassays to test drought stress effects on plant defense proteins and mite enzymatic activities	57
3.2.5 Chemical and biochemical analysis	57
3.2.6 Statistical analysis	59
3.3 Results	60
3.3.1 Effects of drought stress on stomatal conductance, photosynthetic efficiency and tomato plant growth	60
3.3.2 Effects of drought stress on <i>T. urticae</i> (TA and TNA strains) population growth and leaf damage	60
3.3.3 Changes in plant nutritional composition induced by drought stress and <i>T. urticae</i> (TA and TNA)	62
3.3.4 Effect of drought stress and <i>T. urticae</i> (TA and TNA) on plant defense proteins	64
3.4 Discussion	66
3.5 Conclusions	69
3.6 Supporting information	70
Chapter 4 DROUGHT STRESS PROMOTES THE COLONIZATION SUCCESS OF A HERBIVOROUS MITE THAT MANIPULATES PLANT DEFENSES	75
4.1 Introduction	77
4.2 Materials and methods	79
4.2.1 Plant material and mite rearing	79
4.2.2 Drought stress regime	79
4.2.3 Bioassays	80
4.2.4 Chemical and biochemical analysis	81
4.2.5 Quantification of phytohormones by means of LC-MS	82
4.2.6 Quantification of gene expression via qRT-PCR	83
4.2.7 Statistical analysis	84

4.3 Results	85
4.3.1 Effects of drought on <i>Aculops lycopersici</i> population growth and plant damage.	85
4.3.2 Effects of drought on stomatal conductance and tomato plant growth.	85
4.3.3 Changes on plant nutritional composition induced by drought and TRM.	86
4.3.4 Effect of drought and TRM on tomato plant defense: phytohormones, defense genes and defense proteins.	87
4.4 Discussion	90
4.5 Conclusions	93
4.6 Supporting information	94
Chapter 5 PLANT-MEDIATED EFFECTS OF WATER DEFICIT ON THE PERFORMANCE OF <i>Tetranychus evansi</i> ON TOMATO DROUGHT-ADAPTED ACCESSIONS	99
5.1 Introduction	101
5.2 Materials and methods	102
5.2.1 Plant material and mite rearing	102
5.2.2 Drought stress regime	103
5.2.3 Bioassays	103
5.2.4 Chemical and biochemical analysis	104
5.2.5 Statistical analysis	106
5.3 Results	106
5.3.1 Effect of drought on stomatal conductance, photosynthetic efficiency and tomato plant growth.	106
5.3.2 Effect of drought on <i>T. evansi</i> performance.	107
5.3.3 Changes in plant nutritional composition induced by drought and <i>T. evansi</i> .	109
5.3.4 Effect of drought and <i>T. evansi</i> on tomato defense proteins.	110
5.4 Discussion	113
5.5 Conclusions	115
5.6 Supporting information	116
Chapter 6 GENERAL DISCUSSION	119
Chapter 7 CONCLUSIONS	125
Chapter 8 REFERENCES.....	129

Resumen

El incremento de los periodos de sequía como consecuencia del cambio climático afectará a cultivos sensibles a la sequía como el tomate, cuya producción en el área mediterránea se efectuará, previsiblemente, con un menor número de riegos. En esta tesis se ha investigado el efecto de estrés abióticos (sequía) y bióticos (infestación por ácaros) en plantas de tomate. Concretamente, se evaluó: 1) el efecto de la sequía sobre la biología de los ácaros fitófagos *Tetranychus urticae*, *Tetranychus evansi*, y *Aculops lycopersici* en plantas de tomate (cv Moneymaker); 2) los cambios morfológicos, fisiológicos y moleculares que se producen en la planta como respuesta a ambos estreses (por separado y combinados); y 3) el efecto de accesiones de tomate adaptadas a sequía en la biología de *T. evansi*.

Los datos obtenidos revelan que la sequía favorece el desarrollo de *T. evansi* en plantas de tomate, como indican el incremento significativo en el daño foliar (1,5 veces), el número de huevos (2 veces) depositados a 4 días post-infestación (dpi) y en las formas móviles (1,5-2 veces) encontradas a los 10 días en plantas estresadas. Los niveles de algunos aminoácidos esenciales (histidina, isoleucina, leucina, tirosina, valina) y azúcares libres fueron inducidos por sequía en combinación con la infestación con ácaros. Prolina, un aminoácido no esencial, fue altamente inducido, y se pudo demostrar que estimula la fecundidad y la alimentación del ácaro cuando se añadió a discos foliares en niveles similares a los observados a 10 dpi en plantas estresadas. Los niveles de proteínas de defensa del tomate se vieron también afectados por ambos estreses, pero *T. evansi* fue capaz de eludir sus efectos nocivos.

Tanto una línea de *T. urticae* adaptada a tomate (TA) como una no adaptada (TNA) se beneficiaron del incremento en el valor nutricional (aumento de los niveles de aminoácidos y azúcares libres) de las plantas de tomate como respuesta a sequía. La composición nutricional de las hojas de tomate no se alteró por la infestación con ácaros, con la excepción de un aumento de la concentración de azúcares libres. Las proteínas de defensa de tomate fueron inducidas por los ácaros y la sequía. Sin embargo, la inducción de inhibidores de proteasa (IPs) fue superior en plantas infestadas con ácaros TNA que en aquellas infestadas con TA. El mejor desarrollo de la línea TA puede ser explicado por cambios en la fisiología digestiva (actividades cisteín y aspartil proteasa y α -amilasa) y de enzimas de detoxificación (actividad esterasa) de los ácaros, así como por la atenuación de las defensas de la planta (IP).

El crecimiento de la población de *A. lycopersici* fue más rápido y causó mayor daño en plantas de tomate estresadas hídricamente. Ello puede ser debido al incremento en los niveles de proteína total y de varios aminoácidos libres en plantas infestadas por el ácaro. La infestación promueve la respuesta ligada a ácido salicílico (SA) y la expresión de ácido jasmónico (JA) y sus derivados, así como la actividad de inhibidores de cisteín proteasa, polifenol oxidasas y peroxidasas (POD). La sequía redujo la expresión de genes marcadores de JA y la actividad de inhibidores de serín proteasas y POD, alterando, además, los niveles de algunos aminoácidos libres. Cuando ambos estreses se combinan la sequía tuvo un efecto antagónico sobre los niveles de POD y JA inducidos por el ácaro y sinérgico en la acumulación de los niveles de azúcares libres y SA.

Finalmente, se observó que el déficit hídrico tuvo un efecto diferenciado en el desarrollo de *T. evansi* en cuatro accesiones de los tomates adaptados a sequía, 'Tomàtiga de Ramellet' (TR). En las accesiones TR61 y TR154 la sequía aumentaba el desarrollo del ácaro, mientras que en las TR58 y TR126 no. Se encontró una clara relación entre los cambios en los nutrientes de la planta y el desarrollo del ácaro. Los aminoácidos esenciales libres se acumularon en las plantas de tomate en las que la sequía aumentaba el desarrollo del ácaro (TR154 y Moneymaker). Sin embargo, dicha acumulación no se produjo en TR126 en la que la sequía no tuvo efecto en el ácaro. Además, la inducción de la defensa del tomate por *T. evansi* fue superior en TR126 y TR154 que en Moneymaker, lo que puede explicar el peor desarrollo del ácaro en dichas accesiones.

Estos datos revelan que los cambios inducidos en la planta por la sequía y el ácaro aumentan el valor nutricional de la planta y como consecuencia el desarrollo de los ácaros. Además, presenta un marco experimental para que la búsqueda de accesiones de tomate resistente a sequía incluya la resistencia a ácaros herbívoros, en función de los cambios metabólicos de la planta. Estos hallazgos son especialmente importantes en la mitigación de los efectos del cambio climático en las áreas de producción de tomate, y para el manejo y predicción de los brotes de ácaros fitófagos.

Abstract

The increase on drought periods as consequence of climate change will affect crops sensitive to drought, like tomato whose production in Mediterranean areas is expected to be under deficit irrigation schedules. In this thesis, it has been established the effect of both abiotic (drought) and biotic (mite infestation) stresses on tomato plant, specifically: 1) the effect of drought-stressed tomato plants (cv. Moneymaker) on the performance of three key mite pests, the spider mites *Tetranychus urticae* and *T. evansi*, and the eriophyid mite *Aculops lycopersici*; 2) the morphological, physiological and molecular changes induced in tomato plants in response to both stresses (either alone or in combination) and their effects on the mite's performance; and 3) the plant-mediated effects of water deficit on the performance of *T. evansi* on tomato drought-adapted accessions.

Our data reveal that *T. evansi* caused more leaf damage (1.5 fold) to drought-stressed tomato plants. Mite performance was also enhanced, as revealed by significant increases of eggs laid (2 fold) at 4 days post infestation (dpi), and of mobile forms (1.5-2 fold) at 10 dpi. The levels of several essential amino acids (histidine, isoleucine, leucine, tyrosine, valine) and free sugars in tomato leaves were significantly induced by drought in combination with mites. A non-essential amino acid, the osmolite proline, was strongly induced and stimulated mite feeding and egg laying when added to tomato leaf disks at levels recorded at 10 dpi. Tomato plant defense proteins were also affected by drought and/or mite infestation, but *T. evansi* was capable of circumventing their potential adverse effects.

Both, tomato adapted (TA) and non-adapted (TNA) strains of *T. urticae* benefit from the improved nutritional value of tomato plants induced by drought stress (increased concentrations of essential amino acids and free sugars). Mite infestation alone had almost no effect on the nutritional composition of tomato leaves, with the exception of an increase of free sugars. Tomato plant defense proteins were induced by both drought stress and mite infestation. However, the induction of protease inhibitors was higher in tomatoes exposed to mites from the TNA strain than from the TA strain. The better performance of the TA strain could be associated to both changes in the digestive (cysteine and aspartyl protease and α -amylase activities) and detoxification (esterase activity) physiology of the mites and the attenuation of some of the plant's defenses (protease inhibitors).

A. lycopersici population grew faster and caused more damage on drought-stressed tomato plants. This finding can be related to the increased levels of total protein and several free amino acids in tomato-infested plants. Mite infestation promoted the salicylic acid (SA) response and up-regulated the expression of jasmonic acid (JA) and derivatives, as well as the activity of cysteine protease inhibitors, polyphenol oxidase and peroxidase (POD). Drought stress, in turn, reduced the expression of JA marker genes and the activity of serine protease inhibitors and POD, and altered the levels of some free-amino acids. When combined, drought stress antagonized the accumulation of POD and JA by mite infestation and synergized accumulation of free sugars and SA.

Finally, it has been tested the effect on *T. evansi* performance of four accessions of the drought-adapted tomatoes, ‘Tomàtiga de Ramellet’ (TR) under water shortage. In the accessions TR61 and TR154 it was observed an enhancement of mite performance by drought, but not in the TR58 and TR126. A clear link could be established between changes in plant nutritional value and mite performance. Soluble free essential amino were accumulated on tomato plants where mite performance was enhanced under drought stress (TR154 and Moneymaker). This induction did not occur in TR126, where mite performance was not altered. Furthermore, the induction of plant defences in response to *T. evansi* infestation was stronger in TR126 and TR154 than in Moneymaker, which might be a factor contributing to its lower performance on these TR accessions..

These data reveal that the changes induced in the plant by drought and mite infestation increase plant nutritional value and mite performance. Furthermore, it provides an experimental framework to screening for drought-resistant tomato accessions that will be at the same time resistant to herbivore mites, depending on the metabolic changes on the plant. These findings are especially important in the task of adapting area-wide tomato production to mitigate the effects of climate change, and for the management and prediction of herbivore mite proliferation.

Abbreviations

θ	Soil Volumetric Water Content
1-NA	1-naphthyl acetate
aa	Amino Acids
ABA	Absciscic Acid
Ala	Alanine
AMC	7-Amino-4-methylcoumarin
ANOVA	Analysis Of Variance
AOC	Allene Oxide Cyclase
AOS	Allene Oxide Synthase
Arg	Arginine
Asp	Aspartic Acid
CABI	Centre For Agricultural Bioscience International
cDNA	Complementary Deoxyribonucleic Acid
CDNB	1-chloro- 2,4-dinitrobenzene
C:N	Carbon: Nitrogen Ratio
Ct	Cycle Treshold
Cys	Cysteine
D	Drought
DMSO	Dimethyl Sulfoxide
DNA	Deoxyribonucleic Acid
dpi	Days Post Infestation
DTT	Dithiothreitol
DW	Dry Weight
EDTA	Ethylenediaminetetraacetic acid
ESYRCE	Encuesta Sobre Superficies Y Rendimientos De Cultivos/ Spanish Survey of Surfaces and Crop Yields
ET	Ethylene
FAO	Food And Agriculture Organization of the United Nations
Fv/Fm	Maximum Quantum Yield Of Photosystem II Photochemistry
FW	Fresh Weight
Glu	Glutamic acid

Gly	Glycine
gs	Stomatal Conductance
GST	Glutathione S- Transferase
His	Histidine
I	Mite Infestation
Ile	Isoleucine
IPCC	Intergovernmental Panel On Climate Change
JA	Jasmonic Acid
JA-Ile	Jasmonic Acid Isoleucine
JAZ	Jasmonate Zim Domain
<i>JIP-21</i>	Jasmonate Inducible Protein 21 Gene
LC-MS	Liquid Chromatography- Mass Spectrometry
Leu	Leucine
LOX	Lipoxigenases
LpNa	L-leucine p-nitroanilide
LSD	Least Significant Difference
Lys	Lysine
MCA	7-Methoxycoumarin-4-Acetyl -Pro-Leu-Gly
Met	Methionine
MocAc- GKPILFFRLK (Dnp)-D-R NH ₂	7-Methoxycoumarin-4-Acetyl- Gly- Lys- Pro- Ile- Leu- Phe- Phe- Arg- Leu- Lys(Dnp)- D-Arg-NH ₂
MYC-2	Defense Transcription Factor
NADP	Nicotinamide Adenine Dinucleotide Phosphate
NRQ	Normalized Relative Quantity
OPDA	12-Oxo-Phytodienoic Acid
OPR3	OPDA reductase
PE	Primer Efficiency
Phe	Phenylalanine
PI	Protease Inhibitor
<i>PI-II_f</i>	Protease Inhibitor II _f Gene
PPO	Polyphenol Oxidase

<i>PPO-F</i>	Polyphenol Oxidase- F Gene
<i>PR-P6</i>	Pathogenesis Related Protein P6 Gene
Pro	Proline
POD	Peroxidase
qRT-PCR	Quantitative Real Time – Polymerase Chain Reaction
RNA	Ribonucleic Acid
SA	Salicylic Acid
SAR	Systemic Acquired Resistance
SCF ^{COI1}	Coi1 E3 ubiquitin-ligase complex
SDS	Sodium Dodecyl Sulfate
Ser	Serine
SucAAPF-AMC	Suc-Ala-Ala-Pro-Phe-7-amido-4-methylcoumarin
SW	Saturation Weight
TA	Tomato Adapted
TD	Threonin Deaminase
<i>TD2</i>	Threonin Deaminase 2 Gene
TF	Transcription Factor
Thr	Threonine
TNA	Tomato No Adapted
TR	Tomàtiga de Ramellet
Tris	Tris(hydroxymethyl)aminomethane
TRM	Tomato Russet Mite
Tyr	Tyrosine
Val	Valine
v/v	Volume / Volume
v/v/v	Volume/Volume/Volume
w/v	Weight / Volume
Z-FR-AMC	N-carbobenzoxyl-oxy-Phe-Arg-7-amido 4-methylcoumarin
Z-LA-AMC	N-carbobenzoxyl-oxy -L-Arg-7-amido-4-methylcoumarin
Z-RR-AMC	N-carbobenzoxyl-oxy-Arg-Arg-7-amido-4-methylcoumarin
Z-VAN-AMC	N-carbobenzoxyl-oxyVal-Ala-Asn-7-amido-4-methylcoumarin

Chapter 1

GENERAL INTRODUCTION

The expected increase of the global population to about 9.000 million in 2050 will require a major challenge in crop production, as we need to produce about 70-100% more food, with similar arable land, less water availability and without increasing agriculture's environmental footprint (Godfray et al., 2010; Mueller et al., 2012). Water has traditionally been considered the main factor limiting crop production and food security in much of the world where rainfall is insufficient to meet crop demand. Climate change will represent another major risk for long-term food security, as crop yield in many areas will decrease due to the frequency of extended drought periods (Gornall et al., 2010). According to climate prediction models, a gradual increase in temperature with more frequent heat waves and an increase in drought events is expected that, in case of mid-continental and Mediterranean climatic areas, will be more frequent and severe (IPCC, 2013). The intimate relationship between plants and herbivores, which in some cases results in devastating crop losses, will be altered by global climate change. In the case of arthropods, the increase in temperature will shorten their life cycle, having as consequences more generations per year and an earlier appearance of the pest (Luedeling et al., 2011; Delucia et al., 2012; Bebber, 2015). In addition, phytophagous pests will expand their distribution area as a consequence of increases in northern temperatures, a process that will be helped by trade, a factor responsible for introduction of invasive species (Parmesan, 2006; Meynard et al., 2013; Bebber, 2015).

To increase food security it is crucial to reduce crop losses due to biotic and abiotic stresses. Among the different abiotic stresses, water shortage is considered to be the most damaging to crop productivity and it is expected to cause global grain yield losses of up to 30% by 2025 as compared to current yields (Foolad, 2007; World Economic Forum Water Initiative, 2009; Mir et al., 2012). It is considered by the FAO, World Bank and the Red Cross Foundation as the cause for important economic losses worldwide, being especially severe in Sub-Saharan Africa, Central and South America, southern Europe, the Middle East, and southern Australia (Figure 1.1). Despite all pesticide use and other non-chemical pest controls, crop yield losses due to pests are about 40% of all potential food production worldwide. Arthropod pests cause an estimated 14% crop loss, plant pathogens about 13%, and weeds a 13% loss worldwide. The value of total crop losses is estimated to be close to 2,000 billion \$ per year (Pimentel, 2009).

Traditionally, the effect of drought stress and pests on plants has been studied separately, however, in nature biotic and abiotic stresses often co-occur, interacting with each other. This interaction can be positive or negative in terms of plant resistance to herbivory (Maxmen, 2013; Suzuki et al., 2014; Atkinson et al., 2015), since drought induces physiological changes in the plant that can modify its defenses and nutritional value (Mattson and Hack, 1987; DeLucia et al., 2012). Thus, understanding ecological consequences of arthropod pests and water deficit interaction is becoming increasingly important (Tylianakis et al., 2008). In this context, we have selected for this study a summer-irrigated crop, tomato, which is one of the most important vegetable crops worldwide, and highly demanding of water supply (Foolad, 2007; Steduto et al., 2012). Therefore, it is likely that it is affected by deficit irrigation (drought stress) that could induce morphological and nutritional changes with a high impact in its interactions with arthropod pests, such as mites, and might have a negative impact on area-wide tomato production.

Figure 1.1 Global distribution of risk of economic losses due to drought. Data are divided in three categories of risk index represented by three colours: low (blue), orange (medium) and red (high). Adapted from Dilley et al. (2005).

1.1 Tomato

The cultivated tomato, *Solanum lycopersicum* L. (formerly *Lycopersion esculentum* Mill) is a Solanaceous crop originated from the Andean region in South America, and was introduced to Europe in the sixteenth century, where it has a secondary centre of diversity. Tomato cultivation is concentrated in semiarid regions, where it needs irrigation and where drought events associated with climate change are expected to be more frequent (Galmes et al., 2011; Nankishore and Farrell, 2016). Tomato is the most important vegetable crop worldwide (Lin et al., 2014), being cultivated both in the greenhouse and in open field conditions in temperate Mediterranean climates. World production was estimated to be 164.5 million tonnes in 2013. More than 38 million tonnes per year are grown for the processing industry, making tomato the world's leading vegetable for processing (Steduto et al., 2012). Spain is the third most important world exporter (after Mexico and The Netherlands) with about 1 million tonnes exported in 2013 (FAOSTAT Database, www.fao.org/faostat/es/#data, accessed December 2016). In Spain, most of the tomato for industry is produced in the open field with about 99.7 % of the cultivated area under irrigation in 2015. Fresh tomato production is mainly in greenhouses, and the irrigated cultivated area did increase from 80% in 2009 to 97% in 2015 (ESYRCE, 2015). The expected reduction of water availability in the future due to climate change will lead to improve water use efficiency through the application of deficit irrigation, which can have large consequences for tomato production, as deficient irrigation might cause up to a 50% reduction in yield (Cantore et al., 2016). The high sensitivity of tomato to drought under a future climate change scenario will foster the implementation of drought-adapted/tolerant varieties in area-wide tomato production and the investigation of their interactions with mite pests.

1.2 Mite pests

Mites belong to the phylum Arthropoda, subphylum Chelicerata, class Arachnida, the second largest group of terrestrial animals, after insects (Krantz and Lindquist, 1979). Among them, only a few families of the subclass Acari utilize living plants as a food source, the main phytophagous groups being those that belong to the superfamilies Tetranychoida and Eriophyoidea (Vacante, 2016).

***Tetranychus urticae* Koch (Acari: Tetranychidae)**, the two-spotted spider mite, is a very polyphagous pest with a worldwide distribution. This small arthropod is about 0.5 mm long and is usually yellowish-green in colour with two dark patches on the body, though the colour can vary depending of multiple factors like diet or physiological stage (Figure 1.2A). Its life cycle is composed of the egg, the larva, two nymphal stages (protonymph and deutonymph) and the adults. All five mite stages, except eggs, are motile and have four pair of legs except for larvae that have three pairs (Lindquist, 1985). The duration of the life cycle depends mainly on the temperature and the host, with a mean generation time of about 10-15 days under optimal developmental temperatures (27-30 °C) (Kasap, 2004; Riahi et al., 2013). Adult females have an oval shape with a lifespan of 10 to 30 days and an average oviposition rate on tomato of 20-40 eggs during their life cycle, with many overlapping generations per year (Marinosci et al., 2015). Unfertilized eggs (haploid) develop into male offspring (which have a characteristic pointed shape), whereas fertilized eggs (diploid) produce female offspring (Macke et al., 2010). The mites produce webs over the leaf surface of their host plants that protect them against adverse climate conditions, predators and acaricides, and can help communication and dispersal (Sarmiento et al., 2011a).

T. urticae is an extremely polyphagous species, as its host range includes more than 1100 species from 140 different plant families, including 150 plants of economic importance (Migeon and Dorkeld, 2006-2016). However, local populations can become adapted to particular hosts, performing poorly when they are switched to another host (Gotoh et al., 1993; Agrawal et al., 2002, Magalhaes et al., 2009). To feed on plants, *T. urticae* mouthparts are adapted to pierce the plant tissues, consisting of an elongated and retractable cheliceral stylet, which is connected to the pharynx and salivary glands (Van der Geest, 1985; Bensoussan et al., 2016). When feeding, they introduce the stylet between epidermal pavement cells or through a stomatal opening, without damaging the epidermis, and inject saliva on a single parenchyma cell each time, without damaging the surrounding cells (Figure 1.2 C). The saliva predigests the cell content which afterwards is imbibed (Van der Geest et al., 1985; Bensoussan et al., 2016). Moreover, salivary secretions contain a wide range of proteins that can potentially interfere with plant defenses (Jonckheere et al., 2016; Villarroel et al., 2016). Digestion of dietary proteins relies on cysteine and aspartyl proteases and aminopeptidases (Santamaria et al., 2015a) and that of dietary carbohydrates on glycosidases (Nisbet and Billingsley, 2000). The genome of *T. urticae* has been sequenced (Grbic et al., 2011) and specific expansions have been found for gene families involved in digestion and detoxification processes, which may be

involved in the capability of *T. urticae* to cope with the varying nutrients and defense compounds of different host plants.

T. urticae is a notorious pest responsible for yield losses in both greenhouse and open field crops (Lange and Bronson, 1981; Vacante, 2016). In tomato, it feeds mainly on the leaves causing typical chlorotic lesions (Figure 1.2D) and leaf defoliation when mite populations are high (Jayasinghe and Mallik, 2010) leading to severe crop losses. Furthermore, it can cause cosmetic damage on tomato fruit (known as gold fleck) which increases economic losses (Meck et al., 2013). *T. urticae* control is largely based on the use of acaricides (Van Leeuwen and Dermauw, 2016) and/or biological control agents (CABI datasheet, <http://www.cabi.org/isc/datasheet/53366>, accessed on December 2016). However, frequent acaricide applications have often led to development of resistance, as this species is the arthropod with the most pesticide numbered resistance reports (Van Leeuwen and Dermauw, 2016). Besides more generations per year are expected for *T. urticae* induced by global warming according to climatic and distribution models (Luedeling et al., 2011), whereas a negative effect of dry and hot conditions is expected on some of its predators (Ferrero et al., 2010), which might increase the risk of outbreaks in the future.

Figure 1.2 Adult of *Tetranychus urticae* (A) and *T. evansi* (B) under stereomicroscope (images by M. García, CIB-CSIC). Schematic section of a leaf with a spider mite feeding from mesophyll cells (light green) by introducing the stylet (red line) through the epidermic cells (blue) (C) (adapted from Bensoussan et al. (2016)) and Damage caused by Tetranychid mites on tomato leaflet (D).

***Tetranychus evansi* Baker & Pritchard (Acari: Tetranychidae)**, the tomato red spider mite, is an emerging pest of solanaceous plants worldwide. Its morphology is similar to that of *T. urticae* but the body is typically orange red in colour (Figure 1.2B). This species is native to South America and in the last two decades has expanded its geographical distribution range to southern USA, most of sub-Saharan Africa, around the Mediterranean basin and East Asia (Migeon and Dorkeld 2006-2016; Navajas et al., 2013). Two distinct genetic lineages (I and II) have been identified, the distribution of lineage I being wider outside its area of origin (Boubou et al., 2011, 2012). As a tropical species, its optimal developmental temperature is in the range 37-40 °C, with a mean generation time of about 10 days (Gotoh et al., 2010), but it can not tolerate cold winter conditions (Gotoh et al., 2010; Migeon et al., 2015). The feeding behaviour is similar to the two-spotted spider mite, but its digestive physiology is unknown. The production of a profuse web over the leaves gives this species a competitive advantage against competitors such as *T. urticae* (Sarmiento et al., 2011a) and protect them from natural enemies (Ferragut et al., 2013)

T. evansi is considered a solanaceous specialised herbivore, being reported in 53 species of the family Solanaceae, including cultivated crops such as tomato, eggplant, pepper and potato (Navajas et al., 2013). However, it has also been reported to be able to feed on plants of other families, a complete list of hosts can be found in Migeon and Dorkeld (2006-2016). Outbreaks in some African regions have been reported to cause high yield losses in tomato (Saunyama and Knapp 2003), and it has been added to the alert list of the European and Mediterranean Plant Protection Organization (EPPO, 2004). Conventional biological control has failed to be effective against *T. evansi* in the invaded areas, as naturally occurring and commercial available

Figure 1.3 Actual *Tetranychus evansi* distribution and future prospect under a conservative scenario. Adapted from Meynard et al. (2013).

phytoseiids are not able to feed on it (Navajas et al., 2013). Chemical control is also difficult because the dense webs that they produce act as barriers, and it is also prone to develop resistance (Van Leeuwen and Dermauw, 2016). Climatic and distribution models predict that the hot and dry conditions imposed by climate change will increase its potential to spread to northern latitudes in Europe (Meynard et al., 2013; Figure 1.3).

Aculops lycopersici (Masse) (Acari: Eriophyidae), the tomato russet mite, is present worldwide in areas where solanaceous crops are grown. It is the smallest arthropod feeding on tomato plants, as the adult's body is between 100 and 140 μm long (Lindquist 1996). The life cycle consists of egg, two immature stages and adult. The immature stages and adults are yellow, vermiform and with two pairs of legs (Figure 1.4A&B). Its optimal growing conditions are at 25-30 °C and low relative humidity (about 30%), with a mean generation time of about one week and females producing 30-50 eggs under these conditions, giving rise to a succession of overlapping generations (Haque and Kaway 2003; Vacante, 2016). As described for *T. urticae* unfertilized eggs (haploid) develop into male offspring whereas fertilized eggs (diploid) produce female offspring (Lindquist 1996). This species doesn't produce webs, its disperses by wind or on animal carriers (Michalska et al., 2010)

A. lycopersici is an oligophagous mite that has tomato as its primary host but it can feed on other solanaceous crops including potato, tobacco, eggplant and pepper, and some

Figure 1.4 Adult of *Aculops lycopersici* under stereomicroscope (A) (image by J. van Arkel, UvA) and under electro microscope (B) (image by E. Erbe and C. Pooley, composition by R. Ochoa, USDA-ARS). Schematic section of a mite (yellow) feeding by introducing the stylet (red) on a epidermic cell (blue) (C) (adapted from Nuzzazi and Alberty (1996)). Damage caused on tomato leaf observing shriveling on subterminal leaflets (D).

convolvulaceus crops such as sweet potato (CABI Datasheet; www.cabi.org/isc/datasheet/56111, accessed September 2016). The mouthparts are modified into a short stylet that allows them to puncture the plant epidermal cells (Figure 1.4C) but not to reach the parenchyma (Nuzzaci and Alberti 1996). Epidermis cells collapse and a callous layer is formed adjacent to the parenchyma in regions of epidermal cell death, continuous feeding causes a total destruction of the epidermis that is visible as russetting (Royalty and Perring, 1988; Duso et al., 2010). It is unknown if *A. lycopersici* injects saliva into the cells when feeding, but Glas (2014) has identified several putative “secreted salivary gland peptides” in the genome of this species.

On tomato, *A. lycopersici* feeds on foliage, stems, inflorescences and young fruits, causing shriveling and necrosis of leaves (Figure 1.4D), dropping of flowers, russetting of fruit and, if uncontrolled, death of the plant (CABI, <http://www.cabi.org/isc/datasheet/56111>) (Duso et al., 2010, Van Houten et al., 2013). As consequence, russet mite infestation can cause up to 50 % of yield losses on tomato (Duso et al., 2010), the loss being mainly due to the dead of flower buds (Kamau et al., 1992). It is a species that is difficult to control, since it’s small size allows it to hide in the forest of tomato trichomes that protects them from predators and pesticides (Van Houten et al., 2013). The lack of tomato varieties resistant to russet mites and the difficulties in timely identification of damage symptoms, since the damage is similar to that caused by broad mite and thrips, make it seriously noxious to the tomato (Duso et al., 2010). Moreover, the increase in temperatures due to climate change might favour its dispersal to new latitudes (Haque and Kaway 2003), increasing economic losses in the future.

1.3 Tomato defense response to mites

Terrestrial plants have developed a number of defense traits against herbivorous arthropods that can be classified according to the manner in which they are regulated. Constitutive defenses include physical and chemical barriers that exist irrespective of whether herbivores are present, though many such traits are also induced following herbivore attack. In contrast, induced defenses are only expressed in response to herbivory. Constitutive and induced defenses including those associated with glandular trichomes have been well documented in *Lycopersicon* spp (Kennedy, 2003; Howe and Jander, 2008). Eight different types of trichome have been described in tomato, half of them being glandular and half non-glandular. They confer resistance to small arthropods, including *T. urticae* and *T. evansi* (Simmons and Gurr, 2005), and have been used in plant breeding to increase herbivore resistance (Glas et al., 2012). However, a tiny mite like *A. lycopersici* is too small to be hindered

by tomato trichomes, which confer shelter to this mite by creating a competitor-free and natural enemy-free space (Van Houten et al., 2013). The defensive role of trichomes in varieties with low trichome density, such as Moneymaker (Glas et al., 2012), is likely to be negligible. Inducible defenses can be divided into direct defenses, which affect the growth or survival of herbivores negatively, and indirect defenses that attract natural enemies of the herbivores (Kant et al., 2015). Tomato inducible defenses include secondary metabolites and anti-nutritional proteins as direct defenses, and volatile-mediated indirect defenses (Stout et al., 1994; Kennedy, 2003; Kant et al., 2004; Martel et al., 2015). The induction of plant defenses has different steps that are followed successively to trigger the response. First plants should detect herbivory activity, then phytohormone signalling pathways will regulate the reconfiguration of plant metabolism, resulting in the induction of a wide array of plant defense traits (Howe and Jander, 2008; Schuman and Baldwin, 2016). Plant response to mites is summarized in Figure 1.5.

1.3.1 Induced defense: Phytohormone signalling

Plants recognize herbivory by perception of damage patterns and herbivore associated-elicitors (Howe and Jander, 2008; Schuman and Baldwin, 2016). Damage patterns are for instance cell wall fragments like oligogalacturonides that induce expression of defense compounds in tomato (Doares et al., 1995). Elicitors are the compounds present in the herbivore secretions (saliva, regurgitant, oviposition fluid, etc.) that stimulate plant defenses, as reported for tomato (Boughton et al., 2006). Even when both patterns are used for recognition, plants are reported to differentiate between wounding and herbivory, something in which elicitors are crucial (Howe and Jander, 2008). In the case of mites, some potential effectors have been indentified on their saliva, though the mechanism by which plants perceive their presence is unknown.

Two phytohormones are the main regulators of plant defense, Jasmonic acid (JA) and Salicylic acid (SA) (Pieterse et al., 2012; Verma et al., 2016). JA regulates the defenses against herbivorous arthropods and necrotrophic pathogens, while SA regulates the defense against phloem-feeding herbivores and biotrophic pathogens. Several other phytohormones that control other aspects of plant physiology are known to play also a secondary role in plant defense, including abscisic acid (ABA) and ethylene (ET) (Kant et al., 2015; Verma et al., 2016).

The JA-signalling and biosynthesis pathway has been widely studied, with various transgenic lines in *Arabidopsis* and tomato available for its study (Schuman and Baldwin, 2016; Singh et al., 2016). The biosynthesis of JA starts in the chloroplast, where α -linolenic acid released from its membrane is converted into 12-oxo-phytodienoic acid (OPDA) by

lipoxygenases (LOX), allene oxide synthase (AOS) and allene oxide cyclase (AOC). OPDA is transferred to the peroxisome where it will be converted by OPDA reductase (OPR3) and β -oxidation steps into JA, which is released to the cytosol to be transformed and conjugated to different forms, JA-Isoleucine (JA-Ile) being the active bioform. JA-Ile is perceived in the nucleus by the SCF^{COI1} (Skip/Cullin/F-Box) receptor complex which ubiquitinates and subsequently produces degradation of transcriptional suppressors like *JAZ*, resulting in an activation of transcription factors like *MYC2*. This has as consequence transcriptional reconfiguration of plant defense traits. The JA-pathway is activated in tomato in response to *T. urticae* (Glas, 2014; Martel et al, 2015), and *A. lycopersici* (Glas, 2014), inducing both species the accumulation of OPDA, JA and JA-Ile (Glas, 2014; Glas et al, 2014; Alba et al., 2015). Furthermore, the importance of the JA pathway for plant resistance to mites was proven by the use of JA-deficient mutants of *Arabidopsis* that show a higher susceptibility to *T. urticae* (Zhurov et al., 2014) and of tomato to *A. lycopersici* (Glas et al., 2014).

The **SA-pathway** has been mostly studied as response to biotrophic pathogen attack and phloem-feeding herbivores (Pieterse et al. 2012; Kant et al., 2015; Verma et al., 2016). SA is an aromatic compound derived from chorismate. Its biosynthesis takes place in the chloroplast by two distinct routes, one involving isochorismate synthase and the other phenylalanine ammonia lyase. The SA-pathway regulates the expression of a large set of defense-related genes, including pathogen-related proteins (PRs), and is responsible for Systemic Acquired Resistance (SAR). SA is induced in tomato by *T. urticae* (Alba et al., 2015; Glas et al., 2014a) and *A. lycopersici* (Glas, 2014; Glas et al., 2014a), but its role in plant defense response against mites is not clearly understood. SA can be converted into Methyl salicylate, a volatile induced in tomato by spider mites that plays an important role in indirect defense as an attractant of natural enemies (Kant et al., 2004). However, when the whole tomato genome transcriptional response to *T. urticae* was analyzed, none of the genes encoding SA signaling proteins, nor commonly used SA markers, were differentially expressed (Martel et al., 2015). Furthermore, SA-deficient mutants increased the performance of *T. urticae* on tomato (Villarroel et al., 2016) but had no effect on *Arabidopsis* (Zhurov et al., 2014).

ABA is a hormone that is a key regulator of abiotic stress responses in plants (see section 1.5), that also plays a secondary role in plant defense, either alone or in conjunction with the primary defense hormones (Nakashima and Yamaguchi-Shinozaki, 2013). **ET** is a hormone that regulates a large number of plant processes, including defense against necrotrophic pathogens (Kazan and Lyons, 2014). ABA is induced in tomato by *T. urticae* and *A. lycopersici* but not by *T. evansi* (Glas, 2014), whereas the signalling cascade downstream of

ET synthesis showed both up- and down-regulation in the response of tomato to *T. urticae* infestation (Martel et al., 2015). Thus, it is difficult to assess the importance of these hormones in tomato defense against mites.

Antagonistic and/or synergistic interactions between hormone signal transduction pathways can occur, modulating each other's actions (Pieterse et al., 2012; Verma et al., 2016). Crosstalk between SA and JA pathways is well documented in tomato, where the pathways have been frequently observed to antagonize each other with corresponding trade-offs in disease and pest resistance (Doherty et al., 1988; Thaler et al., 1999; El Oirdi et al., 2011). However, neutral and synergistic interactions have been described as well in tomato, including the case of a positive JA-ET interaction in response to wounding (O'Donnell et al., 1996). It has been suggested that hormone crosstalk allows plants to finely regulate plant defenses in a cost-efficient manner, depending on the type of attacker, or in case when the plant is attacked by multiple attackers, depending on the timing and sequence of infestation (Kant et al., 2015).

1.3.2 Defensive secondary metabolites and anti-nutritional proteins

The term '**secondary metabolites**' refers to those metabolites that are not directly involved in the primary biochemical pathways of an organism (growth, development and reproduction). Plants are a remarkably prolific source of structurally diverse secondary metabolites (terpenoids, alkaloids, phenolics, flavonoids, tannins, glycosides saponins, etc.), which are involved in the adaptation of plants to their environment, including direct defense responses against herbivory because of their toxic or deterrent properties (Kant et al., 2015). Secondary metabolites also include volatile compounds that are released to attract the natural enemies of herbivores, playing a role in indirect defense (War et al., 2012). In the case of tomato, different secondary metabolites (e.g., phenolics, flavonoids and terpenoids) are elicited in response to mite feeding, and there is some evidence that both JA and SA pathways are involved in orchestrating this response (Kant et al., 2004; Glas, 2014; Martel et al., 2015). Among them there are volatile terpenoids, and it has been demonstrated that their emission is positively correlated with the attraction of predatory mites (Kant et al., 2004).

Anti-nutritional proteins have as target the disruption of herbivore digestion, reducing the uptake of nutrients (Zhu-Salzman et al., 2008; Howe and Herde, 2015). Studies on the tomato–mite interaction highlight the importance of plant defense proteins that act in the herbivore's gut (e.g., amino acid-degrading enzymes, oxidative enzymes and protease

inhibitors) as prominent components of tomato-induced defenses elicited by spider mite feeding (Li et al., 2002; Kant et al., 2004; Martel et al., 2015).

Threonine desaminase (TD) catabolyzes the amino acid threonine in the herbivor's gut, depleting the availability of this essential amino acid (Chen et al., 2005, 2007). Oxidative enzymes such as **Polyphenol oxidase (PPO)** accumulates on tomato glandular trichomes (Yu et al., 1992) and generates quinones which covalently bind dietary proteins reducing the digestibility of plant tissues (Constabel and Barbenheene; 2008). **Peroxidase (POD)** is another important oxidative enzyme with various roles in plant defense: it contributes to the creation of a physical barrier by catalysing the cross-linking of cell wall compounds, it regulates reactive oxygen species (ROS) and, as PPO, it can generate quinones that interfere with digestion (Passardi et al., 2005; Zhu-Salzman et al., 2008). PPO enzymatic activity and gene expression of TD and PPO are induced in tomato by *T. urticae* and *T. evansi* (Alba et al., 2015; Martel et al. 2015) but not by *A. lycopersici* (Glas et al., 2014), whereas POD is induced by both *T. urticae* (Glas, 2014) and *A. lycopersici* (Stout et al., 1994, 1996; Glas, 2014).

Protease inhibitors (PIs) are the most studied anti-nutritional proteins against mites. The PIs are classified by the nature of the proteases that they inhibit (serine, cysteine, aspartyl, etc.) (Zhu-Salzman and Zeng, 2015). Tomato has a large number of PIs: 95 genes were found while only 38 were found on *Arabidopsis*, being the expression of 25 of them affected by *T. urticae* infestation (Martel et al., 2015). The serine (*PI-I*, *WIPI-II*, *JIP-2I*) aspartyl (*CDI*) and cysteine (*CysPI*) PI genes are induced by tetranychid mites in tomato but not by *A. lycopersici* (Li, et al., 2002; Kant et al., 2014; Glas, 2014; Glas et al., 2014; Alba et al. 2015; Martel et al., 2015). Plant cysteine PIs are harmful for *T. urticae* (Santamaria et al., 2012, 2015b), as they inhibit some of the main mite gut proteolytic enzymes (cathepsins B-, L-, and D- and legumain-like proteases) (Carrillo et al., 2011; Santamaria et al., 2015a). Plant serine PIs are also harmful to *T. urticae* (Santamaria et al., 2012), but they may target other physiological processes, since serine proteases (trypsin and chymotrypsin) do not appear to be directly involved in the hydrolysis of dietary proteins in this species (Carrillo et al., 2011; Santamaria et al., 2015a).

Some of the genes mentioned above are used as plant defense marker genes. Markers for the JA-pathway include the TD (*TD-II*), PPO (*PPO-F*, *PPO-D*) and PI (*PI-I*, *PI-II-f*, *JIP-2I*) genes. The most used SA-pathway marker genes, in turn, are the Pathogenesis Related proteins (*PR-1a* and *PR-P6*) proteins. PR proteins function against herbivores is not clear yet: they might have glucanase, chitinase, PI or peroxidase activities (Kant et al., 2015) or, like PR-

1, sterol-binding activity (Gamir et al., 2016). The timing of the tomato defense response has

Figure 1.5 Plant cell- mite interaction schematic overview. The mite introduces the stylet in the parenchymatic cell injecting saliva, the saliva content elicits that together with DAMP (Damage Associated Molecular Patterns) induce the JA- and SA- signaling pathways (in black). The mite takes the plant nutrients (green) and the defense compounds (in red). The defense compounds interfere on mite digestion. To adapt to plant defenses the mite attenuates plant defense producing effectors, and circumvents their deleterious effects by transcriptional plasticity of gene families involved in detoxification and digestion processes (blue). Abbreviation: TF = Transcription Factor; PI = Protease Inhibitors; PPO = Polyphenol Oxidases; POD = Peroxidases; TD = Threonine Deaminases; PR = Pathogen Related proteins; SAR= Systemic Acquired Resistance

been studied using some of these markers genes (Martel et al., 2015; Alba et al., 2015). The JA response is in general fast, varying between 6h and 4 days depending on the severity of mite infestation (Martel et al., 2015; Alba et al., 2015). SA related gene expression and secondary metabolite production is slower starting at 4 dpi (Kant et al., 2004).

1.4 Mite adaptation to plant defenses.

Phytophagous mites have shown a high ability to adapt to new plant species, as is proven by the oligophagous *T. evansi* and *A. lycopersici* and the extremely polyphagous *T. urticae* (Migeon and Dorkeld, 2006-2016; CABI-Datasheet; www.cabi.org, accessed September 2016). Different studies have shown that local *T. urticae* strains reared on bean or cucumber (the most common host used in laboratory culture) show an initial low performance on tomato but quickly adapt to feed on it (Fry, 1989; Gotoh et al., 1993; Agrawal et al., 2002; Magalhaes et al., 2009). In order to be able to adapt and feed on tomato, mites have developed different strategies, including physiological adaptations to tomato plant defenses by suppressing or circumventing them (Glas et al., 2014; Alba et al., 2015; Van Leeuwen and Dermauw, 2016). In addition, some of these mites have been reported to induce changes in the primary metabolism of tomato plants that might be beneficial for their performance (Wybouw et al., 2015). The mite response to plant defenses is summarized on Figure 1.5.

The **suppression of plant defenses** is, a strategy employed by plant pathogens and arthropod herbivores, characterized by lowering the rate of production of defensive compounds of the plant (Kant et al., 2015). In mites, it has been described on tomato for *T. evansi* (Sarmiento et al., 2011b; Alba et al., 2015), *A. lycopersici* (Glas et al., 2014) and some strains of *T. urticae* (Kant et al., 2008; Alba et al., 2015). Interestingly, in the process of adaptation to tomato, *T. urticae* has been shown to develop the ability to attenuate tomato plant defenses (Wybouw et al., 2015). The suppression is produced by molecules, called effectors, which can interfere with any of the steps of the plant defense signalling cascade (Kant et al., 2015). The interference, in case of mites, happens independently to the phytohormonal crosstalk, being the mechanism unknown (Glas et al., 2014; Alba et al., 2015). Recently, Villarroel et al. (2016) identified four proteins in *T. urticae* and *T. evansi* saliva which may act as effectors. Indeed, they hypothesise that in the mite saliva there is a wide range of proteins that can act as effectors or elicitors, and its presence will determine the tomato defense outcome. In fact, the proteins present in the saliva of *T. urticae* vary depending on the host (Jonckheere et al., 2016).

To **circumvent the deleterious effects** of plant secondary metabolites and anti-nutritional proteins, herbivores can decrease the sensitivity of the target sites or decrease the

exposure by increasing the metabolism of plant defenses (Kant et al., 2015). The proliferation (Grbic et al., 2011) and transcriptional plasticity of gene families involved in detoxification and digestion processes (Dermauw et al., 2013; Wybouw et al., 2015) have been proposed as one of the reasons of the adaption capacity of *T. urticae* to novel hosts. Four main detoxification enzymes families have been described: P450 monooxygenases, carboxyl/cholinesterases, glutathione-S-transferases (GST), and ATP-binding cassette transporters; which respond strongly in *T. urticae* in relation to host plant switching (Van Leuween and Dermauw, 2016). Besides, when *T. urticae* were fed with inhibitors of P-450 detoxification enzymes, mites performance was depressed severely on tomato (Agrawal et al., 2002), highlighting their importance in the adaptation to this host. Remarkably, *A. lycopersici* has only 24 P450 genes, compared with the 81 genes present in *T. urticae*, potentially reflecting host specialization (Van Leuween and Dermauw, 2016). A proliferation of cysteine protease genes (27 cathepsins B, 29 cathepsins L and 19 legumains) has also been reported in the genome of *T. urticae* (Grbic et al., 2011). This species regulates the expression of these genes in response to the ingestion of protease inhibitors to compensate for inhibitor-targeted enzymes, though they may also act as a direct barrier against other ingested plant defensive proteins (Santamaria et al., 2015a). The digestive physiology and the physiological adaptations of *T. evansi* and *A. lycopersici* to plant defenses remain largely unexplored.

Herbivore attack can also induce **changes in plant primary metabolism**, such as the reallocation of carbon and nitrogen resources (Steinbrenner et al., 2011; Zhou et al., 2015). The function of these changes for the plant is still not clear, but it has been suggested that they might be used by the plant as a source of energy, as precursors of defense compounds and to hamper herbivore performance by reducing the nutritive value of plant tissue (Zhou et al., 2015). However, the alteration of plant primary metabolism has been demonstrated to be beneficial for the performance of some herbivores, for example aphids (Sandström et al., 2000; Koyama et al., 2004; Errard et al., 2016). *T. urticae* has been reported to increase the levels of sucrose on damaged cotton tissue (Schmidt et al., 2009). In tomato, *T. urticae* infestation down regulates the transcriptional response associated with anabolic processes, including the expression of amino acid catabolizing enzyme genes (Martel et al., 2015; Wybouw et al., 2015). However, when the effects of *T. urticae* on free amino acid content in tomato were analyzed, inconsistent results were obtained (Nachappa et al., 2013; Errard et al., 2016).

1.5 Tomato response to drought stress

When facing drought stress, plants can adopt two strategies, avoidance or tolerance (Levitt 1980; Chaves and Oliveira, 2004; Verslues et al., 2006). The avoidance mechanism includes a more efficient water uptake by roots, and the capacity of plant cells to hold acquired water and further reduce water loss through reduced leaf area and limited transpiration. For drought stress tolerance, the plant activates mechanisms to withstand water deficit, like improving osmotic adjustments and increasing antioxidant metabolism. Plants often combine both strategies when responding to drought stress, a response that takes place in two phases (Chaves et al., 2003; Harb et al., 2010). The short-term response is characterised by stomatal closure and down-regulation of photochemistry. For the long-term response, morphological, physiological and metabolic changes take place. Protective responses must be triggered quickly to prevent the plant from being irreversibly damaged. Chemical signalling relies mostly on redox-active compounds and plant hormone pathways, ABA being particularly relevant (Verma et al., 2016). Summary of tomato plant response to drought on Figure 1.6.

Figure 1.6 Plant responses to drought stress in short term (left) and long term or acclimation phase (right). Down is represented the process of soil water loss as Saturation Weight percentage (SW). The abbreviation: ABA= Absciscic acid; ROS= Reactive Oxygen Species; POD= Peroxidases. (Stomata image by Jeremy Burgess, SPL, Science Source).

Stomatal closure is among the earliest plant responses to water deficit, which restricts water loss and maintains leaf turgor by modulating the transpiration level (Daszkowska-Golec and Szarejko, 2013). Stomatal conductance (gs) provides a reliable indicator of the intensity of drought stress and it has been successfully used for tomato (Thompson et al., 2007), being able to discriminate between mild, moderate and severe stress (Hummel et al., 2010). It has also been reported as an indicator of drought stress tolerance on tomato plant varieties, which showed reduced stomatal conductance (Galmes et al., 2011). By closing stomata, the pathway through which atmospheric CO₂ enters the leaf is also restricted, affecting **photosynthesis** (Chaves and Oliveira, 2004). An effective way to measure the effect of water deficit on photosynthesis is the maximum quantum yield of photosystem II photochemistry (Fv/Fm), being also an indicator of the health status of the tomato plant (Nankishore and Farrell, 2016). Its optimal value in tomato is around 0.8 (Rigano et al., 2016), but it can be reduced to values under 0.7 after long-term water shortage (Mishra et al., 2012).

Morphological changes mediated through acclimation to drought stress include reductions in growth and leaf area, changes in stomatal density and enhanced leaf thickness to increase water retention (De Micco and Aronne 2012; Clauw et al., 2015). Plants also reduce their growth by modulating both cell division and cell expansion, as ways to save and redistribute resources that can become limited under stress (Skirycz and Inze 2010; Xu et al., 2010;). As a result, a reduction in stem growth rate, length, leaf area, plant weight and biomass has been reported for tomato plants under drought stress (Thompson et al., 2007; Petrozza et al., 2014; Tapia et al., 2016).

Osmotic adjustment is a key factor in drought stress tolerance that allows plants to maintain turgor and keep normal cell activity (Bartels and Sunkar, 2005; Blum, 2017). It involves the accumulation of osmolites, mainly soluble carbohydrates and amino acids, facilitating water uptake and retention, and protecting and stabilizing macromolecules and structures from damage induced by stress conditions. The increase in free sugars and amino acids mostly results in increased starch and protein hydrolysis, respectively, which requires activity of hydrolytic enzymes (Hummel et al., 2010). Both, free sugars and amino acids accumulate on tomato leaves during restricted watering (English-Loeb et al., 1997; Tapia et al., 2016). Among the amino acids, proline is considered to be a compatible solute in many plant species, being a good indicator of drought stress on tomato plants (Claussen, 2005). Apart from acting as an osmolyte, other roles have been proposed for proline in drought stressed plants, such as being a hydroxyl radical scavenger, an energy and nitrogen source, and a signalling molecule to modulate mitochondrial functions, cell proliferation and cell death (Szabados and

Savouré, 2010). Another amino acid that has been reported to play an important role during water stress in tomato is glutamine for the relocation of leaf nitrogen (Bauer et al., 1997).

A consequence of stomatal closure in response to water deficit is an imbalance between the use and the generation of electrons in the chloroplast due to decreased availability of CO₂ for photosynthesis, leading to **overproduction of Reactive Oxygen Species (ROS)** (Cruz de Carvalho, 2008; Miller et al., 2010). This enhanced ROS production, which also occurs in mitochondria and peroxisomes, serves as a stress signal for the activation of drought stress-response (Miller et al., 2010; Verma et al., 2016), but has the potential to cause oxidative damage to plant cells (Sharma et al., 2012). To alleviate this effect, plants have evolved enzymatic and non-enzymatic systems for ROS scavenging and detoxification (Cruz de Carvalho, 2008; Noctor et al., 2014; Sharma et al., 2012). Enzymatic antioxidant activities in tomato include superoxide dismutases, catalases and peroxidases (POD) (Hayat et al., 2008; Sanchez-Rodriguez et al., 2010; Rai et al., 2013). The POD activity, whose anti-herbivore activity has already been mentioned, converts H₂O₂ into water. Its activity becomes increasingly greater the older the tomato plant becomes (Hayat et al., 2008) and in the early stages of drought stress (Rai et al., 2013). Its induction might vary depending on the tomato plant genotype (Sanchez-Rodriguez et al., 2010).

Other **proteins involved in herbivore defense** including protease inhibitors (PIs) and polyphenol oxidases (PPO) are also induced and play a role in the plant response to drought stress. Both serine and cysteine PIs are induced by drought in *Solanum dulcamara*, rice, potato and *Arabidopsis* (Kang et al., 2002; Huang et al., 2007; Zhang et al., 2008; Nguyen et al., 2016a), with not clear pattern in tomato (English-Loeb et al., 1997). They have been proposed to regulate the proteases induced by drought, delaying leaf senescence and thus increasing drought tolerance (Huang et al., 2007; Zhang et al., 2008; Diaz-Mendoza et al., 2014). In turn, PPO activity is reported to be induced by drought in tomato (English-Loeb et al., 1997; Thipayong et al., 2004), however its role in drought tolerance is not yet understood (Thipayong et al., 2004; Mayer 2006).

Hormone-mediated regulation of the plant response to water deficit is mainly dependent on abscisic acid (ABA), which in turn triggers the expression of a cascade of stress-responsive genes (Verma et al., 2016). The fluctuation in ABA levels occurs as an early response to stress with an attenuation if the drought stress continues (Harb et al., 2010). In the case of tomato, ABA has been involved in different drought tolerance mechanisms such as stomatal closure, growth reduction, metabolism reconfiguration and the increase of free sugars and amino acids (Thompson et al., 2007; Asbahi et al., 2012; Iovieno et al., 2016). Nevertheless,

SA and JA, the two main regulators of biotic stress responses, have been also reported to be involved in the plant response to drought (Daszkowska-Golec and Szarejko, 2013; Bartels and Sunkar 2005; De Ollas and Dodd, 2016). Jasmonates play an important role in the plant response to drought, interacting at different levels with the ABA signaling pathway on tomato plants (Muñoz-Espinosa et al., 2015; De Ollas and Dodd, 2016). In contrast, the role SA on plant response to drought is not clearly understood, in tomato is reported to increase drought tolerance and is involved on early signaling (Hayat et al., 2008; Muñoz-Espinosa et al., 2015).

The high sensitivity of tomato to water deficit has prompted different approaches for breeding drought-resistant crops (Hu and Xiong, 2014), including exploiting plant genetic diversity for the search of **drought tolerant/adapted varieties** (Foolad, 2007; Sinclair, 2011). Cultivated tomato germplasm has a narrow base, although it is one of the crops with the highest diversity of genetic sources, most coming from wild species from the Andean region where it originated (Bai and Lindhout, 2007; Foolad 2007). Drought tolerance has been studied on tomato wild relatives such as *S. pennelli*, *S. chilensis* and *S. peruvianum* originating from dry zones in Peru and Chile (Rigano et al., 2016; Tapia et al., 2016). Another source of diversity for drought tolerance are the landraces from the secondary diversity areas such as the semiarid Mediterranean or tropical regions (Galmes et al., 2011; Fullana-Pericas et al., 2016; Nankishore and Farrel, 2016). “Tomàtiga de Ramellet” tomatoes are a good example, they are a group of landraces from the Balearic Islands (Spain) traditionally growth under low water conditions whose adaptation to drought has been studied recently (Galmes et al., 2011, 2013). The link between traits and yield in those studies has been investigated by comparing the changes induce by water shortage on drought-adapted and non-adapted varieties, with the limitation of not being able to prove casual relationship between changes and drought adaptation (Verslues and Juenger 2011). In summary, they conclude that a more efficient use of water by an earlier stomata closure and a higher production of drought tolerance molecules such as the overproduction of antioxidative enzymes, the increase on chlorophyll, the higher accumulation of free amino acids and carbohydrates, are good indicators of drought stress tolerance.

1.6 Drought effect on the tomato-spider mite interaction

As climate change will increase the co-occurrence of abiotic (drought) and biotic (herbivory) stresses, it is crucial to understand how plants react to both stresses together, something that is not yet well understood (Nguyen et al., 2016b). As already mentioned, the response to both stresses is regulated by phytohormones, the interaction between them is crucial for the final response of the plant (Fujita et al., 2006; Nguyen et al., 2016b; Verma et al., 2016).

In tomato, a synergistic effect between ABA and JA has been reported in the response to drought stress and herbivory (Muñoz-Espinosa et al., 2015; Nguyen et al., 2016a), the transcription factor *MYC2* playing a key role in regulating the crosstalk between these two phytohormones (Kazan and Manners, 2013). In *S. dulcamara*, a species close to tomato, drought has been shown to increase the levels of JA and ABA, increasing resistance to the beet armyworm *Spodoptera exigua* (Nguyen 2016a) with a similar effect observed in tomato (English-Loeb et al., 1997). Interestingly, ABA-deficient tomato plants reduced the resistance to *S. exigua* but increased the SA linked defense response increasing the resistance to biotrophic pathogens (Thaler and Bostock, 2004). However, the interaction between biotic and abiotic stresses may not necessarily involve ABA. Thus, drought stress stimulates the resistance of tomato plants to pathogens, whereas exogenous ABA resulted in increased susceptibility (Achuo et al., 2006). It can be concluded that the effect of abiotic factors in plant-biotic interactions is complex and depends on the system studied.

Drought has been historically advocated as a key factor for herbivore outbreaks (Mattson and Haack, 1987). However, experimental studies have shown that although some phytophagous arthropods benefit from drought-stressed plants, others are adversely affected (Huberty and Denno, 2004). Two main hypotheses have been proposed to explain the effect of drought-stressed plants on herbivores. The “Plant stress hypothesis” assumes that the increase in free sugars and amino acids induced by drought stress increases the plant nutritional value, improving as consequence herbivore performance (White, 1984, 2009). The “Plant Vigor Hypothesis” in contrast, states that the reduction in plant growth and the increase in some defense compounds induced by drought stress make the plant less suitable for herbivores (Inbar et al., 2001; Cornelissen et al., 2008). The resulting performance of phytophagous arthropods on drought-stressed plants will then depend on the balance of plant nutrients and defenses, which may vary from plant to plant, depending on the plant species and genotype and in response to the intensity and the mode of the imposition of stress (Huberty and Denno, 2004; White, 2009). For instance, different tomato cultivars and different drought stress levels have positive, negative or no-effect on the aphid *Macrosiphum euphorbiae* (Rivelli et al., 2013). Nevertheless, the consequences of water stressed plants are not equivalent for different feeding guilds, since in general they have positive responses for borers, negative responses for gall-formers, and inconsistent responses for free-living species and leaf miners (Inbar et al., 2001; Huberty and Denno, 2004). Another important factor is the grade of specialization of the

herbivore, polyphagous species being expected to be more affected than specialists by drought-mediated alterations in plant defenses (Gutbrodt et al., 2011).

In the case of mites, drought stress has been reported to be positive, negative or to have no effect on their performance, depending on the mite species, the host plant and the stress level (Table 1.1).

Table 1.1 Effect of drought stress on mite performance on different plants. It can be positive (+) negative (-) or have no effect (=)

Mite	Plant	Stress level	Effect on mites	Reference
<i>T. urticae</i>	Alfalfa	Severe	-	Butler, (1995)
<i>T. urticae</i>	Alfalfa	Moderate	-	Butler, (1995)
<i>T. urticae</i>	Cotton	Severe	-	Sadras et al., (1998)
<i>T. urticae</i>	Cucumber	Severe	-	Gould, (1978)
<i>T. urticae</i>	Soybean	Moderate	-	Mellors et al., (1984)
<i>T. urticae</i>	Bean	Severe	-	English-Loeb, (1989)
<i>T. urticae</i>	Bean	Moderate	+	English-Loeb, (1990)
<i>T. urticae</i>	Corn	Moderate	+	Nansen et al., (2010)
<i>T. urticae</i>	Corn	Severe	+	Nansen et al., (2010)
<i>T. urticae</i>	Pink Delight	Moderate	+	Gillman (1999)
<i>T. urticae</i>	Peppermint	Moderate	+	Hollingsworth and Berry (1982)
<i>T. pacificus</i>	Vineyards	---	+	Stavriniades et al., (2010)
<i>T. pacificus</i>	Almond tree	Moderate	+	Youngman et al., (1988)
<i>T. pacificus</i>	Almond tree	Severe	=	Youngman et al., (1988)
<i>A. lycopersici</i>	Tomato	Moderate	+	Gispert et al., (1989)
<i>A. lycopersici</i>	Tomato	Severe	+	Gispert et al., (1989)

Drought stress triggers both the mobilization of plant nutritional compounds (Bauer et al., 1997) and the induction of plant defenses (English-Loeb et al., 1997; Inbar et al., 2001; Hayat et al 2008) in tomato. Both scenarios are then possible in the case of drought-mediated tomato-mite interactions. On one hand, a positive correlation has been reported between mite fecundity and the amino acid and sugar content on plants (Tulisalo, 1971; Dabrowski and Bielak, 1978; Wermelinger et al., 1991), supporting the “Plant stress hypothesis”. On the other hand, drought induces the accumulation of defense compounds like IPs, PPO and POD in tomato (English-Loeb et al., 1997; Inbar et al., 2001) that are detrimental for mites (Santamaria et al., 2012, 2015b), which is consistent with the “Plant Vigor Hypothesis”. The only study on tomato was performed with *A. lycopersici*, which increased its population and the damage it causes in the plant when irrigation was reduced (Gispert et al., 1989). However, no information is available on how tomato plant physiology is affected by the combined effects of both mites and drought stress and how these changes influence mite performance.

1.7 Objectives and general outline.

The general aim of this thesis is to investigate the effect of drought stress in tomato plants on the performance of three key mite pests. Whether or not mild or moderate drought stress (simulating two deficit irrigation schedules) induces physiological changes in tomato plants, and if they have an effect on the mite's behavior, development and physiology was investigated. Both water stress regimes were over the wilting point associated with severe drought stress. This information is essential to assess risks of mite outbreaks on tomato drought sensitive cultivars like Moneymaker and on drought-adapted tomato accessions under future deficit irrigation scenarios. Specifically, the horizontal objectives of this thesis are:

- To evaluate the effect of drought stress on the biology of the phytophagous mite: *Tetranychus urticae* (tomato adapted and non-adapted), *Tetranychus evansi*, and *Aculops lycopersici*.
- To measure the morphological, physiological and molecular changes induced by abiotic (drought) and biotic (mite infestation) stresses and the combination of both on tomato plants.
- To measure the physiological response (i.e. digestive proteases and detoxification enzymes) of *T. evansi* and adapted and non-adapted *T. urticae* when feeding on drought stressed tomato plants.
- To assess the plant-mediated effects of water deficit on the performance of *T. evansi* on tomato drought-adapted accessions

Chapter 2 and 3 focus on the effect of mild and moderate drought stress on the tomato-Tetranychid interaction. In Chapter 2, the performance of *T. evansi* on drought stressed tomato plant was studied, while in Chapter 3 it was analysed whether tomato adapted and non-adapted *T. urticae* strains perform equally when feeding on drought stressed plants. In both Chapters the changes in plant nutrients and defense compounds, as well as mite digestive proteases was estimated.

Chapter 4 concentrates on how both drought stress and the Eryophid mite, *A. lycopersici*, modify tomato plant metabolism increasing mite performance. In this chapter, changes in nutrients as well as the different layers of defense (phytohormones, genes and protein) were investigated. The mite's digestive physiology was not determined due to the difficulty in collecting mite samples.

Finally, in Chapter 5, the performance of *T. evansi* on four accessions of drought-adapted “Tomàtiga de Ramellet” tomatoes was studied. The differential response of these accessions to drought stress was analyzed to ascertain the differences in mite performance between accessions.

Chapter 2

DROUGHT-STRESSED TOMATO PLANTS TRIGGER BOTTOM-UP EFFECTS ON THE INVASIVE *Tetranychus evansi*

2.1 Introduction

Agricultural production faces the challenge to produce more food while constrained by a number of biotic and abiotic factors. Climate change is predicted to produce an increase in temperature and drought events in the next decades, especially in the mid-continental and Mediterranean climate areas where they are expected to be more frequent and intense (IPCC, 2013). Drought is by far the leading environmental stress in agriculture that limits the global productivity of major crops by directly reducing plant potential yield (Mishra et al., 2012), but also by indirectly influencing their interactions with biotic factors, as a consequence, playing a critical role on the world's food security.

Drought stress has been historically advocated as one key factor for herbivorous outbreaks (Mattson and Haack, 1987; English-Loeb, 1990). Yet, the relationship between arthropod outbreaks and drought is not consistent, depending on the timing, intensity and water stress phenology (Huberty and Denno, 2004) and on the feeding guild that the herbivore belongs to (Inbar et al., 2001). It is widely accepted that drought stress triggers significant alterations in plant biochemistry and metabolism (Hummel et al., 2010) that may alter the physiology of the host plant and modify the nutritional values, affecting herbivore performance (Han et al., 2014). There are several hypotheses concerning the response of the plant to drought stress and how herbivores adapt to those changes (Huberty and Denno, 2004; Cornelissen et al., 2008; White, 2009). Drought induces metabolic changes in the plant, such as increased levels of free sugars and free essential amino acids, which according to the “Plant stress hypothesis” causes the plant to have a higher nutritional value for herbivores (White, 1984; Inbar et al., 2001; White, 2009), and can play an important role in herbivore outbreaks (Guo et al., 2013; Johnson et al., 2014). In contrast, drought is also associated with a reduction in growth and an increase in defense compounds making the plant less suitable for herbivores according to the “Plant Vigor Hypothesis” (Cornelissen et al., 2008). The resulting performance of phytophagous arthropods on drought-stressed plants will then depends on the access they have to an optimal balance of nutrients in the plant according to their feeding habit (Huberty and Denno, 2004), and their adaptation to plant defense compounds according to their grade of specialization (Gutbrodt, 2011).

Climate change is expected to increase the incidence of water shortage in semi-arid environments. Then, deficit irrigation scheduling, yielding mild and moderate drought, might help to improve the efficiency with which water is used in major crops, such as tomato, widely cultivated in semi-arid regions. The tomato agro-ecosystem is threatened by a few major key

pests, such as spider mites, and many minor or secondary pests (Lange and Bronson, 1981). The red tomato spider mite, *Tetranychus evansi* Baker & Pritchard was first recorded in Brazil, and has emerged as a serious invasive agricultural pest in invaded areas such as Africa and Europe (Meynard et al., 2013). In last decade, it has been considered one of the most important pests of solanaceous crops in Africa, causing high yield losses in tomato in some African regions (Saunyama and Knapp, 2003). This species has been reported as highly tolerant to hot and dry conditions. As a result of climate change, the Mediterranean basin is the most threatened area for the potential spread of *T. evansi* (Navajas et al., 2013). In fact, outbreaks have been recorded in Europe, particularly around the Mediterranean basin where *T. evansi* has spread significantly in the last decades (Navajas et al., 2013). The high invasive potential of *T. evansi* and the severity of damage have prompted its addition to the alert list of the European and Mediterranean Plant Protection Organization (EPPO, 2004).

When feeding on tomato leaves, *T. evansi* was found to suppress anti-mite plant defenses by down-regulating the expression of genes involved in the regulation of secondary metabolites and defense proteins, increasing the fitness of these mites in the absence of competitors (Sarmiento et al., 2011; Alba et al, 2015). However, water deficit stress in tomato plants had been reported to increase some of these plant defenses, such as protease inhibitors and the oxidative enzymes polyphenol oxidases and peroxidases (English-Loeb et al., 1997; Inbar et al., 2001; Ripoll et al., 2014). On the other hand, water deficit stress can elicit accumulations of nutrients in tomato plants. Among such nutrients, free amino acids, which are known to be instrumental in the maintenance of host plant osmotic balance, and free sugars are usually detected at elevated levels, and can therefore contribute to improve mite performance (Showler, 2013). Moreover, free proline, a non-essential amino acid for arthropods that accumulates in most drought stressed plants, has been reported to be a feeding stimulant for many phytophagous species (Mattson and Hack, 1987). However, there is little information on the combined effects of both biotic (mites) and abiotic (drought) stress on the tomato plant physiology and on how these changes influence *T. evansi* performance.

This study represents an attempt to establish if mild or moderate drought stress (simulating two scenarios of deficit irrigation) induces physiological changes (i.e. nutritional values and chemical defenses) in tomato plants, and if they have an effect on *T. evansi* behavior, development and physiology. This holistic approach provides insight into the bottom-up effects that may influence mite performance on drought-stressed tomato plants and the implications

for outbreak risks of *T. evansi*, which might strongly affect tomato production on area-wide scales.

2.2 Materials and Methods

2.2.1 Plant material and mite rearing

A colony of *T. evansi* collected in Beausoleil (South of France) was provided by Dr. Maria Navajas (CBGP, France). Mites were maintained on detached tomato leaves (*Solanum lycopersicum* L., cv. Moneymaker) placed in ventilated plastic cages (22x30x15 cm) for about 30 generations. The petiole of the leaves was in contact with a thin layer of water in the bottom of the cages to avoid mites escaping and to maintain leaf turgor. Tomato plants were grown from seeds in 40 well trays. Plants with 3 expanded leaves were transferred to 2.5 liter pots (diameter: 16 cm, height: 15 cm) (Maceflor, Valencia, Spain) filled with 600 g of Universal growing medium 'Compo sana' (Compo GmbH, Münster, Germany) and watered to saturation. The mite colony and the tomato plants were maintained on a climate room at 25°C±1°C, 50±5% relative humidity and a 16 h light/8 h dark photoperiod.

2.2.2 Drought stress regime and experimental design

Water stress status of single tomato plants was quantified gravimetrically by frequent recording of pot weights (balance BSH 6000, PCE Iberica, Tobarra, Spain), and transpired water was replaced by intermittently watering them such that soil water content was maintained within a narrow range (Figure 2.1). Measurements of soil water status was determined by using two parameters: percentage of saturation weight [$SW = 1 - ((\text{weight of 100\% water saturated soil} - \text{soil weight}) / \text{weight of 100\% water saturated soil})$]; and the volumetric soil water content [$\Theta = (\text{soil weight} - \text{soil dry weight}) / (\text{H}_2\text{O density} * \text{soil volume})$]; soil dry weight was determined by drying 600 g of soil in a stove at 70°C for 3 days]. Tomato plants with 4–5 expanded leaves were watered every 2–3 days to maintain the SW in the range of 80–95% (corresponding to Θ between 55–74%) for the control plants. We imposed two water stress regimes, defined as mild and moderate, by stopping irrigation for 4 or 7 days, respectively, and thereafter watering every 2–3 days to maintain the SW in the range of 60–73% ($\Theta = 36$ –50%) for mild stress and between 45–53% ($\Theta = 16$ –36%) for the moderate stress (Figure 2.1). Both water stress regimes were above the wilting point associated with severe drought stress (Fitter and Hay, 2002) that was established at 40% of SW for Moneymaker in our experimental conditions.

Figure 2.1 Water stress status (well-watered, mild and moderate drought) of tomato plant. Drought induction was started 7 days before mite infestation and was continued until the end of the experiment [10 days post infestation (dpi)]. It is expressed as percentages of saturation weight (SW) and volumetric soil water content (Θ). The turgor loss point was established as 40% of SW.

Mild and moderate stressed plants were coupled with well-watered (control) plants in two independent experiments. In each experiment, plants were divided into eight different groups combining two different treatments: Drought stress (control or drought) and *T. evansi* infestation (infested or non-infested); and two sampling times (4 or 10 days post infestation = dpi). Spider mite performance and leaf damage were assessed in both mild and moderate drought stress experiments. Chemical and biochemical analysis of plant material was only performed for moderate drought stress (the utmost applied).

Tomato plants (6–7 expanded leaves) were infested when steady stress conditions were reached (Figure 2.1) by placing 8 spider mite females on each of the two leaflets next to the apical one of the leaves 3, 4 and 5 (48 mites per plant). Mites were collected from the laboratory colony using a vacuum pump D-95 (Dinko S.A., Barcelona, Spain) with a sucking power of 10–50 mmHg connected to a modified Eppendorf. All plants were confined with a ventilated metacrylate cylinder fitting the pot diameter and set up in a climate chamber in a complete randomized block design. Temperature and humidity inside the cylinders were recorded by introducing USB dataloggers Log 32 (Dostmann electronic GmbH, Wertheim, Germany), on average the relative humidity was $76 \pm 2\%$ inside the well-watered cylinders, and $73 \pm 2\%$, and

56±2% for mild and moderate drought conditions, respectively. The average temperature was 24±1°C in all cases.

The stress intensity of all plants at infestation and at each sampling time (4 or 10 dpi) was assessed by measuring: a) stomatal conductance (gs) and b) variations in maximum quantum yield of photosystem II photochemistry (Fv/Fm), as they are reported to be good drought indicators (Verslues et al., 2006; Mishra et al., 2012). Between sampling times, the stress status was not recorded to avoid mite disturbance. Stomatal conductance was measured using a leaf porometer (SC-1 Decagon-T, Pullman, USA) and the Fv/Fm was estimated using a FluorPen FP 100 (PSI, Drasov, Czech Republic). Both parameters were measured on the 4th leaf, as preliminary experiments showed that it is representative of the plant status. Plant growth was estimated by measuring the stem length (distance between the soil and the terminal bud) and by weighting the aerial part of the plant (transformed to dry weight by using the water content data calculated as referred to below).

2.2.3 Performance of *T. evansi* and leaf damage

The performance of *T. evansi* was determined by estimating mite population growth at 4 and 10 dpi. At each time, 15 infested plants per treatment were analyzed in the moderate stress experiment, and 9 infested plants per treatment in the mild stress one. All leaves were detached from the plant, and the number of eggs and mobile mite stages (larvae, nymphs and adults) were counted under a stereomicroscope M125 (Leica Mycrosystem, Wetzlar, Germany). To determine the leaf damage area (mm² of chlorotic lesions), damaged leaflets were scanned using hp scanjet (HP Scanjet 5590 Digital Flatbed Scanner series, USA). The scanned leaflets were analyzed by the program GIMP 2.8 (www.gimp.org), for which we created a new layer, and the damaged area was painted in black color, then the number of pixels in black was counted using the histogram-tool. Finally, the number of pixels was transformed to mm² using a scanned ruler as a reference.

2.2.4 Chemical and biochemical analysis of tomato plant material

Leaves from control and drought stress plants were collected and analyzed (6 plants/treatment). The collection was as follows: a) the left leaflets from leaves 3, 4 and 5 were collected together and frozen at -72°C; ground using a mortar and pestle in the presence of liquid nitrogen to a fine powder and stored for free amino acids, protein, protease inhibitors and oxidative enzymes analysis; and b) the right leaflets from the same leaves were collected together, dried in an oven at 70°C for 3 days, weighed before and after drying to assess the

percentage of water and ground using a mortar and pestle to obtain a fine powder and stored for C, N and free sugar analysis. Unless otherwise specified, all chemical compounds used here and in the mite enzymatic activities assay were from Sigma-Aldrich (St Luis, USA).

Total C and N composition. Samples of 1 mg of dried leaf powder were analyzed to determine total nitrogen and carbon concentration at the Elemental Microelement Center of Complutense University (Madrid, Spain) by using a microelement analyzer LECO CHNS-932 (LECO, St Joseph, MI, USA).

Free sugars. Samples of 3 mg of dried leaf powder were homogenized in 650 μ l of ethanol 95% (v/v), heated at 80°C for 20 min, centrifuged at 10000 rpm for 10 min, and the supernatant collected. The process was repeated two more times and the three supernatants were pooled. A volume of 750 μ l of the mixture was dried on a SpeedVac Concentrator Savant SVC-100H (ThermoFisher scientific, Willmington, DE, USA) and redissolved in 500 μ l of water. Soluble carbohydrate concentration was estimated by the anthrone method (Maness, 2010) using glucose as standard. In brief, 1 ml of anthrone reagent (0.2% v/v anthrone on 95% sulfuric acid) was added to the extract, heated at 90°C for 15 min, and the absorbance at 630 nm was measured using a VERSAmax microplate reader (Molecular Devices Corp., Sunnyvale, USA).

Free amino acids. The extraction of the free amino acids was done as described by (Hacham et al., 2002). Samples of 50 mg of leaf frozen powder were homogenized with 600 μ l of water:chloroform: methanol (3:5:12 v/v/v). After centrifugation at 12000 rpm for 2 min, the supernatant was collected and the residue was re-extracted with 600 μ l of the same mixture, pooling the two supernatants. A mixture of 300 μ l of chloroform and 450 μ l of water was added to the supernatants, and after centrifugation the upper water:methanol phase was collected and dried in a SpeedVac. The samples were dissolved on 100 μ l of sodium citrate loading buffer pH 2.2 (Biochrom, USA) and 10 μ l were injected on a Biochrom 30 Amino Acid Analyser (Biochrom, USA) at the Protein Chemistry Service at CIB (CSIC, Madrid, Spain).

Protein extracts. Samples of 100 mg of leaf frozen powder were homogenized in 500 μ l of 0.15 M NaCl, ground with fine sand (Sigma, USA) in 1.5 ml tubes, the supernatant was frozen and stored at -20°C. The homogenate was centrifuged at 12000 rpm for 5 min at 4°C, and the total protein concentration was quantified using a Nanodrop 1000 spectrophotometer (ThermoFisher scientific, Willmington, USA) with an absorbance ratio of A260/A280.

Protease inhibitors. The inhibitory activity of plant protein extracts was tested *in vitro* against commercial enzymes: papain (EC 3.4.22.2), cathepsin B from bovine spleen (EC

3.4.22.1), trypsin from bovine pancreas (EC 3.4.21.4) and α -chymotrypsin from bovine pancreas (EC 3.4.21.1). As substrates, Z-FR-AMC (N-carbobenzoxyloxy-Phe-Arg-7-amido 4-methylcoumarin) was used for papain, Z-RR-AMC (N-carbobenzoxyloxy-Arg-Arg-7-amido-4-methylcoumarin) for cathepsin B, Z-LA-AMC (Z-L-Arg-7-amido-4-methylcoumarin) for trypsin and SucAAPF-AMC (Suc-Ala-Ala-Pro-Phe-7-amido-4-methylcoumarin) for chymotrypsin, all of them purchased from Calbiochem (MerkMilipore, Billerica, USA). Samples of 20 μ g of plant protein extracts were preincubated for 10 min with 100 ng of the commercial enzyme. The incubation was carried out at 28°C in 100 mM sodium phosphate buffer pH 6.0 (10 mM L-cysteine, 10 mM EDTA, 0.01% (v/v) Brij 35) for the papain and cathepsin B assays and at 35°C in 0.1 M Tris-HCl buffer, pH 7.5, for the trypsin and chymotrypsin assays. Subsequently, the substrates were added at a final concentration of 0.2 M and incubated for 1 h at 28°C or 35°C depending on the enzyme. Fluorescence was measured using an excitation filter of 350 nm and an emission filter of 465 nm (Tecan GeniusPro, Mannedorf, Switzerland). Double blanks were used to account for spontaneous breakdown of substrates and the plant protease activity, and all assays were done in duplicate. Results were expressed as a percentage of protease activity inhibited.

Oxidative enzymes. Samples of 100 mg of leaf frozen powder were ground in a 1.5 ml tube with sand and 500 μ l of extraction buffer (0.1 M phosphate buffer, pH 7.0; 5% w:v polyvinylpolypyrrolidone). The homogenate was centrifuged at 12000 rpm for 10 min and the supernatant was collected and frozen until used. Polyphenol oxidase (PPO) activity was analyzed by incubating 20 μ l of enzyme extract with catechol (40 mM final concentration) in 160 μ l of TrisHCl pH 8.5 buffer at 30°C for 1 h. Absorbance was read at 420 nm. Peroxidase (POD) activity was determined incubating 20 μ l of a 1:10 dilution of the enzyme extract with guaiacol (5 mM final concentration) and H₂O₂ (2.5 mM final concentration) in 150 μ l of potassium phosphate pH 6 buffer at 30°C for 10 min. Absorbance was read at 470 nm. In both cases, a Varioskan Flash reader (ThermoFisher Scientific, Willmington, USA) was used.

Mite enzymatic activities. To analyze the effect of tomato drought-stressed plants on mite enzymatic activities an additional experiment was performed. Control and drought-stressed plants (6–7 leaves) were inoculated with 200 mites using the pump system previously described (6 replicates per treatment). Mites were collected at 10 dpi and frozen in liquid nitrogen. Mite protein extracts were obtained by homogenizing 1 mg of mites in 100 μ l of 0.15M NaCl, centrifuged at 12000 rpm for 5 min and collecting the supernatant. Total protein content was determined according to the method of Bradford (1976).

Cathepsin B-, cathepsin L- and legumain-like activities were assayed as described by (Santamaria et al., 2015a), using Z-RR-AMC, Z-FR-AMC (Calbiochem, USA) and Z-VAN-AMC (N-carbobenzoxycarboxyVal-Ala-Asn-7-amido-4-methylcoumarin) (Bachem, Bubendorf, Switzerland) as substrates, respectively. The assays were performed by incubating 10 μ l of mite protein extracts with 85 μ l of 100 mM sodium citrate buffer (0.15 M NaCl, 1 mM DTT, pH 5.5 for cathepsin L- and B-like activities and pH 4.5 for legumain-like assay, containing 0.2 M of substrate) at 30°C for 15 min. Fluorescence was measured using an excitation filter of 350 nm and an emission filter of 465 nm on a Varioskan Flash reader (ThermoFisher Scientific, Willington, USA). A calibration curve was obtained with known amounts of AMC (7-amino-4-methylcoumarin) (Bachem, Switzerland).

Cathepsin D-like activity was determined using MocAc-GKPILFFRLK(Dnp)-D-R NH₂ (PeptaNova GmbH, Germany) as substrate. The assays were performed incubating 5 μ l of the mite protein extract with 85 μ l of 100 mM sodium citrate buffer (0.15 M NaCl, 1 mM DTT, 10 μ M E-64, pH 3.5, containing 20 μ M substrate) at 30°C for 15 min. Fluorescence was measured using an excitation filter of 328 nm and an emission filter of 393 nm on a VarioskanFlash reader (ThermoFisher Scientific, Willington, USA), and MCA (MoCAC-Pro-Leu-Gly) (Peptanova GmbH, Germany) was used as standard.

Leucine aminopeptidase-like activity was determined using LpNa (L-leucine p-nitroanilide) as substrate. Samples of 10 μ l of mite extracts were incubated with 160 μ l of 1 mM LpNa in 0.1 M Tris-HCl buffer (0.15 M NaCl, 5 mM MgCl₂, pH 7.5) at 30°C for 4 h. The reaction was stopped with 100 μ l of 30% acetic acid and the absorbance was measured at 410 nm, using a molar extinction coefficient of 8800 M⁻¹ cm⁻¹ for LpNa.

α -Amylase activity was determined using starch as a substrate as described by Valencia et al. (2000). Samples of 20 μ l of mite extract were incubated with 80 μ l of 0.1 M Tris-HCl buffer (40 mM CaCl₂, 20 mM NaCl, pH 6.0) and 100 μ l of 0.5% starch solution, at 30°C for 4 h. The reaction was stopped by adding 1 ml of lugol solution (0.02% I₂; 0.2% KI). The mixture was centrifuged at 6000 rpm for 5 min and the supernatant's absorbance was measured at 580 nm. A starch standard curve was used as a reference.

Esterase activity was determined using 1-naphthyl acetate (1-NA) as a substrate. Samples of 10 μ l of a dilution 1/40 of the mite protein extract were incubated with 160 μ l of 0.25 mM 1-NA in 0.1 M Tris-HCl buffer (0.15 M NaCl, 5 mM MgCl₂ pH 7.0) for 1 h at 30°C. The reaction was stopped by adding 80 μ l of a solution (0.04% (w/v) of Fast blue salt BN

(tetrazotized) and 1.5% (w/v) SDS) and incubating at room temperature for 1 h. A standard curve of 1-naphthol was used as a reference. Absorbance was measured at 600 nm.

Glutathione S-transferase (GST) activity was determined incubating 40 μ l of mite protein extract with 220 μ l of 0.1 M Tris-HCl buffer (pH 8.0), containing 0.4 mM CDNB (1-chloro- 2,4-dinitrobenzene) as substrate and 5 mM reduced glutathione as cofactor for 15 min at 30°C. The increment in absorbance at 340 nm was recorded every minute for 5 min, using a molar extinction coefficient of 9.6 mM⁻¹ cm⁻¹ for conjugated CDNB.

P450 activity was assayed by incubating 60 μ l of mite protein extract with 120 μ l of 100 mM Tris-HCl buffer, (pH 7.0; containing the NADPH generating system [0.5 mM NADP, 2.5 mM glucose 6-phosphate and 0.3 units of glucose 6-phosphate dehydrogenase]) and 20 μ l of a 50 μ M cytochrome c solution at 30°C for 4 h. The reaction was stopped with 100 μ l of methanol and the absorbance measured at 550 nm, using a molar extinction coefficient of 27.6 mM⁻¹ cm⁻¹ for cytochrome c reduced.

Hydrolytic activities were expressed as nmol of substrate hydrolyzed per min and mg of mite protein, GST activity as nmol of CDNB conjugated per min and mg of protein and P450 activity as nmol of cytochrome c reduced per min and mg of protein.

2.2.5 Proline feeding stimulant test

Assays were performed in plastic Petri dishes (30x90 mm), coated in their bottom half with about 30 ml of a 0.625% agar solution to prevent leaf desiccation. The Petri dishes had a hole in the top, and were covered with filter paper to allow ventilation and to prevent the escape of mites. Tomato leaf disks (3.14 cm²) were cut from well-watered tomato plants with a corkborer ($\varnothing$: 2 cm) and weighed. The abaxial and the adaxial surfaces of the leaf disks were dispensed with 25 μ l of an ethanol solution of L-proline (Sigma, St Louis, USA) or ethanol alone for the control, and allowed to dry before mite inoculation. Concentrations were chosen to simulate the L-proline induced by both drought and *T. evansi* infestation on tomato plants at 4 and 10 dpi (see Section 2.4), corresponding to 0.070 mg and 0.202 mg of L-proline/g of fresh leaf weight, respectively. Twenty *T. evansi* adult females from the colony, replicated ten times, were placed on each leaf disk (one per plate), being the number of eggs laid and the leaf damaged area recorded after 48 h. All Petri dishes were kept in a growth chamber (Sanyo MLR-350-H, Sanyo, Japan) at 25°C $\pm$ 1°C, 70 $\pm$ 5% relative humidity and a 16 h light/8 h dark photoperiod.

2.2.6 Statistical analysis

All plant and mite parameters analyzed were checked for the assumptions of normality and heteroscedasticity, and transformed if necessary. Stem length and stomatal conductance were $\log_{10}(x)$ transformed, Fv/Fm was $\log_{10}(x+1)$ transformed; and these two parameters and stem dry weight data were statistically analyzed using a three-way ANOVA (using as fixed factors drought treatment, *T. evansi* infestation and time). The number of *T. evansi* eggs and mobile forms and leaf damage area were $\log_{10}(x)$ transformed and analyzed by a two way ANOVA (drought treatment, time). For stem length, plant aerial dry weight, stomatal conductance, Fv/Fm, mite population and leaf damage a Bonferroni *post hoc* test was performed to compare drought-stress treatments within each time. Percentage of nitrogen, protein, free amino acids and free sugars, as well as C:N ratio and protease inhibition were $\arcsin(\text{squareroot}(x))$ transformed. These data and the oxidative enzyme activities were analyzed using a two-way ANOVA for each time separately using as fixed factors drought treatment and *T. evansi* infestation. A Newman-Keuls *post hoc* test was performed to see differences between treatments mean. A Dunnet *post hoc* test was performed for every single amino acid data at 10 dpi, using as reference the control non-infested plant. Spider mite enzymatic activity data were analysed by a t-student test. For the proline feeding assay, data were $\log_{10}(x)$ transformed and analysed by one-way ANOVA using proline concentration as a factor. A Bonferroni *post hoc* test was performed to determine the difference between treatments.

2.3 Results

2.3.1 Effects of drought on tomato plant growth, stomatal conductance and photosynthetic efficiency

Data from both infested and non-infested plants were pooled as no significant differences were found between them (data not shown). Moderate drought stress had a strong effect on stomatal conductance, which was significantly reduced between 4 and 5 times with respect to the control plants at 4 and 10 dpi (Figure 2.2a). The Fv/Fm of moderate drought stress plants was also significantly smaller than the control at infestation and at 10 dpi (Figure 2.2b). Similar effects, but less pronounced were obtained with mild stressed plants, with the stomatal conductance progressively reduced from about 20% less than in control plants at mite infestation to about half at 10 dpi, whereas Fv/Fm was only significantly lower at 10 dpi (Figure S2.1). Significant differences in the growing pattern (aerial dry weight and/or stem length) were

obtained at 10 dpi for both drought levels (Figure 2.2c, 2.2d and S2.1), with the reductions more acute for moderate stressed plants.

Figure 2.2 Effect of moderate drought at mite infestation and at 4 and 10 days post infestation (dpi) on: a) tomato stomatal conductance (g_s); b) maximum quantum yield of PSII photochemistry (F_v/F_m); c) stem length; d) plant aerial dry weight. Data are mean $\pm$ SE. * indicates a statistically significant difference within each time (Three-way ANOVA, Bonferroni *post hoc* test, $P < 0.05$).

2.3.2 Effect of drought on *T. evansi* population growth and leaf damage

The population of *T. evansi* increased faster when feeding on both moderate and mild drought-stressed plants than on control tomato plants (Figure 2.3). The number of eggs laid under moderate stress was significantly higher at 4 (2 fold) and 10 dpi (1.6 fold) than on control plants (Figure 2.3a). No differences on mobile forms were observed at 4 dpi, as these individuals correspond to the surviving adult females (F0) that were used to infest the plant

Figure 2.3 Performance of *T. evansi* in the moderate (a,b&c) and mild (d,e,f) drought stress experiments. The number of eggs (a&d), total mobile forms (b&e) and leaf damaged area (c&f) on control and drought stressed (mild or moderate) tomato plants at 4 and 10 days post infestation (dpi) were measured. Data are mean $\pm$ SE. * indicates a statistically significant difference within each time (Two-way ANOVA, Bonferroni *post hoc* test, $P < 0.05$).

(Figure 2.3b). However, a significant increase (2.1 fold) was observed at 10 dpi, with most of these individuals resulting from the F1 progeny. Interestingly, the leaf damage area was 1.3 and 1.6 times significantly larger than on controls at both 4 dpi and 10 dpi, respectively (Figure 2.3c). When mites were reared on mild drought-stressed tomato plants, significant increases in eggs laid (Figure 2.3d), mobile forms (Figure 2.3e) and leaf damage area (Figure 2.3f) were also recorded.

2.3.3 Effects of drought and *T. evansi* on plant nutritional composition and defense proteins

Differences in the levels of nutrients in the plants were found at 4 and 10 dpi, with drought stress the most significant factor (Table 2.1). At 4 dpi, drought, mite infestation and the combination of both factors produced a significant decrease in nitrogen and an increase of free sugars. Total protein was also reduced by drought at 4 dpi, but the difference was only significant in infested plants. At 10 dpi, a decrease in nitrogen and total protein, and an increase in free sugars were obtained with drought alone or in combination with mite infestation. Drought was also a significant factor for total free amino acids at 10 dpi, although no significant differences among treatments were found (Table 2.1). However, it could be demonstrated that drought drives the accumulation of specific amino acids (Figure 2.4). Proline was the amino

Table 2.1 Effect of moderate drought and *T. evansi* infestation on key nutrients in tomato leaves at 4 and 10 days post infestation.

	Non-infested		Infested		ANOVA (p<0.05)
	Control	Drought	Control	Drought	
4 Days Post Infestation					
Nitrogen ¹	6.7±0.1 a	5.7±0.2 b	5.8±0.2 b	5.7±0.1 b	D;I; D*I
Protein ¹	31±2 ab	26±3 b	35±2 a	25±2 b	D
Free aa ¹	1.0±0.2	1.2±0.2	1.0±0.1	1.1±0.2	-
C:N	5.9±0.1	6.4±0.2	6.5±0.3	6.3±0.2	-
Free Sugars ¹	3.3±0.1 b	3.9±0.3 a	4.2±0.1 a	4.3±0.2 a	I
10 Days Post Infestation					
Nitrogen ¹	6.3±0.2 a	5.7±0.1 b	5.9±0.2 ab	5.3±0.1 b	D
Protein ¹	40±5 a	19±4 b	27±6 ab	15±2 b	D
Total free aa ¹	0.9±0.2	1.7±0.3	1.2±0.3	1.8±0.4	D
C:N	6.3±0.1	6.4±0.1	6.1±0.2	6.6±0.2	-
Free Sugars ¹	3.2±0.2 c	4.8±0.3 a	3.7±0.1 cb	4.3±0.3 ab	D;D*I

(1) Data, as % of dry weight, are mean ± SE. D (drought) and I (mite infestation) indicate significant factors in the Two-Way ANOVA. Different lower case letters within rows indicates significant differences (Newman-Keuls test at $p < 0.05$)

acid more clearly induced by drought, increasing significantly by > 20 fold by drought alone or in combination with mite infestation (Figure 2.4). The levels of several essential (valine, isoleucine, leucine, tyrosine, histidine) and non-essential (glutamine) amino acids were also significantly induced by drought in combination with mites (Figure 2.4). The infestation by *T. evansi* did not have significant effects on any of the amino acids analyzed at 10 dpi.

Figure 2.4 Levels of free amino acids in tomato leaves subjected to moderate drought stress and *T. evansi* infestation at 10 days post infestation (dpi). Data represent the ratio $\pm$ SE of the different treatments with respect to non-infested control plants at 10 dpi. = [(Treatment – Control 10dpi) / Control 10dpi]. The division between essential and nonessential amino acids is based on a study with the closely related species *Tetranychus urticae* (Rodriguez and Hampton, 1966). * Indicates significant difference of the treatment with the control determined by Dunnet *post hoc* test $p < 0.05$.

Tomato plant defense proteins were also affected by drought stress and mite infestation, but different responses were obtained depending on the post-infestation time (Table 2.2). At 4 dpi, the inhibitory activity of tomato leaf extracts against papain was reduced by mite infestation under both drought and control conditions, whereas the inhibitory activity against chymotrypsin was reduced by drought conditions on non-infested plants. By contrast, the inhibitory activity

Table 2.2 Effect of moderate drought and *T. evansi* infestation on plant defense proteins in tomato leaves at 4 and 10 days post infestation

	Non-infested		Infested		ANOVA (p<0.05)
	Control	Drought	Control	Drought	
4 Days Post Infestation					
Protease inhibitors (% Inhibition)					
Papain	48±2 ab	64±2 a	31±5 c	45±5 b	D;I
Cathepsin B	49±3	62±6	63±8	66±4	-
Trypsin	40±8	44±2	49±8	43±2	-
Chymotrypsin	61±7 a	45±6 b	64±2 a	63±4 a	I
Oxidative enzymes (specific activity)					
Polyphenol oxidases ¹	1.1±0.1 b	2.1±0.1 a	2.5±0.2 a	2.6±0.3 a	D;I;D*I
Peroxidases ²	2.7±0.2 b	5±0.5 a	6.8±1.1 a	5.1±0.2 a	I; D*I
10 Days Post Infestation					
Protease inhibitors (%Inhibition)					
Papain	57±3 c	81±3 a	71±4 b	86±3 a	D;I
Cathepsin B	40±5 c	60±7 b	63±6 b	83±5 a	D;I
Trypsin	37±3 ab	54±4 a	31±8 b	52±5 a	D
Chymotrypsin	69±4	65±6	74±6	77±4	-
Oxidative enzymes (specific activity)					
Polyphenol oxidases ¹	2.2±0.1 b	2.5±0.4 b	3.7±0.5 a	3±0.4 ab	I
Peroxidases ²	4.3±0.5 b	4±0.5 b	8.1±0.9 a	5.8±0.6 b	D; I

(1) PPO: nmol hydrolyzed Cathecol/ mg Protein*min (2) POD: nmol hydrolyzed Guaiacol/ mg Protein*min.
Data are mean ± SE. D (drought) and I (mite infestation) indicate significant factors in the Two-Way ANOVA.
Different lower case letters within rows indicates significant differences (Newman-Keuls test at *p*<0.05).

of tomato leaf extracts at 10 dpi was enhanced against papain and cathepsin B as a response to both drought and mite infestation, whereas the inhibitory activity against trypsin increased in response to drought. The specific activity of the oxidative enzymes PPO and POD was significantly increased in response to mite infestation, drought stress and a combination of both at 4 dpi, but only in response to mite infestation at 10 dpi

2.3.4 Enzymatic activities of *T. evansi* fed on drought-stressed and well-watered tomato plants

Hydrolytic and detoxification enzyme activities in *T. evansi* were analyzed after feeding for 10 days on moderate drought-stressed or control tomato plants (Table 2.3). A significant increase in total mite protein and a reduction in the specific activity of cathepsin D- and legumain-like proteases and of α -amylase were observed in mites fed on tomato plants exposed to drought stress. The activity of the other digestive proteases (cathepsin B- and L-like and

aminopeptidase) and detoxification enzymes (esterase, GST and P450) analysed were not affected by the drought status of the plant on which they fed.

Table 2.3 Enzymatic activities in *T. evansi* feeding on moderate drought-stressed or control tomato plants for 10 days

	Control	Drought-stressed
Protein¹	55±2	69±5*
Digestive enzymes (specific activity)		
Cathepsin B²	7.8±1.0	8.7±1.1
Cathepsin L²	1.9±0.2	1.8±0.2
Cathepsin D²	13±2	5.5±0.4*
Legumain²	0.29±0.03	0.20±0.02*
Aminopeptidase²	23±3	24±3
α-Amylase²	162±7	130±8*
Detoxification enzymes		
Esterase²	1774±146	1397±216
GST³	197±30	179±17
P450⁴	0.2±0.02	0.2±0.01

Data are mean ± SE. * indicates statistically significant difference between treatments (t-student, $p < 0.05$).

(1) $\mu\text{g}/\text{mg}$ fresh weight

(2) nmol substrate hydrolyzed per min and mg of protein

(3) nmol CDNB conjugated per min and mg of protein

(4) nmol cytochrome c reduced per min and mg of protein.

Fig 2.5 Performance of *T. evansi* with regard to the number of eggs laid (a) and leaf damaged area (b), on leaf disks treated with L-proline. Concentrations applied are equivalent to that estimated on drought-stressed tomato plants at 4 and 10 days post infestation. Data are mean ± SE. Different letters indicate significant differences (One-way ANOVA, Bonferroni post hoc test, $P < 0.05$).

2.3.5 Proline feeding stimulant assay

The proline feeding stimulant assay showed a significant increase in eggs laid (2.5 fold) and leaf damage area (1.8 fold) on leaf disk treated with L-proline at the highest concentration tested (equivalent to that estimated for drought- and mite-stressed tomato plants at 10 dpi). In contrast, the effect was not significant when disks were treated with a concentration of proline equivalent to that estimated for drought- and mite-stressed tomato plants at 4 dpi (Figure 2.5). This result suggests a phagostimulant effect on *T. evansi* at the highest concentration of proline tested.

2.4 Discussion

Under a climate change scenario, invasive mite species, such as *T. evansi*, raise further concerns because they might expand to new geographical areas (Migeon et al., 2009) and have more generations per year, causing severe damage on tomato (Azandémè-Hounmalon et al., 2014). Our data revealed that mild and moderate drought stressed tomato plants trigger bottom-up effects on *T. evansi*, with these plants physiologically more suitable for mite development. Mite performance was enhanced on both mild and moderate drought-stressed tomato plants, as revealed by a significant increase in the number of eggs laid and of mobile forms, resulting in more leaf damage. It has been reported that plants under drought stress mobilize proteins into amino acids and carbohydrates into free sugars for osmotic adjustments (Hummel et al., 2010). We have found that both drought and *T. evansi* damage triggered the decline of foliar soluble protein and total nitrogen content, suggesting that mites and drought are accelerating leaf senescence and likely inducing the transference of nutrients to younger leaves (Mody et al., 2009), as reported in cotton for the combined effects of water-stress and *T. urticae* infestation (Sadras et al., 1998). In contrast, the amount of several free amino acids and free sugars increased significantly in drought-stressed tomato leaves, as already reported for tomato (Bauer et al., 1997), with proline the amino acid most clearly induced by drought. Interestingly, we have demonstrated that L-proline, a known phago-stimulant for insects (Mattson and Haack, 1987), also has a significant feeding stimulant effect on *T. evansi*, increasing both eggs laid and leaf damage area when added to tomato leaf discs. In addition, mite-infested, drought-stressed tomato plants showed significantly increased concentrations of five essential amino acids (valine, isoleucine, leucine, tyrosine and histidine), which have been reported to enhance the fecundity of the closely-related species *T. urticae* on strawberry, cucumber and chrysanthemum plants (Tulisalo, 1971; Dabrowski and Bielak, 1978). Therefore, *T. evansi* is probably taking advantage of the increase in these plant-derived essential amino acids to synthesize structural

proteins and enzymes, reducing the metabolic costs associated with digestion that can be reallocated for growth and development. A similar role can be proposed for the increase in free sugars, since it has been reported that fecundity of *T. urticae* correlated with sugar content in apple leaves (Wermelinger et al., 1991), and free sugars can also act as phagostimulants for some herbivores (Showler, 2013). Altogether, our data suggests that significant increases of available free sugars and essential amino acids, jointly with their phagostimulant effects, created a favourable environment for a better *T. evansi* performance on drought-stressed tomato leaves.

Another important aspect concerning the palatability of plants for mites is the level of antinutritional plant defense proteins that accumulate in response to abiotic and biotic stresses. As reported for other species (Mosolov and Valueva, 2011), drought consistently induced the inhibitory activity of tomato leaf extracts against cysteine (papain and cathepsin B) and serine (trypsin) proteases. This response was obtained for both *T. evansi* infested and non-infested plants, indicating that the response was driven by drought stress. By contrast, chymotrypsin inhibitory activity and the activity of PPO and POD varied depending on the time of plant exposure to drought stress, which may be a consequence of the complex regulatory pathways for some of these inducible defense genes in plants (Atkinson et al., 2015). It has been reported that *T. evansi* suppresses plant defenses in tomato by down-regulating the induction of both the salicylic acid and jasmonic acid signalling pathways (Kant et al., 2015). In particular, this was reflected in a reduction in the expression of serine protease inhibitors and PPO (Sarmiento et al., 2011; Alba et al., 2015), which are normally induced in tomato leaves in response to the attack by other spider mites (Kant et al., 2004). Likewise, we have shown that the inhibitory activity against serine proteases (trypsin and chymotrypsin) was not significantly induced in leaves attacked by *T. evansi*, when compared to non-infested tomato plants. However, we found that mite infestation induced the inhibitory activity against cysteine proteases (cathepsin B and papain), which were not considered in previous studies. These results are especially relevant, since proteolytic digestion in spider mites relies on cysteine proteases (including *T. evansi*, see below) and cysteine protease inhibitors have been shown to be harmful against the spider mite *T. urticae*, *in vivo* (Santamaria et al., 2012). We have also found that the specific activities of both PPO and POD were significantly increased in response to *T. evansi* infestation. These results may be considered somewhat contradictory with those obtained by Alba et al. (2015), who reported that *T. evansi* suppressed the expression of two PPO genes (PPO-D and PPO-F) in tomato. However, this suppression only occurred when compared with tomato plants infested

with mites from a non-suppressor *T. urticae* strain, but the expression of these genes in plants infested with *T. evansi* was slightly but significantly higher than on non-infested tomato plants (Alba et al., 2015). Thus, further studies to determine the *in vivo* effect of tomato cysteine protease inhibitors and PPOs on *T. evansi* are required.

Plant-feeding arthropods have developed a remarkable diversity of physiological adaptations to respond to changes in the nutritional composition of their host plants and to counteract the adverse effects of plant defenses including the regulation of digestive and detoxification enzymes (Grbic et al., 2011; Ortego, 2012; Zhu-Salzman, 2015). The hydrolysis of dietary proteins in spider mites relies on cysteine (cathepsin B-, cathepsin L- and legumain-like) and aspartyl (cathepsin D-like) proteases and aminopeptidases (Carrillo et al., 2011; Santamaria et al., 2015a). Our results indicate that *T. evansi* has a similar digestive proteolytic profile, hydrolyzing specific substrates for cathepsin D, cathepsin B, cathepsin L and legumain proteases and aminopeptidases. In addition, we determined the presence of α -amylase activity for the hydrolysis of carbohydrates in *T. evansi*, as already reported in other tetranychid mite species (Akimov and Barabanova, 1997). We found that cathepsin D- and legumain-like proteases and α -amylase activities were significantly reduced when *T. evansi* mites were fed on drought-stressed plants, but this reduction did not seem to affect their performance. The decrease of these mite hydrolytic activities could be explained by the ingestion of hydrolytic enzyme inhibitors from tomato plants. However, cathepsin B- and L-like activities in *T. evansi* were not affected, despite the induction of specific inhibitors for these two types of proteases in drought-stressed plants. These results suggest that *T. evansi* is capable of circumventing the adverse effects of at least some plant protease inhibitors. An alternative explanation for the reduction of some digestive protease and α -amylase activities in *T. evansi* could be related to the higher availability of nutrients in a readily assimilated form (essential amino acids and free sugars) in tomato plants subjected to drought conditions. Tomato plants are known to induce an array of secondary metabolites during periods of water deficit (Ripoll et al., 2014), thereby potentially increasing the concentration of compounds involved in plant defense. However, no significant changes in esterase, GST or P450 specific activities were observed when *T. evansi* was fed on drought-stressed plants. Secondary metabolites were not analyzed in this study, but it has been reported that *T. evansi* suppresses the expression of genes involved in regulation of secondary metabolites (Alba et al., 2015) and the release of inducible volatiles (Sarmiento et al., 2011). This down-regulation of plant defenses may render unnecessary the induction of the mite detoxification enzymes, and the metabolic resources saved diverted toward growth and

reproduction. Nevertheless, further studies will be needed to determine the combined effects of drought and *T. evansi* infestation in the regulation of secondary metabolites in tomato.

2.5 Conclusions

Our data reveal that both drought and *T. evansi* infestation induced significant changes in the nutritional quality of tomato plants, as more essential amino acids and free sugars are available. These changes trigger a bottom-up effect on key biological traits of *T. evansi* causing a highly significant increase in leaf damage and mite performance. Plant defense proteins were also induced by drought and/or mite infestation, but the mite physiological responses suggest that *T. evansi* is capable of circumventing their potential adverse effects. These findings support the “Plant stress hypothesis” and suggest that drought-stressed tomato plants even at a mild level may be more prone to *T. evansi* outbreaks in a climate change scenario, which might strongly affect tomato production on area-wide scales. Moreover, abiotic stresses can also exert a strong impact at higher tropic levels. Thus, it has been demonstrated recently that plant water status may also trigger bottom-up effects on the reproductive behaviour and fitness of some natural enemies, especially the omnivorous predators (Han et al., 2015; Seagraves et al., 2011), which might shape the arthropod structure. Therefore, this complexity of food-web structures should be considered when using species distribution models to infer climate change effects on crop pests.

2.6 Supporting information

Figure S2.1 Effect of mild drought at mite infestation and at 4 and 10 days post infestation (dpi) on: a) tomato stomatal conductance (gs); b) maximum quantum yield of PSII photochemistry (F_v/F_m); c) stem length; d) plant aerial dry weight. Data are mean $\pm$ SE. * Indicates significant difference within each time (Three-way ANOVA, Bonferroni *post hoc* test, $P < 0.05$).

Chapter 3

DROUGHT STRESS IN TOMATO INCREASES THE PERFORMANCE OF ADAPTED AND NON-ADAPTED STRAINS OF *Tetranychus urticae*

3.1 Introduction

The two-spotted spider mite, *Tetranychus urticae* Koch, is a polyphagous pest with a worldwide distribution. While this species has been reported feeding on more than 1000 plant species (Migeon and Dorkeld, 2006–2015), local mite populations do not perform equally well on all potential hosts (Gotoh et al., 1993; Kant et al., 2008). The capability of *T. urticae* to cope with the varying nutritional value and defense compounds of different host plants has been explained by the proliferation of gene families involved in digestion and detoxification processes (Grbic et al., 2011; Van Leeuwen and Dermauw, 2016) and by its transcriptional plasticity in response to host plant shifts (Dermauw et al., 2013; Wybouw et al., 2015). Moreover, it has been demonstrated that horizontally transferred genes from microorganisms facilitate the metabolism of toxic xenobiotics (Ahn et al., 2014).

T. urticae populations initially show low acceptance for tomato (*Solanum lycopersicum* L.) as a host, but they can rapidly become adapted (Fry, 1989; Agrawal et al., 2002), causing crop losses worldwide (Jayasinghe and Mallik, 2010; Meck et al., 2013). Studies on tomato–*T. urticae* interactions highlight the importance of secondary metabolites (e.g. phenylpropanoids, flavonoids and terpenoids) and anti-digestive proteins (e.g. protease inhibitors, polyphenol oxidases, amino acid catabolizing enzymes and peroxidases) as prominent components of tomato-induced defenses elicited by spider mite feeding (Li et al., 2002; Kant et al., 2004; Martel et al., 2015). Induction of these defenses is especially relevant, since the ingestion of protease inhibitors has been shown to be harmful to *T. urticae* (Carrillo et al., 2011; Santamaria et al., 2012). However, some genotypes of *T. urticae* are able to manipulate the main defense pathways in tomato, resulting in an attenuated induction of defense genes to levels at which they are less detrimental (Kant et al., 2008; Alba et al., 2015). Interestingly, Wybouw et al. (2015) demonstrated that during their adaptation to tomato, *T. urticae* extensively rearranged their xenobiotic metabolism through differential expression of genes that code for detoxifying enzymes and xenobiotic transporters, and acquired the ability to interfere with plant defenses. Candidate effector proteins in *T. urticae* that attenuate plant defenses have been identified in their saliva (Villarroel et al., 2016).

Because of their tight relationship with host plants, *T. urticae* has been reported to be differentially affected by water deficit. For example, *T. urticae* performance has been shown to increase (Hollingsworth and Berry, 1982; Nansen et al., 2010), decrease (Gould, 1978; Sadras et al., 1998) or vary (English-Loeb, 1990) depending on the plant species and the intensity of the water stress. Plants under drought stress are potentially more suitable as food because of

increased nutrient availability (Brodbeck and Strong, 1987; Showler, 2013), and can therefore contribute to improve herbivore performance (Huberty and Denno, 2004; Showler, 2013). A positive correlation has been reported between *T. urticae* fecundity and amino acid and sugar content on plants (Dabrowski and Bielak, 1978; Wermelinger et al., 1991). Aside from acting as direct sources of available nutrients, free sugars and amino acids have also been reported to be feeding stimulants for many phytophagous species (Showler, 2013). On the other hand, drought stress has been also associated with an increase in the production of plant defense compounds making stressed plants potentially less suitable for herbivores (Gould, 1978; Mattson and Haack, 1987). The resulting performance of *T. urticae* on drought stressed plants will then depend on the balance of induced nutrients and chemical defenses in the plant, and how the mites adapt to these changes. Drought stress triggers both the mobilization of plant nutritional compounds (Bauer et al., 1997) and the induction of plant defenses (English-Loeb et al., 1997; Inbar et al., 2001) in tomato. Recently, it has been reported that the tomato red spider mite *T. evansi*, an oligophagous mite feeding mainly on solanaceous plants, takes advantage of the changes in the nutritional quality of drought-stressed tomato plants and it is capable of circumventing the potential adverse effects of tomato plant defenses (Ximénez-Embún et al., 2016).

In the present study, we tested the effect of drought stressed tomato plants in both, tomato adapted and tomato non-adapted *T. urticae* strains, since its performance might vary depending on the rate of adaptation of mite populations to this host (Kant et al., 2008). Furthermore, changes in plant nutritional composition and defense proteins, as well as mite enzymatic (hydrolytic and detoxification) activities, were analyzed to understand how mite and plant physiological responses interact against each other under drought stress, which might provide insights for mite outbreaks under future deficit irrigation scenarios.

3.2 Materials and methods

3.2.1 Plant material and mite rearing

Tomato (cv. Moneymaker) plants were grown from seeds in 40 well trays. Plants with three expanded leaves were transferred to 2.5 L pots (diameter: 16 cm, height: 15 cm) (Maceflor, Valencia, Spain) filled with 600 g of Compo Sana Universal potting soil (Compo GmbH, Münster, Germany) and watered to saturation. Bean (*Phaseolus vulgaris* L. cv California Red Kidney Beans) seeds were germinated and transferred to 1.6 L pot (diameter: 15 cm, height: 13 cm) (Plásticos Alber, Granada, Spain) filled with 500 g of the same growing

medium. All plants were maintained in a rearing room at $25\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$, $50 \pm 5\%$ relative humidity and a 16 h light/8 h dark photoperiod. Two colonies of *T. urticae* derived from the London strain that was originated in Ontario (Canada) and kept in laboratory conditions for over 100 generations were used: tomato adapted (TA) and tomato non-adapted (TNA). The TA colony was maintained on detached tomato leaves for about 30 generations. The petiole of the leaves was in contact with a thin layer of water in the bottom of ventilated plastic cages (22 x 30 x 15 cm) to contain the mites and to maintain the leaf turgor. The TNA colony was maintained on potted bean plants for the same period of time. Both colonies were kept in two separate growth chambers (Sanyo MLR-350-H, Sanyo, Japan) at $25\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$, $70 \pm 5\%$ relative humidity and a 16 h light/8 h dark photoperiod.

3.2.2 Drought stress regime

Drought stress was attained by water deficiency as described by Ximénez-Embún et al. (2016). In brief, tomato plants were well-watered until they developed four-five expanded leaves, then we imposed three watering treatments, defined as control, and mild and moderate drought stress, in order to establish three irrigation regimes. Control plants were watered every two-three days to maintain the soil volumetric water content (H) up to 74%. For mild and moderate drought stress, watering was stopped for four and seven days, respectively, and thereafter plants were watered to maintain H between 36 and 50% in case of the mild stress and between 21 and 30% in case of the moderate stress. The H was determined gravimetrically by recording single plant pot weight (balance BSH 6000, PCE Iberica, Tobarra, Spain). Steady stress conditions were reached at about seven-nine days after ceasing irrigation. Both water stress regimes were above the wilting point associated with severe drought stress that was established at $H = 16\%$ for Moneymaker in our experimental conditions (Ximénez-Embún et al., 2016). The severity of drought stress was assessed by measuring the following parameters on the sub-terminal leaflet of the 4th leaf: a) stomatal conductance (gs) using a leaf porometer (SC-1 Decagon-T, Pullman, USA); and b) variations in maximum quantum yield of photosystem II photochemistry (Fv/Fm), using a FluorPen FP 100 (PSI, Drasov, Czech Republic). Plant growth was estimated by measuring the stem length (distance between the soil and the terminal bud) and by weighting the aerial part of the plant (transformed to dry weight by using the water content data calculated as referred below).

3.2.3 Bioassays to test drought stress effects on mite performance and plant nutritional composition

A factorial design with two levels of water stress (control or drought), two levels of *T. urticae* infestation (infested or non-infested) and two sampling times (4 or 10 days post infestation = dpi) was conducted. Mild and moderate drought-stressed plants were coupled with well-watered (control) plants in independent experiments. When drought stress conditions were reached a subset of tomato plants were infested. Mite females, of random age, were collected from the laboratory colony using a vacuum pump D-95 (Dinko S.A., Barcelona, Spain) with a sucking power of 10–50 mmHg connected to a modified Eppendorf. They were placed on the two sub-terminal leaflets of tomato leaves three, four, and five (eight mites per leaflet, each plant receiving 48 mites in total). All plants (infested and non-infested) were confined with a ventilated metacrylate cylinder fitting the pot and set up in a rearing chamber in a complete randomized block design (each experiment was repeated two-three times). Temperature and humidity inside the cylinders was recorded introducing USB dataloggers Log 32 (Dostmann electronic GmbH, Wertheim, Germany), on average the relative humidity was $79 \pm 2\%$ inside the well-watered cylinders, and $71 \pm 2\%$, and $58 \pm 2\%$ for mild and moderate drought stress conditions, respectively. The average temperature was 24 ± 1 °C in all cases. In each experiment, a subset of 74 plants was used to assess mite performance and the other subset for plant nutritional composition. Mite performance was assessed at 4 and 10 dpi for the two levels of drought stress (moderate and mild) when infested with the TA strain, whereas only moderate drought stress was tested with the TNA strain. These sampling times were chosen to study the effect of drought stress on two consecutive generations of spider mites. At each time, 15 infested plants per treatment (control or drought) were analyzed in the moderate stress experiments, and nine infested plants per treatment in the mild stress one. All leaves were detached from the plants, and the number of eggs and mobile mite stages (larvae, nymphs and adults) counted under a stereomicroscope M125 (Leica Mycosystem, Wetzlar, Germany). The leaf damaged area (mm² of chlorotic lesions) was determined as described by Ximénez-Embún et al. (2016) by scanning the damaged leaflets using hp scanjet (HP Scanjet 5590 Digital Flatbed Scanner series, USA) and analyzing the scanned leaflets with the program GIMP 2.8 (www.gimp.org). The nutritional composition of plant material was analyzed in control and moderate drought-stressed plants, infested and non-infested (six plants per treatment). The samples were obtained at 4 and 10 dpi with the TA strain, but only at 4 dpi with the TNA strain since most mites from the TNA strain have already moved away from the plant at 10 dpi (see results section). The left leaflets from leaves three, four, and five were pooled and grounded in

liquid nitrogen to a fine powder and stored for free amino acid and protein analysis. The right leaflets from the same leaves were dried together in an oven at 70 °C for 3 days, weighed before and after drying to assess the percentage of water and ground using a mortar and pestle to obtain a fine powder and stored for C, N and free sugar analysis.

3.2.4 Bioassays to test drought stress effects on plant defense proteins and mite enzymatic activities

A factorial design with two levels of water stress (control or moderate drought) and three levels of *T. urticae* infestation (infested with TA, infested with TNA or non-infested) was conducted. When drought stress conditions were reached, a subset of tomato plants were infested by placing 150 adult female mites on leaves three, four, and five (450 mites per plant), laying a thin barrier of Lanolin (Manuel Riesgo S.A., Madrid, Spain) on the petiole to contain the mites. Plants (infested and non-infested) were placed in a rearing chamber in a complete randomized block design (the experiment was repeated two times in time) and maintained at 25 ± 1 °C, $50 \pm 5\%$ relative humidity and a 16 h light/8 h dark photoperiod. The experiment was terminated at 4 dpi. Plant material (six per treatment) was collected by sampling together infested leaflets from leaves three, four and five. Mites were collected from a different subset of plants (six per treatment) by using the pump system previously described. Plant and mite material was immediately frozen in liquid nitrogen and stored at -80 °C. Mites from the TNA strain reared on bean plants were also collected for comparison.

3.2.5 Chemical and biochemical analysis

Unless otherwise specified, all chemical compounds used were from Sigma-Aldrich (St Luis, USA). Fluorimetric measurements were made using a Varioskan Flash reader (ThermoFisher Scientific, Willmington, USA), and spectrophotometric measurements with a VERSAmax microplate reader (Molecular Devices Corp., Sunnyvale, USA).

Total C and N composition. Samples of 1 mg of dried leaf powder were analyzed to determine total nitrogen and carbon concentration at the Elemental Microelement Center of Complutense University (Madrid, Spain) by using a microelement analyzer LECO CHNS-932 (LECO, St Joseph, MI, USA).

Free sugars. Samples of 3 mg of dried leaf powder were homogenized in 650 µl of ethanol 95% (v/v), heated at 80 °C for 20 min, centrifuged at 10,000 rpm for 10 min, and the supernatant collected. The process was repeated two more times and the three supernatants were pooled. A volume of 750 µl of the mixture was dried on a SpeedVac Concentrator Savant SVC

100H (ThermoFisher scientific, Willmington, DE, USA) and redissolved in 500 µl of water. Soluble carbohydrate concentration was estimated by the anthrone method (Maness, 2010) using glucose as standard. In brief, 1 ml of anthrone reagent (0.2% v/v anthrone on 95% sulfuric acid) was added to the extract, heated at 90 °C for 15 min, and the absorbance measured at 630 nm.

Free amino acids. The extraction of the free amino acids was done as described by Hacham et al. (2002). Samples of 50 mg of leaf frozen powder were homogenized with 600 µl of water:chloroform:methanol (3:5:12 v/v/v). After centrifugation at 12,000 rpm for 2 min, the supernatant was collected and the residue was re-extracted with 600 µl of the same mixture, pooling the two supernatants. A mixture of 300 µl of chloroform and 450 µl of water were added to the supernatants, and after centrifugation the upper water:methanol phase was collected and dried in the SpeedVac. The samples were dissolved on 100 µl of sodium citrate loading buffer pH 2.2 (Biochrom, USA) and 10 µl were injected on a Biochrom 30 Amino Acid Analyser (Biochrom, USA) at the Protein Chemistry Service at CIB (CSIC, Madrid, Spain).

Soluble protein. Samples of 100 mg of leaf frozen powder were homogenized in 500 µl of 0.15 M NaCl, ground with fine sand. The homogenate was centrifuged at 12,000 rpm for 5 min at 4 °C, and the soluble protein quantified by absorbance at 280 nm on a Nanodrop 1000 spectrophotometer (ThermoFisher scientific, Willmington, USA).

Plant defense proteins. Samples of 100 mg of leaf frozen powder were homogenized with 500 µl of extraction buffer (0.15 M NaCl for protease inhibitors, and 0.1 M phosphate buffer, pH 7.0; 5% w:v polyvinylpolypyrrolidone for oxidative enzymes) and soluble protein quantified as explained above. The inhibitory activity of plant protein extracts was tested against commercial enzymes: papain (EC 3.4.22.2), cathepsin B from bovine spleen (EC 3.4.22.1), trypsin from bovine pancreas (EC 3.4.21.4), a-chymotrypsin from bovine pancreas (EC 3.4.21.1), cathepsin D from bovine spleen (EC 3.4.23.5) and leucine aminopeptidase from porcine pancreas (EC 3.4.11.1); as described by Ximénez-Embún et al. (2016). Reaction conditions are summarized in Table S3.1. Results were expressed as a percentage of protease activity inhibited. Polyphenol oxidase (PPO) and Peroxidase (POD) activities of plant extracts were analyzed as described by Ximénez-Embún et al. (2016). Reaction conditions are summarized in Table S3.2.

Mite enzymatic activities. Samples of 1 mg of mites were homogenized in 100 µl of 0.15 M NaCl, centrifuged at 12,000 rpm for 5 min and the supernatant was collected. Total

protein content was determined according to the method of Bradford (1976). Hydrolytic enzymes (cathepsin B-, cathepsin L-, cathepsin D- and legumain-like) activities were assayed as described by Santamaria et al. (2015a), and leucine aminopeptidase- and α -amylase-like activities and detoxification enzymes (esterase, glutathione S-transferase and P450) activities as described by Ximénez-Embún et al. (2016). Reaction conditions are summarized in Table S3.3.

3.2.6 Statistical analysis

All plant and mite parameters analyzed were checked for the assumptions of normality and heteroscedasticity, and transformed if necessary. Stem length and stomatal conductance were $\log_{10}(x)$ transformed, Fv/Fm was $\log_{10}(x + 1)$ transformed; and these three parameters and stem dry weight data were statistically analyzed using a three-way ANOVA (using as fixed factors drought stress treatment, *T. urticae* infestation and time). The number of *T. urticae* eggs and mobile forms and leaf damaged area were $\log_{10}(x)$ transformed and analyzed by a two-way ANOVA (drought stress treatment, time). For stem length, plant aerial dry weight, stomatal conductance, Fv/Fm, mite eggs and mobile forms, and leaf damage a Bonferroni *post hoc* test was performed to compare drought-stress treatments within each time. Percentage of water, nitrogen, protein, free amino acids and free sugars, as well as C:N ratio and protease inhibition were $\arcsin(\sqrt{x})$ transformed. These data and the oxidative enzyme activities were analyzed using a two-way ANOVA for each time separately using as fixed factors drought stress treatment and *T. urticae* infestation. A Newman-Keuls *post hoc* test was performed to see differences of mean between treatments, except for essential and non-essential amino acids that were analyzed by a Dunnett *post hoc* test to compare the different treatments with respect to control non-infested plants. Spider mite enzymatic activity data were $\log_{10}(x)$ transformed and analyzed by a one-way ANOVA, using Newman-Keuls as *post hoc* test.

3.3 Results

3.3.1 Effects of drought stress on stomatal conductance, photosynthetic efficiency and tomato plant growth

Drought stress significantly affected leaf stomatal conductance, which was reduced between five and seven times in the case of moderate drought stress and between 1.3 and 1.8 times in the case of mild drought stress when compared to control plants (Figure S3.1a). The photosynthetic efficiency determined as F_v/F_m was also significantly lower on drought stressed plants in all experiments at 10 dpi (Figure S3.1b). These results corroborated that the severity of the stress was different for the two water stress regimes imposed (moderate and mild), but severe drought stress conditions were never reached, since F_v/F_m values were never below 0.7 (Ritchie, 2006). Moderate drought stress slowed plant growth, as both stem length and aerial dry weight were significantly smaller compared to control plants at most measured points, whereas mild drought stress only produced a significant reduction in aerial dry weight at 10 dpi (Figure S3.2a, b). The infestation by *T. urticae* (TA or TNA) did not affect stomatal conductance, photosynthetic efficiency and tomato plant growth. Thus, data from infested and non-infested plants were pooled within each experiment.

3.3.2 Effects of drought stress on *T. urticae* (TA and TNA strains) population growth and leaf damage

The performance of mites from the TA strain was enhanced when reared on both moderate and mild drought-stressed tomato plants. Females laid a significantly higher number of eggs on moderately stressed plants than on control plants at 4 dpi (2.4-fold) and 10 dpi (1.5 fold) (Figure 3.1d). The F1 generation (mobile forms) was significantly more abundant on moderately stressed plants at 4 and 10 dpi (1.4 and 2-fold respectively) (Figure 3.1e). As a result, damaged leaf area doubled in moderately stressed plants compared to control tomatoes (Figure 3.1f). The same trend was observed under mild drought stress, but the differences among treatments were less pronounced and only significant in the case of number of eggs laid (1.5 fold) at 4 dpi, and mobile forms (1.4 fold) and leaf damage (1.5 fold) at 10 dpi (Figure 3.1a, b and c). Population growth in the case of the TNA strain was negative, with lower number of mobile forms at 4 and 10 dpi than at the beginning of the assay. Only a few dead mites were found on the tomato leaves, indicating that mites were most likely dispersing from the plant. Nevertheless, they laid more eggs (about 2 fold) and the number of mobile forms was higher (about 1.5 fold) on drought-stressed tomato plants than on control plants (Figure 3.1g and h).

Damaged leaf area as a result of TNA infestation was markedly lower than that produced by TA infestation and could not be reliably measured.

Figure 3.1 Performance of *T. urticae* tomato adapted strain (TA) on mild (a, b and c) and moderate (d, e and f) drought stressed tomato plants and of the non-adapted strain (TNA) on moderate stressed plants (g and h). Plants were infested with 48 adult females (day 0) and eggs laid on plants (a, d and g), total mobile forms (b, e and h) and leaf damaged area (c and f) assessed at 4 and 10 days post infestation (dpi). Leaf damage was negligible in TNA. Data are mean $\pm$ SE. * indicates statistically significant differences between drought stress treatments within each time (Two-way ANOVA, Bonferroni *post hoc* test, $p < 0.05$).

3.3.3 Changes in plant nutritional composition induced by drought stress and *T. urticae* (TA and TNA)

Drought stress was the most significant factor in plant nutritional composition (Table 3.1), resulting in a decrease in the relative amount of water (all cases) and protein (TA at 10 dpi), and an increase on total free amino acids (TA at 10 dpi and TNA at 4 dpi) and sugars (all

Table 3.1 Effect of moderate drought stress and infestation by *T. urticae* tomato adapted (TA) and non-adapted (TNA) strains on nutritional composition of tomato leaves.

	Non-infested		Infested		ANOVA (p<0.05)
	Control	Drought	Control	Drought	
TA strain					
4 Days Post Infestation					
Water ¹	93.0±0.4 a	91.0±0.6b	93.0±0.4 a	91.0±0.7b	D
Nitrogen ²	6.7±0.3	5.9±0.3	6.8±0.2	6.0±0.3	D
Protein ²	21.7±1.8	18.5±2.1	24.0±1.2	25.8±2.7	I
Free aa ²	1.1±0.2	1.3±0.2	1.1±0.1	1.2±0.3	-
C:N ³	5.8±0.1	6.6±0.3	5.8±0.1	6.3±0.3	D
Free Sugars ²	3.3±0.1 a	3.6±0.1 a	4.5±0.1 b	4.9±0.2 b	D; I
10 Days Post Infestation					
Water ¹	93.0±0.7 a	91.0±0.5 b	93.0±0.3 a	92.0±0.4 ab	D
Nitrogen ²	6.2±0.1	6.1±0.3	6.0±0.3	5.9±0.2	-
Protein ²	17.7±1.3 ab	14.5±1.5 b	25.7±2.6 a	13.6±1.9 b	D
Free aa ²	0.9±0.2	1.2±0.1	0.9±0.1	1.4±0.2	D
C:N ³	6.3±0.1	6.3±0.2	6.1±0.1	6.3±0.1	-
Free Sugars ²	3.5±0.1	4.2±0.1	3.8±0.2	4.2±0.4	D
TNA strain					
4 Days Post Infestation					
Water ¹	94.0±0.0 a	89.0±1.0 ab	91.0±2.0 ab	87.0±2.0 b	D
Nitrogen ²	5.7±0.1	6.1±0.2	5.7±0.1	6.1±0.1	D
Protein ²	28.0±6.0	23.0±2.0	18.0±4.0	14.0±3.0	I
Free aa ²	0.6±0.1 ab	1.0±0.1 b	0.5±0.1 a	1.0±0.2 b	D
C:N ³	6.2±0.2	6.1±0.2	6.1±0.1	6.0±0.1	-
Free Sugars ²	3.6±0.2 a	4.3±0.2 a	4.1±0.1 a	5.7±0.6 b	D; I
Data are mean ± SE: (1) % fresh weight (2) % dry weight (3) ratio. D (drought stress) and I (Mites infestation) indicates significant factor in a two-way ANOVA. Different lower case letters within rows indicates significant differences (Newman-Keuls <i>post hoc</i> at p<0.05)					

cases) and C:N ratio (TA at 4 dpi). The effect of drought stress on nitrogen was not consistent, as it decreased in the TA experiment at 4 dpi and increased in the TNA experiment. Concentrations of free sugars increased in plants exposed to both mite strains at 4 dpi, while total protein quantities increased in tomatoes exposed to mites from the TA strain and decreased when tomatoes were fed upon by mites from the TNA strain.

The levels of specific amino acids, classified as essential or non-essential for *T. urticae* according to Rodriguez and Hampton (1966), were also analyzed (Figure 3.2). Proline was the amino acid most highly induced by drought stress alone or in combination with mites (about 10–16 times depending on the experiment). Likewise, most of the essential amino acids for this species (valine, isoleucine, leucine, tyrosine, phenylalanine, histidine, lysine and arginine) and

Figure 3.2 Levels of free amino acids in tomato leaves from control plants and plants under moderate drought stress and/or infested with *T. urticae* tomato adapted (TA) and non-adapted (TNA) strains at 10 and 4 days post infestation (dpi), respectively. Data are mean $\pm$ SE of amino acid contents (% dry weight) represented at logarithmic scale. The division between essential and non-essential amino acids for *T. urticae* is based on Rodriguez and Hampton (1966). * Indicate significant difference of the treatment with respect to control plants (Two-way ANOVA, Dunnet *post hoc* test, $p < 0.05$).

the non-essential amino acid serine were also induced by drought stress alone or in combination with TA or TNA mite infestation. However, none of the amino acids analyzed were significantly affected when plants were stressed only by mite infestation.

3.3.4 Effect of drought stress and *T. urticae* (TA and TNA) on plant defense proteins

Tomato plant defense proteins were affected by both drought stress and mite infestation. The response was generally higher to the TNA strain than to the TA strain (Table 3.2). The inhibitory activity against chymotrypsin was significantly induced by both TA and TNA strains and the inhibitory activity against trypsin by TNA, independently of the watering conditions. The infestation with TA only induced trypsin inhibitory activity when combined with drought stress, and aminopeptidase inhibitory activity was only induced by TNA under control conditions. Drought stress induced peroxidase activity, but it was only significant when combined with TA mite infestation, whereas the interaction drought stress-infestation was significant for poliphenol oxidase activity. No significant effect was found on the levels of cathepsin B, D, and papain inhibition.

Table 3.2 Effect of moderate drought stress and infestation with *T. urticae* tomato adapted (TA) and non-adapted (TNA) strains on tomato plant defense proteins at 4 days post infestation.

	Non-infested		Infested with TA		Infested with TNA		ANOVA (p<0.05)
	Control	Drought	Control	Drought	Control	Drought	
Protease inhibitors (% inhibition)							
Cathepsin B	38±7	27±3	36±6	35±5	45±9	29±5	-
Papain	54±6	39±7	45±2	44±5	50±7	45±9	-
Cathepsin D	52±4	53±2	51±3	56±3	52±4	52±2	-
Trypsin	40±3 a	33±5 a	41±7 a	61±7 b	64±8 b	84±5 c	D, I
Chymotrypsin	47±4 a	46±6 a	71±10 b	91±2 c	86±4 c	97±1 c	D,I
Aminopeptidase	32±5 ab	23±1 a	25±3 a	27±4 a	43±4 b	33±2 ab	D,I
Oxidative enzymes (specific activity)							
PPO ¹	4.2±0.3	3.3±0.3	3.3±0.5	4.3±0.2	4.0±0.5	3.2±0.4	D*I
POD ²	1.9±0.2 ab	3.0±0.5 ab	1.6±0.3 a	3.4±0.5 b	2.2±0.4 ab	2.7±0.5 ab	D

Data are mean ± SE. D (drought stress) and I (mite infestation) indicate significant factors in a two-way ANOVA. Different lower case letters within rows indicates significant differences (Newman-Keuls *post hoc* test at p<0.05).

(1) PPO (poliphenol oxidases): nmol hidolyzed Cathecol/ mg Protein*min

(2) POD (peroxidases): nmol hidolyzed Guaiacol/ mg Protein*min.

3.3.5. Physiological response of *T. urticae* (TA and TNA) to drought-stressed tomato plants

None of the hydrolytic enzymes activities analyzed in mites of the TA and TNA strain was affected by the water status of the tomato plant host, whereas there were significant differences between mite strains (Table 3.3). When feeding on tomato, the TNA strain had lower activity than the TA strain for cathepsin B and legumain-like proteases, in both drought stressed and well-watered plants, and for cathepsin L- and D-like proteases and α -amylase activities in well-watered plants. In contrast, aminopeptidase activity was higher on the TNA strain. When the TNA strain was fed on bean (the habitual host plant for this strain), all the hydrolytic activities, except cathepsin D-like and aminopeptidase, were significantly higher

Table 3.3 Enzymatic activities of mites from *T. urticae* tomato adapted (TA) and non-adapted (TNA) strains when feeding for 4 days on control or moderate drought stressed tomato plants. TNA mites feeding on beans were also analyzed.

	TA strain		TNA strain		
	Tomato control	Tomato drought	Tomato control	Tomato drought	Bean
Protein ¹	43±3	46±3	48±4	50±5	44±2
Hydrolytic enzymes ²					
Cathepsin B	4.9±0.2 b	3.9±0.3 b	0.6±0.1 c	0.7±0.1 c	9.0±1.0 a
Cathepsin L	26±2 b	21±1 bc	14±2 c	21±4 bc	39±3 a
Cathepsin D	23±2 a	20±3 ab	14±2 b	13±1 b	18±2 ab
Legumain	1.4±0.1 a	1.2±0.1 a	0.3±0.1 b	0.6±0.1 b	1.1±0.1 a
Aminopeptidase	5.5±0.4 b	4.7±0.2 b	8.5±0.3 a	7.8±0.7 a	8.7±0.7 a
α-Amylase	62±7 ab	59±7 ab	26±5 c	40±8 bc	79±8 a
Detoxification enzymes ²					
Esterase	263±35 a	181±22 ab	99±34 bc	75±35 bc	22±8 c
GST	904±211	1126±298	667±168	743±210	1347±350
P450	0.75±0.09	0.74±0.06	0.60±0.08	0.49±0.11	0.63±0.10

Data are mean $\pm$ SE. Different lower case letters within rows indicate significant differences (One-way ANOVA, Newman-Keuls *post hoc* test at $p < 0.05$).

(1) $\mu\text{g}/\text{mg}$ fresh weight

(2) Specific activity as nmol substrate hydrolyzed per min and mg of protein for Cathepsin B, L, D, legumain, aminopeptidase, α -amylase and esterase. nmol CDNB conjugated per min and mg of protein for GST and nmol cytochrome c reduced per min and mg of protein for P450.

than when fed on tomato, and even higher than for TA in the case of cathepsin B-, L-like proteases and aminopeptidase.

Among the detoxification enzymes studied, we only found significant differences between the strains in the case of esterase activity (Table 3.3). The TA strain showed the highest activity, the TNA strain reared on bean the lowest, and the TNA strain feeding on tomato an intermediate level.

3.4 Discussion

Performance of *T. urticae* on tomato depends on the rate of adaptation of mite populations to this particular host (Agrawal et al., 2002; Kant et al., 2008). Moreover, some of the changes induced by drought stress on tomato nutritional composition and plant defenses (Bauer et al., 1997; English-Loeb et al., 1997; Inbar et al., 2001) have been identified as key factors affecting mite host preferences and performance (Kant et al., 2008; Wybouw et al., 2015). Our data revealed that drought-stressed tomato plants increased the performance of *T. urticae*, enhancing population growth of a tomato adapted TA strain and being more suitable as a host for a non-adapted TNA strain. These findings could have significant implications for mite outbreaks under future climate change scenarios, when longer periods of drought and less water availability are expected for irrigated crops like tomato in semiarid environments (IPCC, 2013). Moreover, *T. urticae* is expected to have more generations per year as a direct effect of rising temperature on mite developmental rates, increasing mite pressure (Luedeling et al., 2011), which might contribute to exacerbate mite damage.

We have found that the increases of available free sugars and essential amino acids, which are limiting nutrients for mite growth and reproduction, seemed to improve the nutritional value of drought-stressed tomato plants for *T. urticae*. Plants under drought stress mobilize existing proteins and complex carbohydrates into amino acids and simple sugars, respectively, for osmotic adjustments and for the transference of available plant nitrogen and carbon to younger leaves and reproductive organs (Hummel et al., 2010; Showler, 2013). Drought-stressed tomato leaves showed an increase in the concentrations of total free amino acids and free sugars and a decrease of total protein, as already reported for tomato (Bauer et al., 1997; Ximénez-Embún et al., 2016). However, the effect of drought stress on nitrogen was not consistent, as it decreased in the assay performed with the TA strain but increased in the assay performed with the TNA strain. Likewise, nitrogen content of tomato plants has been reported to increase under drought stress in some cases (English-Loeb et al., 1997) and decrease

in others (Inbar et al., 2001). Interestingly, when particular amino acids were analyzed, most of those considered as essential for *T. urticae* (Rodriguez and Hampton, 1966) rose in drought stressed tomato leaves. Some of these essential amino acids have been reported to stimulate *T. urticae* feeding (Tulisalo, 1971; Dabrowski and Bielak, 1978). Nonetheless, the amino acid most clearly induced by drought stress was proline, which is not an essential nutrient. Proline has been shown to stimulate feeding in many different phytophagous arthropods (Showler, 2013; Ximénez-Embún et al., 2016). In addition, proline is unique among the amino acids because can be used by arthropods as a direct energy substrate for glycolysis and the production of ATP (Scaraffia and Wells, 2003). A similar role as sources of available nutrients can be proposed for the increase in free sugars, since it has been reported that fecundity of *T. urticae* correlated with sugar content in plants (Wermelinger et al., 1991).

Herbivore attack can also induce changes in the primary metabolism of plants, including the reallocation of carbon and nitrogen resources (Zhou et al., 2015; Martel et al., 2015). We observed that the levels of free sugars were increased in tomato leaves infested with mites from both the TA and the TNA strains. However, mite infestation had different effects on total protein concentration, as it was reduced by TNA while it increased after 4 dpi with TA, though no differences in the levels of total or specific free amino acids were observed. Likewise, Ximénez-Embún et al. (2016) reported that a tomato adapted strain of *T. evansi* also triggered the levels of free sugars in tomato leaves. As a result, more edible carbohydrates are apparently available in mite-exposed tomato plants. Our results do not allow distinguish whether these effects correspond to a plant defensive response, which may result in subsequent carbon remobilization to other tissues, or that the mites are manipulating plant primary metabolism for their own benefit.

The induction levels of several tomato defense genes were found to correlate negatively with *T. urticae* performance (Kant et al., 2008; Wybouw et al., 2015). Our data revealed that tomato plants responded to mite infestation by the induction of serine (trypsin and chymotrypsin) and aminopeptidase protease inhibitors (PIs), resulting the inhibitory activity higher in response to the TNA than to the TA mites, despite the damage inferred by the adapted strain was higher. These results are in line with those previously reported, in which serine PIs has been shown to be consistently induced upon *T. urticae* attack, but this response is attenuated by tomato adapted strains (Li et al., 2002; Kant et al., 2004, 2008; Alba et al., 2015). However, we did not observe mite induction of tomato polyphenol oxidase (PPO) and peroxidase (POD) activities, as found in some of these studies (Alba et al., 2015; Martel et al., 2015). This could

be related to the fact that these genes are induced in tomato as an early response to *T. urticae* infestation, maximum induction occurring within 24 h post infestation, whereas our experiment run for at least 4 days. Regardless, our results appear to be species specific since contrast with those reported for tomato-*T. evansi* interactions, in which this mite infestation induce tomato cysteine PI activity and the specific activities of both PPO and POD, but has no effect on the plant inhibitory activity against serine proteases (Ximénez-Embún et al., 2016).

Proteolytic digestion in *T. urticae* relies mostly on cysteine (cathepsin B-, cathepsin-L and legumain-like) and aspartyl (cathepsin D-like) proteases and aminopeptidases, whereas serine proteases do not appear to be directly involved in the hydrolysis of dietary proteins (Carrillo et al., 2011; Santamaria et al., 2015a). Thus, tomato serine PIs may probably be targeting other physiological processes in mites, such as the regulation of growth and development (Santamaria et al., 2012). Indeed, both cysteine and serine PIs can be harmful when ingested by *T. urticae* (Carrillo et al., 2011; Santamaria et al., 2012, 2015b). Interestingly, we have found that when feeding on tomato, cysteine and aspartyl protease and α -amylase activities were higher in *T. urticae* mites of the TA strain, whereas aminopeptidase activity was higher in the TNA strain. The increase of the aminopeptidase activity in the TNA strain can be associated to the induction of aminopeptidase PIs in tomatoes infested with mites from this strain. However, the rise of the other hydrolytic activities in the TA strain seems to be related to its higher feeding rate, since tomato cysteine and aspartyl PIs were not affected by *T. urticae* feeding. Moreover, cathepsin B- and L-like protease and α -amylase activities increased significantly in TNA when feeding on bean, its preferred host, resulting even higher than for TA in tomato. Nonetheless, we can not discard that the higher activity of these digestive proteases in the TA strain may have an adaptive value, since gut proteases have the potential to inactivate plant defense proteins targeting the digestive system or other tissues (Ortego, 2012). In this sense, it has been shown that the expression of cysteine and aspartyl protease genes increased in *T. urticae* after feeding on plants over-expressing the HvCPI-6 cystatin, that specifically targets cathepsins B and L, but also when feeding on plants over-expressing the CMe trypsin inhibitor that targets serine proteases (Santamaria et al., 2015a). However, we did not find differences for any of the *T. urticae* hydrolytic enzyme activities analyzed when fed on drought-stressed plants, despite drought stress also induced serine PIs and POD when combined with mite infestation.

T. urticae produces large amounts of detoxifying enzymes to circumvent plant secondary metabolites (Grbic et al., 2011; Van Leeuwen and Dermauw, 2016), and

transcriptional changes in genes coding for P450 s, GSTs, and esterases has been reported to occur during the adaptation process of *T. urticae* to tomato (Dermauw et al., 2013; Wybouw et al., 2015). We have found that esterase activity was significantly higher in mites of the TA strain, which can be related to their adaptation to tomato plants. An induction of the esterase activity was also observed when mites of the TNA strain maintained on beans were transferred to tomato, probably in response to some of the xenobiotics present in tomato. However, the level of activity was lower than that of the TA strain on tomato, which may partially explain their lower performance.

3.5 Conclusions

Our data indicate that tomato adapted and non-adapted *T. urticae* benefit from the improved nutritional value of tomato plants induced by drought stress (increased concentrations of essential amino acids and free sugars). Yet, the improved performance of the tomato adapted strain could be associated to changes on the digestive (higher cysteine and aspartyl protease and α -amylase activities) and detoxification (higher esterase activity) physiology of the mites and the attenuation of some of the defense responses (PIs) of the plant. Adaptation to tomato and drought stress favored mite performance, having an additive effect on most of the physiological parameters analyzed. However, assessing the relative impacts of plant nutrients versus defenses will require further research. As drought events are expected to be more frequent and prolonged due to climate change (IPCC, 2013), our results imply an increase in the risk of outbreaks of *T. urticae* on tomato by significantly enhancing population growth of adapted populations and increasing the suitability of tomato as a host for non-adapted ones. Moreover, breeding programs aimed at improving drought tolerance in tomato might take into consideration varieties whose nutritional composition is not enhanced by drought stress for avoiding or mitigating spider mite damage.

3.6 Supporting information

Figure S3.1 Effect of drought stress (moderate and mild) on **A)** tomato stomatal conductance (gs) and **B)** maximum quantum yield of PSII photochemistry (Fv/Fm) at mite infestation and at 4 and 10 days post infestation (dpi). Data (mean $\pm$ SE) shown are average of the values on infested and non-infested plants on experiments with *T. urticae* adapted (TA) and no adapted (NTA) strains, as mite infestation didn't show a significant effect.* indicates a statistically significant differences between drought strain treatments within each time (Three-way ANOVA, Bonferroni *post hoc* test, $p < 0.05$).

Figure S3.2 Effect of drought stress (moderate and mild) on tomato plant growth, expressed as A) aerial dry weight at 4 and 10 days post infestation (dpi) and B) stem length through the experiment. Data (mean $\pm$ SE) shown are average of infested and non-infested plants on experiments with *T. urticae* adapted (TA) and no adapted (NTA), as mite infestation didn't have a significant effect. * indicates a statistically significant difference between drought treatments within each time (Three-way ANOVA, Bonferroni *post hoc* test, $p < 0.05$).

Table S3.1 Summary of analytical methods to assess the inhibitory activity of plant protein extracts ¹

Commercial enzyme ²	Substrate ³	Buffer ⁴	Incubation	Measurement ⁵
Cathepsin B from bovine spleen (EC 3.4.22.1)	Z-RR-AMC	100 mM NA phosphate, pH 6.0 (10 mM L-cysteine, 10 mM EDTA, 0.01% (v/v) Brij 35)	1 h at 28 °C	excitation filter 350 nm
Papain (EC 3.4.22.2)	Z-FR-AMC	100 mM Na phosphate, pH 6.0 (10 mM L-cysteine, 10 mM EDTA, 0.01% (v/v) Brij 35)	1 h at 28 °C	excitation filter 350 nm
Cathepsin D from bovine spleen (EC 3.4.23.5)	MocAc-GKPILFFRLK	100 mM sodium citrate, pH 3.5 (0.15 M NaCl, 5 mM MgCl ₂)	10 min at 30 °C	excitation filter 328 nm
Trypsin from bovine pancreas (EC α -Chymotrypsin from bovine pancreas (EC	Z-LA-AMC	100 mM Tris-HCl, pH 7.5 (0.15 M NaCl, 5 mM MgCl ₂)	1 h at 35 °C	excitation filter 350 nm
	SucAAPF-AMC	100 mM Tris-HCl, pH 7.5 (0.15 M NaCl, 5 mM MgCl ₂)	1 h at 35 °C	excitation filter 350 nm
Leucine aminopeptidase from	LpNa	100 mM Tris-HCl, pH 8 (0.15 M NaCl, 5 mM MgCl ₂)	1 h at 30 °C	absorbance at 410 nm

¹ Procedures adapted from Ximénez-Embún et al. (2016). Samples of 20 µg of plant protein extracts (40 µg in case of leucine aminopeptidase) were preincubated for 10 min with 100 ng of the commercial enzyme.

² All purchased from Sigma-Aldrich (St Luis, USA).

³ The substrates were added at a final concentration of 20 µM. Z-RR-AMC (N-carbobenzoxyloxy-Arg-Arg-7-amido-4-methylcoumarin) for cathepsin B, Z-FR-AMC (N-carbobenzoxyloxy-Phe-Arg-7-amido-4-methylcoumarin) for papain, Z-LA-AMC (Z-L-Arg-7-amido-4-methylcoumarin) for trypsin, SucAAPF-AMC (Suc-Ala-Ala-Pro-Phe-7-amido-4-methylcoumarin) for chymotrypsin, all purchased from Calbiochem (MerkMilipore, Billerica, USA), MocAc-GKPILFFRLK(Dnp)-D-R- NH₂ from Peptanova (Germany) for cathepsin D, and LpNa (L-leucine p-nitroanilide) from Sigma-Aldrich (St Luis, USA) for leucine aminopeptidase.

⁴ Concentrations are expressed at molarity in the reaction mixture.

⁵ AMC (7-amino-4-methylcoumarin) (Bachem, Switzerland) as standard for all fluorescent substrates, except MCA (MoCAC-Pro-Leu-Gly) (Peptanova GmbH, Germany) for cathepsin D. Double blanks were used to account for spontaneous breakdown of substrates and the plant protease activity, and all assays were done in duplicate.

Table S3.2 Summary of analytical methods to assess the enzymatic activity of plant protein extracts

Plant Oxidative Enzymes ¹	Substrate ²	Buffer ³	Incubation	Measurement
Polyphenol oxidase (PPO)	catechol (40 mM)	Tris-HCl pH 8.5	1 h at 30 °C	absorbance at 420 nm
Peroxidase (POD)	guaiacol (5 mM)	Potassium phosphate pH 6, with 2.5 mM H ₂ O ₂	10 min at 30 °C	absorbance at 470 nm

¹ Procedures adapted from Ximénez-Embún et al. (2016). Assays were performed with 20 µl (PPO activity) and 20 µl of a 1:10 dilution (POD activity) of the plant protein extract

² The substrates were added at a final concentration of 40 mM (PBO) and 5 mM (POD). Both from Sigma-Aldrich (St Luis, USA).

³ Concentrations are expressed at molarity in the reaction mixture.

Table S3.3 Summary of analytical methods to assess the enzymatic activity of mite extracts

	Substrate ²	Buffer ³	Incubation	Measurement ⁴
Hydrolytic Enzymes ¹				
Cathepsin B-like	Z-RR-AMC	100 mM sodium citrate, pH 5.5 (0.15 M NaCl, 1 mM DTT)	15 min at 30 °C	excitation filter 350 nm emission filter 465 nm
Cathepsin L-like	Z-FR-AMC	100 mM sodium citrate, pH 5.5 (0.15 M NaCl, 1 mM DTT)	15 min at 30 °C	excitation filter 350 nm emission filter 465 nm
Cathepsin D-like	MocAc-GKPILFFRLK (Dnp)-D-R- NH ₂	100 mM sodium citrate, pH 3.5 (0.15 M NaCl, 1 mM DTT, 10µM E-64)	15 min at 30 °C	excitation filter 328 nm emission filter 393 nm
Legumain-like	Z-VAN-AMC	100 mM sodium citrate, pH 4.5 (0.15 M NaCl, 1 mM DTT)	15 min at 30 °C	excitation filter 350 nm emission filter 465 nm
Leucine aminopeptidase-like	LpNa	100 mM Tris-HCl, pH 7.5 (0.15M NaCl, 5mM MgCl ₂)	4 h at 30 °C	absorbance at 410 nm
α-Amylase-like	Starch	100 mM Tris-HCl, pH 6.0 (40 mM CaCl ₂ , 20 mM NaCl)	4 h at 30 °C	absorbance at 580 nm
Detoxification Enzymes ¹				
Esterase	1-NA	100 mM Tris-HCl, pH 7.0 (0.15M NaCl).	1 h at 30 °C	absorbance at 600 nm
Glutathione S-transferase	CDNB	100 mM Tris-HCl, pH 8.0 (0.15M NaCl, 5 mM reduced glutathione)	15 min at 30 °C	absorbance at 340 nm
P450	cytochrome c	100 mM Tris-HCl, pH 7.0 (0.15M NaCl, containing the NADPH generating system ⁵	4 h at 30 °C	absorbance at 550 nm

¹ Procedures for cathepsin B-, L- and D- and legumain-like activities adapted from Santamaría et al. (2015a); and for leucine aminopeptidase- and α-amylase-like activities and esterase, glutathione S-transferase and P450 activities from Ximénez -Embún et al. (2016). Assays were performed with 60 µl (P450), 40 µl (glutathione S-transferase), 20 µl (α-amylase-like), 10 µl (cathepsin B- and L-, legumain- and leucine aminopeptidase-like activities), 5 µl (cathepsin D-like activity) and 0.25 µl (esterase) of the mite protein extract.

² The substrates were added at a final concentration of 20 µM (cathepsin B- L-, D- and legumain-like activities), 1mM (LpNa), 0.25% starch, 0.25 mM (1-NA), (0.4 mM CDBN) and 50 µM (cytochrome c). Z-RR-AMC, Z-FR-AMC, MocAc-GKPILFFRLK(Dnp)-D-R- NH₂ and LpNa as in Table S; Z-VAN-AMC (N-carbobenzoxyl-Val-Ala-Asn-7-amido-4-methylcoumarin) from Bachem (Bubendorf, Switzerland); starch, 1-NA (1-naphthyl acetate), CDBN (1-chloro-2,4-dinitrobenzene) and cytochrome c from Sigma-Aldrich (St Luis, USA).

³ Concentrations are expressed at molarity in the reaction mixture.

⁴ AMC and MCA were used as standards for fluorescent substrates as in Table S3.1; starch and 1-naphthol were used as standards for α-amylase-like and esterase activities

⁵ The NADPH generating system consisted of 0.5 mM NADP, 2.5 mM glucose 6-phosphate and 0.3 units of glucose 6-phosphate dehydrogenase

Chapter 4

DROUGHT STRESS PROMOTES THE COLONIZATION SUCCESS OF A HERBIVOROUS MITE THAT MANIPULATES PLANT DEFENSES

4.1 Introduction

Global agriculture faces a big challenge as climate change will affect crop production in the near future. According to the IPCC (2013), temperatures will increase and there will be more periods of drought, especially in areas with a Mediterranean climate. Thus, less water will be available for irrigation and for key summer crops, like tomato, often only a suboptimal amount of water will be available. Furthermore, plant drought stress will also have consequences for herbivores and their interactions with plants (Cornelissen, 2011). Drought and high temperature conditions are often associated with herbivore outbreaks (Mattson and Haack 1987), but both positive and negative effects on herbivores have been reported depending on the severity of the stress and differing across species and across the plants they are attacking (Huberty and Denno, 2004; Cornelissen et al., 2008; White 2009; Gutbrodt et al., 2011). Yet, plant responses to a combination of abiotic (e. g. drought stress) and biotic (e. g. herbivory) stresses and its impact on herbivore performance are poorly documented (Inbar et al., 2001; Huberty and Denno, 2004; Gutbrodt et al., 2011).

When a plant detects drought, it activates a series of tolerance mechanisms. First it will close the stomata but if the stress continues it will stop growing while it may reset its metabolism (Harb et al., 2010; Hummel et al., 2010). In order to prevent desiccation, cells undergo an osmotic adjustment, increasing the amount of free sugars and free amino acids, especially proline (Hummel et al., 2010; Showler, 2013). The “plant stress hypothesis” assumes that such a plant will be of higher nutritional quality for herbivores and therefore can promote their performance (Huberty and Denno, 2004; White, 2009). In contrast, the “plant vigor hypothesis” (Huberty and Denno, 2004; Cornelissen et al., 2008) assumes that drought stress may induce the production of defense compounds (English-Loeb et al., 1997; Inbar et al., 2001) which, together with the reduction in plant growth, might have a negative effect on herbivore performance. However, herbivores appear not to be passive and may themselves manipulate the plant’s primary metabolism (Zhou et al., 2015) and secondary metabolism i.e. defenses (Kant et al., 2015) to their own benefit.

Plant stress responses are regulated by a complex network of phytohormones, with Jasmonic acid (JA) and Salicylic acid (SA) as the central players assisted by ancillary hormones such as Absciscic acid (ABA), auxins and ethylene. The response to herbivory is predominantly regulated by the JA pathway (Howe and Jander, 2008). This pathway generates the active component Jasmonic Isoleucine (JA-Ile) via oxophytodienoic acid (OPDA) and JA (Schuman and Baldwin, 2016). In contrast, SA is the key regulator of defense responses induced by

phloem feeding insects and biotrophic pathogens (Howe and Jander, 2008). Finally, drought stress gives rise to accumulation of ABA, which regulates processes like stomatal closure (Verma et al., 2016). Phytohormones can crosstalk and thereby modulate each other's actions. JA and SA pathways are often antagonistic and some herbivores abuse this to manipulate plant defenses (Howe and Jander, 2008; Schuman and Baldwin, 2016). The herbivorous spider mites *Tetranychus evansi* and some strains of *T. urticae*, in contrast, have been reported to suppress plant defenses downstream of JA and SA and independent of crosstalk when feeding on tomato (Alba et al., 2015). This defense suppression could be attributed to salivary effector-proteins secreted into plants during feeding (Villarroel et al., 2016). Also abiotic (drought) stress can change the plant's hormonal balances during herbivory (Atkinson et al., 2015) as ABA was reported to suppress the SA pathway while inducing the JA pathway (Thaler and Bostock, 2004), although its effects can be varying and operate indirectly via other hormones (Verma et al., 2016).

The Tomato Russet Mite (TRM), *Aculops lycopersici* (Massee) is a cosmopolitan pest on solanaceous crops, mainly on tomato (*Solanum lycopersicum*). TRM causes massive yield losses of tomato (Duso et al., 2010), one of the most important horticultural crops worldwide. TRM is an eriophioid mite, a family that includes the world's smallest terrestrial animals (Keifer, 1946; Sabelis and Bruin, 1996) and, therefore, are often detected too late by growers. It has a short life completing its life-cycle within 7 days, depending on the temperature (Haque and Kaway, 2003). Once detected, TRM is difficult to control since it hides in the forest of tomato leaf hairs (trichomes) that protects it from predators (van Houten et al., 2013). Early studies on the TRM-tomato interaction reported that TRM induces accumulation of oxidative enzymes like peroxidases (POD) but not protease inhibitors (PI) or polyphenol oxidases (PPO) when feeding on the plant (Stout et al., 1996; Petanovic and Kielkiewicz, 2010). Recently, Glas et al. (2014) described that TRM manipulates tomato plant defenses by suppressing JA-defenses, but not SA-defenses, downstream of phytohormone accumulation and independent from JA-SA antagonistic crosstalk. TRM outbreaks might be promoted directly by climate change as it's optimal growth conditions are at 27 °C and 30% relative humidity (Duso et al., 2010). However, we were primarily interested in how drought affects herbivores that suppress plant defenses. Recently we showed that *T. evansi*, which suppresses both JA and SA defenses simultaneously (Alba et al., 2015), is promoted on plants under mild and moderate watering regimes probably due to increased levels in free sugars and essential amino acids, indicating that indirect plant-mediated effects independent from defenses may promote population growth

of this mite (Ximénez-Embún et al., 2016). For the current study we used TRM since it selectively suppresses only JA-defenses (Glas et al., 2014) while, in theory, increased ABA could modulate JA-defenses beyond the control of the mite (Golldack et al., 2014).

The overall aim of this study was to assess the extent to which drought affects TRM-induced changes in the primary and secondary metabolism of plants. Moreover, we aimed to assess the extent to which their combination affects the physiological status (i.e. nutritional value and chemical defenses) of tomato in order to estimate the magnitude of the interaction between these two stresses. This information is a useful instrument for predicting outbreaks of pests that manipulate plant resistance during a changing climate.

4.2 Materials and methods

4.2.1 Plant material and mite rearing

Tomato (cv. Moneymaker) seeds were germinated in soil (commercial peat) and transferred to 0.66 L pots (diameter: 12 cm) filled completely with the same soil and grown in a greenhouse at 25 °C and a 15h light /9 h dark regime.

A tomato russet mite population collected in summer 2008 from a greenhouse in the Westland area (The Netherlands) was supplied by Koppert Biological System (Berkel en Rodenrijs, The Netherlands). They were reared in insect cages (Bug-Dorm-44590DH, Bug Dorm Store, MegaView Science, Taichung, Taiwan) on 3-5 week old tomato plants (cv Castelmart), and maintained in a climate room at 27 °C/25 °C day/night under a 16h light /8 h dark regime at 60 % Relative Humidity (RH).

4.2.2 Drought stress regime

Drought stress was attained using the protocol described by Ximénez-Embún et al. (2016) with minor modifications. In brief, tomato plants were maintained well-watered in the greenhouse as described previously until they had developed three expanded leaves (in about 3 weeks). Then plants were transferred to a climate room (same conditions as above) and were randomly divided in two groups: one group for control and the other for the moderate drought stress treatment. Control plants were watered every two to three days to maintain the soil volumetric water content (Θ) up to 74 %. For moderate stress, watering was stopped for seven days and thereafter plants were watered to maintain Θ between 21 and 30 %. The wilting point was avoided as it happens at Θ : 16 %. Θ was determined gravimetrically by recording single plant pot weight.

The severity of drought stress was assessed by measuring the stomatal conductance (gs) of the sub-terminal leaflet of the third leaf using a leaf porometer (SC-1 Decagon-T, Pullman, USA). Plant growth was estimated by measuring the stem length (distance between the soil and the terminal bud).

4.2.3 Bioassays

Two different experiments were carried out: the first to measure the effect of drought on mite population growth, the second to obtain plant material for the different metabolite analyses and to evaluate plant damage. Both experiments were carried out in a climate room under the same environmental conditions as used for the mite rearing.

Mite population growth. Tomato plants were assigned to four different groups: two served as controls and two were used for the drought stress treatment. For each, one group was sampled at 7 and the other at 14 days post infestation (dpi). When drought stress conditions had stabilized (at about 7-9 days after stopping irrigation for Moneymaker in our experimental conditions: see Ximénez-Embún et al., 2016), this moment coincided with the plants having four expanded leaves), 12 plants per group were infested with TRM by placing 20 individual adult mites on leaf 2, 3 and 4 using a fine-bristle paint brush. Thus, plants were inoculated with a total 60 TRM each. A thin barrier of Lanolin (Sigma_aldrichChemie, Zwijndrecht, The Netherlands) was prepared, using a needleless syringe, around the petioles of the leaf to prevent the mites from escaping. The TRM population density was assessed at the two time points using the protocol described by Glas et al. (2014). In brief, the three infested leaves of each plant were detached and washed one by one for 20 seconds in a single volume of 25 ml of Ethanol 100%. TRM were counted by running 2 ml of leaf washes through a particle count system (PAMAS SVSS, PAMAS, Rutesheim, Germany). The number of particles measured was in the range of 50 to 200 μm as TRM adults size is around 120-150 μm .

Plant sampling and plant damage evaluation. Tomato plants were divided in four different groups combining two treatments: drought stress (control or drought) and TRM infestation (infested or non-infested). When drought stress conditions had stabilized (see “mite population growth” section), plants were infested with TRM by placing a small piece of an infested leaf (ca. 0.5 cm^2) on each of the two subterminal leaflet on leaves 2, 3 and 4, resulting in 6 pieces per plant. These pieces were cut from highly infested tomato plants and each piece contained around 200-250 mobile stages as determined by stereomicroscope (1200 - 1500 mites per plant). A thin barrier of Lanolin was prepared, using a needleless syringe, around the

petioles of the leaf to prevent the mites from escaping. Seven days after infestation plant material was collected by pooling the subterminal leaflets of leaves 2, 3 and 4 of each plant. Samples were flash frozen in liquid nitrogen, then ground in liquid nitrogen using a mortar and pestle to a fine powder and stored at -72 °C. Six plants were sampled for each of the four treatments, except for phytohormone analysis for which eleven plants were sampled per treatment.

To determine the damage produced by TRM on tomato, a plant injury index was established between 0 (healthy leaf) and 5 (dead leaf) as described in Figure S4.1. The plant injury index was assessed at 7 dpi, and at 10 dpi and was averaged for three infested leaves.

4.2.4 Chemical and biochemical analysis

Chemicals and Equipment. Unless specified otherwise, all chemical compounds were obtained from Sigma-Aldrich (St Luis, USA). Fluorimetric measurements were made using a Varioskan Flash reader (ThermoFisher Scientific, Willmington, USA), and spectrophotometric measurements with a VERSAmaxmicroplate reader (Molecular Devices Corp., Sunnyvale, USA).

Free sugars. Samples of 40 mg of frozen leaf powder were dried in an oven at 70 °C for three days and 2.5 mg of the resulting material was homogenized in 650 µl of ethanol 95% (v/v), heated at 80 °C for 20 min, centrifuged at 10,000 rpm for 10 min, and the supernatant collected. The process was repeated two more times and the three supernatants were pooled. A volume of 750 µl of the mixture was dried on a SpeedVac Concentrator Savant SVC-100H (ThermoFisher scientific, Willmington, DE, USA) and redissolved in 500 µl of water. Soluble carbohydrate concentration was estimated by the anthrone method (Maness, 2010) using glucose as a standard. In brief, 1 ml of anthrone reagent (0.2% v/v anthrone on 95% sulfuric acid) was added to the extract, heated at 90 °C for 15 min, and the absorbance measured at 630 nm.

Free amino acids. The extraction of the free amino acids was done as described by Hacham et al. (2002). Samples of 50 mg of frozen leaf powder were homogenized with 600 µl of water:chloroform:methanol (3:5:12 v/v/v). After centrifugation at 12,000 rpm for 2 min, the supernatant was collected and the residue was re-extracted with 600 µl of the same mixture, pooling the two supernatants. A mixture of 300 µl of chloroform and 450 µl of water were added to the supernatants, and after centrifugation the upper water:methanol phase was collected and dried in the SpeedVac. The samples were dissolved on 100 µl of sodium citrate

loading buffer pH 2.2 (Biochrom, USA) and 10 µl were injected on a Biochrom 30 Amino Acid Analyser (Biochrom, USA) at the Protein Chemistry Service at CIB (CSIC, Madrid, Spain).

Soluble Protein. Samples of 100 mg of leaf frozen powder were homogenized in 500 µl of 0.15M NaCl, ground with fine sand. The homogenate was centrifuged at 12,000 rpm for 5 min at 4 °C, and the soluble protein quantified by absorbance at 280 nm on a Nanodrop 1000 spectrophotometer (ThermoFisher scientific, Willmington, USA).

Plant defense proteins. Samples of 100 mg of leaf frozen powder were homogenized with 500 µl of extraction buffer (0.15 M NaCl for protease inhibitors, and 0.1 M phosphate buffer, pH 7.0; 5% w:v polyvinylpolypyrrolidone for oxidative enzymes) and soluble protein quantified as explained above.

Protease inhibitors. The inhibitory activity of plant protein extracts was tested against commercial enzymes: papain (EC 3.4.22.2), cathepsin B from bovine spleen (EC 3.4.22.1), trypsin from bovine pancreas (EC 3.4.21.4), α-chymotrypsin from bovine pancreas (EC 3.4.21.1), cathepsin D from bovine spleen (EC 3.4.23.5) and leucine aminopeptidase from porcine pancreas (EC 3.4.11.1), as described by Ximénez-Embún et al. (2016). In brief, samples of 20 µg of plant protein extracts (40 µg in case of leucine aminopeptidase inhibition assay) were preincubated for 10 min with 100 ng of the commercial enzyme, subsequently substrate is added and incubated for a specific time and absorbance is measured. Reaction conditions are summarized in Table S4.1. Results were expressed as a percentage of protease activity inhibited.

Oxidative enzymes. Polyphenol oxidase (PPO) activity was analyzed by incubating 20 µl of enzyme extract with catechol (40 mM final concentration) in 160 µl of Tris-HCl pH 8.5 buffer at 30 °C for 1 h. Absorbance was read at 420 nm. Peroxidase (POD) activity was determined incubating 20 µl of a 1:10 dilution of the enzyme extract with guaiacol (5 mM final concentration) and H₂O₂ (2.5 mM final concentration) in 150 µl of potassium phosphate pH 6 buffer at 30 °C for 10 min. Absorbance was read at 470 nm. PPO and POD activities were expressed as nmol substrate metabolized relative to time and total protein content.

4.2.5 Quantification of phytohormones by means of LC-MS

Phytohormones were extracted adapting the procedure described by Alba et al. (2015). In brief 100 mg of frozen leaf powder was homogenized using a GenoGrinder (Precellys24 Tissue Homogenizer, Bertin Technologies, Aix-en-Provence, France) in 1 ml of ethyl acetate spiked with 100 ng of D₆-SA and D₅-JA (C/D/N Isotopes Inc., Pointe-Claire, Quebec, Canada) as internal standards. After centrifugation for 20 minutes at 13,000 rpm and 4 °C, the

supernatant was collected and the residue was re-extracted with 0.5 ml of ethyl acetate without internal standards. After centrifugation the supernatant was combined with the previous one and evaporated on a vacuum concentrator (CentriVap Centrifugal Concentrator, Labconco, Kansas City, MO, USA) at 30 °C. The pellet was re-suspended in 250 µL of 70% LCMS-grade methanol (v/v), centrifuged for 10 minutes and transferred to liquid chromatography-mass spectrometry (LC-MS) vials (Fisher Scientific, Hampton, NH, USA). A serial dilution of pure standards of OPDA, JA, JA-Ile, SA and ABA was run separately. Phytohormone measurements were conducted on a liquid chromatography tandem mass spectrometry system (Varian 320 Triple Quad LC/MS/MS, Agilent Technologies, Santa Clara, CA, USA). The mobile phase comprised solvent A (0.05% formic acid in water; Sigma-Aldrich, Zwijndrecht, the Netherlands) and solvent B (0.05% formic acid in methanol; Sigma-Aldrich). The program was set as follows: 95% solvent A for 1 minute 30 seconds, followed by 6 minutes in which solvent B increased till 98% which continued for 5 minutes, subsequently returning to 95% solvent A for 1 minute until the end of the run (18 minutes in total). The flow was 0.2 ml/min. during all the run. Compounds were detected in the electrospray ionization negative mode. The parent ions, daughter ions used for these analyses are listed in Supporting information Table S2. For all oxylipins and ABA we used D₅-JA to estimate the recovery rate and D₆-SA for SA. Their *in planta* concentrations were subsequently quantified using the external standard series. Phytohormone amounts were expressed as ng per gram of fresh leaf material.

4.2.6 Quantification of gene expression via qRT-PCR

Samples of 100 mg of frozen leaf powder were taken for total RNA extraction using the hot phenol method (Verwoerd et al., 1989). The integrity of RNA was checked on 1% agarose gels and subsequently quantified using a NanoDrop 100 spectrophotometer. DNA was removed with DNase (Ambion, Huntingdon, UK) according to the manufacturer's instructions, after which a control PCR was carried out to confirm the absence of genomic DNA contamination. cDNA was synthesized from 2 µg total RNA using a poly(dT) primer and M-MuLV Reverse Transcriptase (Fermentas, St. Leon-Rot, Germany) according to the manufacturer's instructions. cDNA dilutions (1:10) were used as the template in quantitative reverse-transcriptase PCR (qRT-PCR). Reactions were carried out in a total volume of 20 µl containing 0.25 µM of each primer, 0.1 µl ROX reference dye and 1 µl of cDNA template. Two technical replicates were performed per measurement. qRT-PCR was performed with Platim SYBR Green qPCR Super Mix (Invitrogen, Paisley, UK) using an ABI 7500 (Applied Biosystems, Foster City, CA, USA) system. The program was set to 2 minutes at 50 °C, 10 minutes at 95

°C, 45 cycles of 15 seconds at 95 °C and 1 minute at 60 °C, followed by a melting curve analysis. Target gene expression levels were normalized to those of actin. The normalized relative quantity (NRQ) data were calculated by the ΔC_t method $NRQ = (PE_{\text{target}}^{C_{t_target}})/(PE_{\text{reference}}^{C_{t_reference}})$ (PE = primer efficiency; Ct = cycle threshold). The PEs were determined by fitting a linear regression line on the Ct-values of a standard cDNA dilution series. Specific amplification was ensured by melting curve analyses and generated amplicons were sequenced. The primers used are listed in Table S4.3.

4.2.7 Statistical analysis

All plant and mite data were checked for the assumptions of normality and heteroscedasticity, and transformed if necessary. Stem length and stomatal conductance were $\log_{10}(x)$ transformed and analyzed using a three-way ANOVA (using as fixed factors drought treatment, TRM infestation and time) performing a Bonferroni *post hoc* test to compare drought-stress treatments within each time. The TRM population size was $\log_{10}(x)$ transformed and analyzed by a two-way ANOVA (drought treatment, time). The significant differences in leaf damage index in control versus moderate drought plants were determined by the non-parametric Mann-Whitney-Wilcoxon test (U-test). The percentage of protein, free amino acids, free sugars and protease inhibition were arcsine square root transformed, phytohormone data were $\log_{10}(x)$ transformed and gene-expression data (NRQ values) were $\ln(x)$ transformed. These data and the oxidative enzyme activities were analyzed using a two-way ANOVA using as fixed factors drought treatment and TRM infestation. A Newman-Keuls *post hoc* test was performed to see differences of mean between treatments, except for gene expression data where Fisher LSD (Least significant difference) *post hoc* tests were performed.

4.3 Results

4.3.1 Effects of drought on *Aculops lycopersici* population growth and plant damage.

When feeding on drought-stressed tomato plants the TRM population grew faster than on control plants (Figure 4.1a). However, the difference in population size was significant only at 14 days dpi. Likewise, plant damage was significantly higher on stressed plants from 10 dpi onwards (Figure 4.1b).

Figure 4.1 Population growth (A) of *Aculops lycopersici* (TRM) and plant injury (B) on moderately drought stressed tomato plants. Plants were infested with 60 individuals for the population growth assay (A) and 1250 individuals for the leaf damage index assay (B). Population size was measured at 7 and 14 days post infestation (dpi) while plant damage index was measured at 7 and 10 dpi. Data points represent the mean $\pm$ SE. An asterisk indicates a significant differences between drought treatments (population assay: Two-way ANOVA, Bonferroni *post hoc* test, $p < 0.05$; Injure index: Mann-Whitney-Wilcoxon test).

4.3.2 Effects of drought on stomatal conductance and tomato plant growth.

The impact of drought stress on stomatal conductance was similar in the two experiments, and it was between 3 and 8 times smaller for stressed plants than for control plants (Figure S4.2 a). We observed a reduction in stomatal conductance for the control plants during the course of the experiment. Moderate drought also affected plant growth, as stem length was significantly shorter for stressed plants than for control plants at all measured points (Figure S4.2 b). The TRM infestation didn't induce a significant effect on stomatal conductance and tomato plant growth. Therefore, data from infested and uninfested plants were pooled within each experiment for producing Figure S4.2.

4.3.3 Changes on plant nutritional composition induced by drought and TRM.

The levels of nutrients (protein, free sugars and amino acids) in tomato leaves changed in response to drought stress as well as to TRM infestation (Table 4.1 and Figure 4.2). TRM induced a significant increase of the amount of total protein. The amino acids analyzed were separated as essential and non-essential ones according to the division made by Rodriguez and Hampton (1966) for the phytophagous mite *T. urticae*. Moderate drought induced a reduction of the amount of the non-essential amino acids aspartic acid and alanine but increased the amount of the essential histidine. Proline was the only non-essential amino acid that accumulated to higher levels in response to both TRM infestation and TRM combined with drought stress, whereas alanine levels were also reduced when both stresses were combined. In contrast, TRM alone and combined with drought stress induced the essential ones, valine, isoleucine, tyrosine, lysine and leucine, the latter at four times higher levels than in the control. The essential amino acid arginine was induced by TRM but not when combined with drought stress. Alone drought or TRM did not affect free sugar levels but combined they significantly increased these levels.

Table 4.1 Effect of moderate drought and infestation by *A. lycopersici* on nutritional composition of tomato leaves.

	Non-infested		TRM infestation	
	Control	Moderate drought	Control	Moderate drought
Protein	17±2 a	18±1 a	27±2 b	27±3 b
Free aminoacids	0.37±0.05 a	0.27±0.02 a	0.44±0.09 a	0.28±0.03 a
Free sugar	4.8±1.0 b	5.2±0.5 b	3.8±0.2 b	8.1±1.1 a

Data are mean ±SE of % dry weight. Different lower case letters within rows indicates significant differences (Two-way ANOVA, Newman-Keuls *post hoc* test at $p<0.05$).

Figure 4.2 Levels of free amino acids in tomato leaves from plants under moderate drought stress and/or infested with *Aculops lycopersici* (TRM) at 7 days post infestation (dpi). The bars represent the mean amount amino acid (μg) per gram of dry weight (DW) $\pm$ SE represented on a logarithmic scale. The division between essential and non-essential amino acids for *T. urticae* is based on Rodriguez and Hampton (1966). Different letters indicate significant difference between treatments (Two-way ANOVA, Newman-Keuls *post hoc* test, $p < 0.05$).

4.3.4 Effect of drought and TRM on tomato plant defense: phytohormones, defense genes and defense proteins.

The defense response of the plant was analyzed at three levels: phytohormone accumulation (Figure 4.3), the transcript levels of marker genes linked to the JA and SA pathways (Figure 4.4) and the activity of defense proteins (Table 4.2).

TRM induced a two-fold increase in the accumulation of JA, a 2.5-fold increase in OPDA (the precursor of JA) and an 8.7-fold increase in Ja-Ile (the bioactive form of JA) in tomato plants. In addition, TRM induced a fourfold increase in SA accumulation. When combining TRM infestation with drought stress the amount of OPDA increased 4-fold and those of SA 12-fold. However, while TRM alone induced JA and JA-Ile accumulation significantly, combining it with drought antagonized this effect. Moderate drought affected ABA

accumulation significantly ($P = 0.007$ in the two-way ANOVA), but the *post hoc* analysis did not detect significant differences among treatments.

Figure 4.3 Phytohormone levels in tomato leaves under moderate drought stress and/or infested with *Aculops lycopersici* (TRM) at 7 days post infestation. The bars represent the mean ng of phytohormone per gram of fresh weight (FW) $\pm$ SE of endogenous OPDA (A), JA (B), JA-Ile (C), ABA (D), SA (E). Different letters in each figure indicate significant difference between treatments (Two-way ANOVA, Newman-Keuls *post hoc* test, $p < 0.05$).

TRM infestation did not significantly induce the expression of the JA-marker genes but it did elevate transcript accumulation for the SA-marker gene *PR-P6* more than 20 times. Moderate drought, in contrast, down regulated the expression of the JA marker genes *JIP-21* and *PPO-F* by 2-fold and 4-fold respectively. When combining drought and TRM, expression levels decrease even further albeit not to significantly lower levels.

The defense proteins were divided in two categories: protease inhibitors and oxidative enzymes. From the different protease inhibitors, TRM infestation increased the total activity of cysteine protease (cathepsin B and papain) inhibitors. Drought decreased the total inhibitory activity of serine protease (trypsin and chymotrypsin) inhibitors. Cathepsin D and aminopeptidase inhibitory activities were not affected by any of the treatments. The total activity of the oxidative enzymes PPO and POD was significantly increased by TRM. Drought

stress decreased the POD activity, but when combined with TRM infestation PPO and POD activities were equal to those of the control.

Figure 4.4 Relative transcript abundance of tomato plants under moderate drought stress and/or infested with *Aculops lycopersici* (TRM) at 7 days post infestation. Selected genes mark the JA pathway: *TD-II* (A), *PPO-F* (B), *JIP-21* (C) and *PI-IIf* (D) and the SA pathway: *PR-P6* (E). The bars represent the mean normalized relative quantity (NRQ) $\pm$ SE. Different letters indicate significant difference between treatments (Two-way ANOVA, Fisher LSD *post hoc* test, $p < 0.05$).

Table 4.2 Effect of moderate drought and infestation with *Aculops lycopersici* on tomato plant defense proteins at 7 days post infestation.

	Non-infested				TRM infestation			
	Control		Drought		Control		Drought	
Cathepsin B	54±2	b	47±5	b	82±2	a	86±2	a
Papain	56±3	c	57±3	c	72±4	b	83±2	a
Cathepsin D	43±1	a	39±3	a	40±4	a	35±3	a
Trypsin	40±4	ab	23±5	b	50±10	a	35±6	ab
Chymotrypsin	91±2	a	45±12	b	95±2	a	75±8	a
Aminopeptidase	14±3	a	10±3	a	16±4	a	19±4	a
Poliphenol oxidases¹	5.2±0.5	a	4.0±0.6	a	7.8±1.3	b	4.5±0.4	a
Peroxidases²	9.4±2.4	b	3.9±0.9	c	29.9±6.4	a	12.1±2.6	b

Data are mean ± SE. Different lower case letters within rows indicates significant differences (Two-way ANOVA, Newman-Keuls *post hoc* test at $p < 0.05$).

(1) PPO: nmol Cathecol metabolized/ mg Protein*min

(2) POD: nmol Guaiacol metabolized/ mg Protein*min.

4.4 Discussion

We showed that drought-stressed tomato plants reconfigure their metabolism (Bauer et al., 1997; English-Loeb et al., 1997) and become a better host for TRM. This finding is in agreement with the “plant stress hypothesis” that assumes that stress, such as drought stress, promotes herbivore performance due to an improvement of the nutritional value of the stressed plant (Huberty and Denno, 2004; White, 2009). Interestingly, drought and TRM synergized accumulation of free sugars which have been shown to act as a phagoestimulant (Wermenlinger et al., 1991; Showler, 2013). Also in a previous study we observed that drought-induced accumulation of free sugars and amino acid in tomato plants coincides with improved performance of the herbivorous mite *T. evansi* (Ximénez-Embún et al., 2016). In contrast, here we did not observed a reduction in proteins, indicating that there was not a mobilization of protein into free amino acids, as has been observed in other plant-herbivore interactions previously (Bauer et al., 1997; Ximénez-Embún et al., 2016). Accordingly, the amounts of most amino acids remained unaltered in drought stresses plants, except for a reduction of aspartic acid and alanine contents and an increase of histidine levels. Proline accumulation, an indicator of drought stress (Claussen 2005; Showler 2013), was not affected by our drought stress treatment. Leaf age or a low level of light radiation may explain the absence of a proline response, since we also observed a reduction in stomatal conductance and in plant growth which

clearly indicates that the drought treatment did affect the plants (Harb et al., 2010; Hummel et al., 2010).

The first important aspect that determines plant palatability is its nutritional composition. Reconfiguration of the plant's primary metabolism during herbivory (Zhou et al., 2015), i.e. mainly that of free carbohydrates and amino acids, has been reported to affect the pool of precursors for defense compounds, the amount of available energy for the plant and may indicate reallocation of nutrients e.g. to storage tissues (Steinbrenner et al., 2011; Zhou et al., 2015). Our data show that TRM infestation does not significantly affect the accumulation of free sugars and total free amino acids. In contrast, we observed increased accumulation of total protein. Something similar was observed for tomato plants infested with the generalist leaf feeder *Helicoverpa zea* (Steinbrenner et al., 2011). Nevertheless, the concentrations of several essential amino acids increased during TRM infestation significantly, with isoleucine and leucine levels being elevated the strongest. Possibly this is causally related to the increase in the JA conjugate JA-Ile which plays a decisive role in plant defense signalling (Ataide et al., 2016; Schuman and Baldwin, 2016). Proline, a non-essential amino acid, was also induced by TRM reminiscent of *Manduca sexta* on tomato (Gomez et al., 2012) and the pea aphid on *Medicago truncatula* (Guo et al., 2013). Proline can be used by mosquitoes as a direct energy substrate for glycolysis and ATP production (Scaraffia and Wells, 2003) and it would be interesting to test its effect on TRM population growth.

The second important aspect that determines plant palatability and mite population growth is plant resistance as determined by its defense responses. As mentioned before, we investigated plant defense at three levels or organization: at the phytohormone-level, the marker-gene expression level and at the defense-protein activity level. We found that TRM elevated the accumulation of JA-precursor OPDA, JA, the JA-derivative JA-Ile and of SA. However, in response to TRM only the transcript levels of the SA-marker gene (*PR-P6*) increased but not those of the JA-marker genes *PI-IIf*, *JIP-21*, *PPO-F* and *TD-II*. This is in line with the results of Glas et al. (2014) who showed that TRM suppresses selectively the expression of JA-dependent defense genes downstream of phytohormone accumulation. *JIP-21* and *PI-IIf* are serine protease inhibitors and, accordingly, also the plant's protease inhibition activity against trypsin and chymotrypsin was not induced by TRM. In contrast, TRM it induced an increase in the activities of cysteine (papain and cathepsin B) protease inhibitors and of the oxidative enzymes PPO and POD, similar to *T. evansi* (Ximénez-Embún et al., 2016). So far, the phytophagous mites analysed belonging to the family Tetranychidae (*T. urticae* and *T.*

evansi) and Tenuipalpidae (*Brevipalpus chilensis*) rely on cysteine proteases for digestion (Carrillo et al., 2011; Ximénez-Embún et al., 2016). In addition, cysteine protease inhibitors have been shown to be detrimental to tetranychid mites (Santamaria et al., 2015b). However, there is no information for digestive physiology for TRM. This raises an interesting question: while it was shown that suppression of defenses - i.e. JA- and SA-defenses by *T. evansi* and JA-defenses by *A. lycopersici* – it beneficial for mites and decreases plant resistance yet all three mite species – including the inducer of JA and SA defenses *T. urticae* – induce accumulation of cysteine protease inhibitors. Hence it would be interesting to see if these mites have a different sensitivity to this type of protease inhibitor or if the role of cysteine protease inhibitors within the collective defense response of the plant is minor. Tomato plants silenced for this class of inhibitors could be instrumental for answering this question.

The hormonal responses in plant when experiencing a mixture of biotic and abiotic stresses are complex as many of these pathways interact via cross-talk (Thaler and Bostock, 2004; Atkinson et al., 2015). The accumulation of hormones is dynamic in time and space and the quantities one measures inside of a single leaf or leaflet will depend on several factors such as the time of day (Atamian and Harmer, 2016), on where the stress started (i.e. is it a local or a systemic stress?) and on when it started (Eckardt, 2015). For instance ABA accumulation typically peaks during the first phase of a drought response but ceases again while the stress continues (Thompson et al., 2007). This might explain why we didn't observed differences in ABA accumulation between treatments at the 14 dpi. Moreover, also JA and SA are involved in the early drought response of tomato (Muñoz-Espinosa et al., 2015), though there is not much information on their role in the later phases of the response. In our experiments, drought stress didn't induce JA or SA. However, when drought stressed plants were also infested with TRM the accumulation of SA increased to almost threefold compared to the levels induced by the mite on control plants. This synergistic effect may be explained by hormonal crosstalk between JA and SA (Howe and Jander, 2008; Schuman and Baldwin, 2016) since the combination of TRM and drought reduces the levels of JA with a similar trend for JA-Ile. Hence the effects we observed on JA and SA may be causally linked. Although the effect of drought stress alone on JA and JA-Ile was not significant, it decreased the transcript levels of the JA-marker genes *JIP-21* and *PPO-F*, and reduced the activity of serine protease (trypsin and chymotrypsin) inhibition, and of POD. This indicates that the inducibility of JA-related responses by herbivores is strongly reduced in drought stressed tomatoes. This leads us to suggest that, under our conditions, the reduction of tomato defenses due to drought stress contributes to the increase

in the plant's palatability to TRM. This implies that for TRM effects of drought on defenses may play a bigger role in promoting the mite's population growth-response than for the spider mite *T. evansi* for which especially changes in nutritional quality were associated with an increase in the mite's performance (Ximénez-Embún et al., 2016). However, in literature results on the effect of drought stress on tomato plant defenses are not always consistent (English-Loeb et al., 1997), possibly as a consequence of the spatiotemporal dynamics of (cross-talking) hormonal responses in relation to differences between sampling protocols and experimental conditions. Therefore, the most important readout is the performance of the herbivore since clearly not only spider mites (Ximénez-Embún et al., 2016) but also russet mites benefit from drought and become a more aggressive pest on tomato although the plant-physiological basis for these effects can be different. We feel that altering the nutritional quality of tomatoes under drought stress through breeding may be more challenging than breeding for resistances that perform well also under stressful conditions. Therefore we feel that identifying "stress-resistant" defenses will be necessary to ensure crop protection also under a changing climate in times that pesticides are increasingly banned from the production process.

4.5 Conclusions

Our data indicate that the joint action of both drought stress and tomato russet mite infestation can have additive, synergistic and antagonistic effects on the plant's nutritional quality and defense responses. These effects are characterized by a disproportional increase in the plant's levels of free sugars, and some essential amino acids and also by a weakened JA-response paralleled by an amplified SA-response. The predicted increase in the number, the duration and the extent of future periods of drought due to climate change will therefore likely promote TRM performance. Hence one may expect TRM to become a more problematic pest in tomato than it already is. To be ahead of this danger we should search for drought-stress resistant defenses that can be used to protect tomato cultivation also for future generations.

4.6 Supporting information

Figure S4.1 Leaf damage index explanation

Figure S4.2 Effect of moderate drought on A) tomato stomatal conductance (gs) and B) stem length at mite infestation (mi) at 7 and 14 days post infestation (dpi). Data (mean $\pm$ SE) are average of the values on infested and non-infested plants on TRM population growth and plant material experiments, as mite infestation didn't show a significant effect. An asterisk indicates a significant difference between drought and control treatments at each time (Three-way ANOVA, Bonferroni *post hoc* test, $p < 0.05$).

Table S4.1 Summary of analytical methods to assess the inhibitory activity of plant protein extracts ¹

Commercial enzyme ²	Substrate ³	Buffer ⁴	Incubation	Measurement ⁵
Cathepsin B from bovine spleen (EC 3.4.22.1)	Z-RR-AMC	100 mM NA phosphate, pH 6.0 (10 mM L-cysteine, 10 mM EDTA, 0.01% (v/v) Brij 35)	1 h at 28 °C	excitation filter 350 nm emission filter 465 nm
Papain (EC 3.4.22.2)	Z-FR-AMC	100 mM Na phosphate, pH 6.0 (10 mM L-cysteine, 10 mM EDTA, 0.01% (v/v) Brij 35)	1 h at 28 °C	excitation filter 350 nm emission filter 465 nm
Cathepsin D from bovine spleen (EC 3.4.23.5)	MocAc-GKPILFFRLK (Dnp)-D-R- NH ₂	100 mM sodium citrate, pH 3.5 (0.15M NaCl, 5 mM MgCl ₂)	10 min at 30 °C	excitation filter 328 nm emission filter 393 nm
Trypsin from bovine pancreas (EC 3.4.21.4),	Z-LA-AMC	100 mM Tris-HCl, pH 7.5 (0.15M NaCl, 5 mM MgCl ₂)	1 h at 35 °C	excitation filter 350 nm emission filter 465 nm
α -Chymotrypsin from bovine pancreas (EC 3.4.21.1),	SucAAPF-AMC	100 mM Tris-HCl, pH 7.5 (0.15M NaCl, 5 mM MgCl ₂)	1 h at 35 °C	excitation filter 350 nm emission filter 465 nm
Leucine aminopeptidase from porcine pancreas (EC 3.4.11.1).	LpNa	100 mM Tris-HCl, pH 8 (0.15M NaCl, 5 mM MgCl ₂)	1 h at 30 °C	absorbance at 410 nm

¹ Procedures adapted from Ximénez-Embún et al. (2016). Samples of 20 µg of plant protein extracts (40 µg in case of leucine aminopeptidase) were preincubated for 10 min with 100 ng of the commercial enzyme.

² All purchased from Sigma-Aldrich (St Luis, USA).

³ The substrates were added at a final concentration of 20 µM. Z-RR-AMC (N-carbobenzoxycloxy-Arg-Arg-7-amido-4-methylcoumarin) for cathepsin B, Z-FR-AMC (N-carbobenzoxycloxy-Phe-Arg-7-amido-4-methylcoumarin) for papain, Z-LA-AMC (Z-L-Arg-7-amido-4-methylcoumarin) for trypsin, SucAAPF-AMC (Suc-Ala-Ala-Pro-Phe-7-amido-4-methylcoumarin) for chymotrypsin, all purchased from Calbiochem (MerkMilipore, Billerica, USA), MocAc-GKPILFFRLK(Dnp)-D-R- NH₂ from Peptanova (Germany) for cathepsin D, and LpNa (L-leucine p-nitroanilide) from Sigma-Aldrich (St Luis, USA) for leucine aminopeptidase.

⁴ Concentrations are expressed at molarity in the reaction mixture.

⁵ AMC (7-amino-4-methylcoumarin) (Bachem, Switzerland) as standard for all fluorescent substrates, except MCA (MoCAC-Pro-Leu-Gly) (Peptanova GmbH, Germany) for cathepsin D. Double blanks were used to account for spontaneous breakdown of substrates and the plant protease activity, and all assays were done in duplicate.

Table S4.2 Parameters used for detection of phytohormones and related compounds by LC-MS/MS

Compound	Molecular ion [M-H] (<i>m/z</i>)	Fragment ion (<i>m/z</i>)
OPDA	291	165
JA	209	59
D₅-JA (Internal Standard)	213	61
JA-Ile	322	130
SA	137	93
D₆-SA (Internal Standard)	141	97
ABA	263	153

Table S4.3 Nucleotide sequence of primers used for qRT-PCR analysis

Target Gene	Name	GenBank (GB) accession	Gen Model ITAG2.3:	Forward Primer 5' → 3'	Reverse Primer 5' → 3'
PPO-F	<i>Polyphenol-oxidase-F</i>	AK247126.1	Solyc08g074630.1.1	CGGAGTTTGCAGGGAGTTATAC	TTGATCTCCACACTTTCAATG
JIP-21	<i>Jasmonate-inducible protein 21</i>	AJ295638.1	Solyc03g098790.1.1	ACTCGTCCTGTGCTTTGTCC	CCCAAGAGGATTTTCGTTGA
TD2	<i>Threonine Deaminase-2</i>	M61915.1	Solyc09g008670.2.1	TGCCGTATAAAATGTCACCA	ACTGGCGATGCCAAAATATC
PI-II_f	<i>Proteinase Inhibitor II_f</i>	AY129402.1	Solyc03g020080.2.1	GACAAGGTACTAGTAATCAAT TATCC	GGGCATATCCCGAACCCAAGA
PR-P6	<i>Pathogenesis-related protein P6</i>	M69248.1	Solyc00g174340.1.1	GTACTGCATCTTCTTGTTTCCA	TAGATAAGTGCTTGATGTCCA
Actin	<i>Actin</i>	XM_004235020.1	Solyc03g078400.2.1	TTAGCACCTTCCAGCAGATGT	AACAGACAGGACACTCGCACT

Chapter 5

PLANT-MEDIATED EFFECTS OF WATER DEFICIT ON THE PERFORMANCE OF *Tetranychus evansi* ON TOMATO DROUGHT-ADAPTED ACCESSIONS

5.1 Introduction

During cultivation, crop plants are exposed to a combination of biotic and abiotic stresses, among which soil water deficiency and arthropod pests are among the most critical. Drought is considered the main environmental factor limiting plant growth and yield worldwide, especially in semiarid areas like the Mediterranean (Chaves et al., 2003). Tomato (*Solanum lycopersicum* L) is a major vegetable crop grown all over the world in outdoor fields and greenhouses, and it is also one of the most water demanding. Thus, it needs to be cultivated under irrigation in most growing areas (Rivelli et al., 2013). Its cultivation is mainly concentrated in semiarid zones, where drought events associated with climate change are expected to be more frequent (Nankishore and Farrell, 2016). Thus, water shortage caused by drought periods can have important consequences for tomato production, as a deficient irrigation might produce yield reduction of up to a 50% in the case of an equivalent reduction in irrigation (Cantore et al., 2016).

The high sensitivity of tomato to water deficit and the need for irrigation in the Mediterranean basin has prompted different approaches for breeding drought-resistant crops, including the search for drought tolerant/adapted varieties (Hu and Xiong, 2014). The term “drought-adapted” refers to higher yield in crop plants under water shortage conditions (Verslues and Juenger, 2011). Remarkably, tomato is one of the crops with the highest diversity of genetic sources, most coming from wild species from arid zones in the Andean region where it came from (Bai and Lindhout, 2007). Furthermore, the landraces formed in secondary centres of diversification like the Mediterranean area can be used as sources of genetic diversity. In this regard, finding tomato drought-adapted varieties might become an important approach to improve water-use efficiency, under future climate change scenarios. Hereof, the ‘Tomàtiga de Ramellet’ tomatoes, which represent a population of landraces from the Balearic Islands (Spain), have been traditionally cultivated outdoors under low water availability conditions during the Mediterranean summer, and thus they represent tomatoes adapted to cultivation under water deficit (Galmes et al. 2011, 2013). Several studies have compared groups of tomato varieties, landraces or wild species observing a differential expression on some drought-associated traits between varieties, but without having a consistent response in the literature. For instance, proline is highly induced by drought in some varieties and poorly in others, whereas leaf water content is generally less reduced on tolerant plants (Sanchez-Rodriguez et al., 2010; Tapia et al., 2016). Other traits like stomatal conductance and the maximum quantum yield of photosystem II photochemistry (Fv/Fm) are the parameters most used as drought

response indicators. In general, a fast decrease in stomatal conductance and a long maintenance of the Fv/Fm value, are indicative of tolerance (Thompson et al., 2007; Mishra et al., 2012; Nankishore and Farrell, 2016; Tapia et al., 2016).

The red tomato spider mite, *Tetranychus evansi* Baker & Pritchard, first recorded in Brazil, has emerged as a serious invasive pest in some areas of Africa and Europe (Navajas et al., 2013). It is described as a species highly tolerant to hot and dry conditions being thus expected to spread northwards across Europe (Meynard et al. 2013). In addition, it has been shown that *T. evansi* is able to suppress tomato plant defenses (Alba et al., 2015), and we have found that both drought and *T. evansi* infestation induced significant changes in the nutritional quality of tomato plants, as more essential amino acids and free sugars are available (Ximénez-Embún et al., 2016). These changes trigger a bottom-up effect on key biological traits of *T. evansi* causing a highly significant increase in leaf damage and mite performance, thus it represents an increasing threat to tomato crop production in the face of climate change.

Nevertheless, to our knowledge, there is little information on how drought-adapted local populations will face the combination of both moderate water deficit, which is the core for the application of deficit irrigation, and the infestation of *T. evansi*. We report here on a holistic approach that considers both drought and spider mites. Accordingly, we investigate the performance of *T. evansi* on drought-adapted (“Tomàtiga de Ramellet” landraces) tomatoes subjected to a moderate water deficit. The mite performance response when feeding on some of the selected landraces was linked to the observed changes in plant nutritional composition and compared to the commercial cultivar, Moneymaker. Furthermore, by imposing these two stresses in combination, the possible directions (synergistic or antagonistic) of the multiple interactions could be examined.

5.2 Materials and methods

5.2.1 Plant material and mite rearing

Four tomato accessions (provided by Dr. Granell, IBMCP, Valencia, Spain) were used in this study: TR58, TR61, TR126 and TR154 (from a population of local landraces ‘Tomàtiga de Ramellet’ (TR) from Mallorca, Balearic Islands, Spain) traditionally grown under non-irrigated conditions during the Mediterranean summer and selected on the basis of their adaptation to water deficit conditions; as well as the commercial cultivar Moneymaker. Tomato plants were grown from seeds in 40 well trays. Plants with 3 expanded leaves were transferred to 2.5 liter pots (diameter: 16 cm, height: 15 cm) (Maceflor ©, Valencia, Spain) filled with 600

g of Universal growing medium 'Compo sana ®' (Compo GmbH, Münster, Germany) and watered to saturation.

A colony of *T. evansi* derived from the Nice strain collected in Beausoleil (South of France) was provided by Dr. Maria Navajas (CBGP, France). Mites were maintained on detached tomato leaves (cv. Moneymaker) placed on ventilated plastic cages (22x30x15 cm) for about 50 generations. The petiole of the leaves was in contact with a thin layer of water in the bottom of the cages to avoid mites escaping and to maintain the leaf turgor. Plants and mite cages were maintained in a climate room at 25±1 °C, 50±5 % relative humidity and a 16 h light/8 h dark photoperiod.

5.2.2 Drought stress regime

Drought stress was attained by water deficiency as described by Ximénez-Embún et al. (2016). In brief, tomato plants were well-watered until they developed four-five expanded leaves, then we imposed two watering treatments, defined as control and moderate drought stress, in order to establish two irrigation regimes. Control plants were watered every two-three days to maintain the soil volumetric water content (Θ) up to 74%. For moderate drought stress, watering was stopped for seven days and thereafter plants were watered to maintain Θ between 21 and 30%. Θ was determined gravimetrically by recording single plant pot weight (balance BSH 6000, PCE Iberica, Tobarra, Spain). Steady stress conditions were reached at about seven to nine days after ceasing irrigation. Water stressed plants from the four accessions were over the wilting point associated with severe drought stress, established at $\Theta \leq 16\%$ for Moneymaker in our experimental conditions (Ximénez-Embún et al., 2016).

The severity of drought stress was assessed by measuring the following parameters on the sub-terminal leaflet of the 4th leaf: a) stomatal conductance (g_s) using a leaf porometer (SC-1 Decagon-T, Pullman, USA); and b) variations in maximum quantum yield of photosystem II photochemistry (F_v/F_m), using a FluorPen FP 100 (PSI, Drasov, Czech Republic). Plant growth was estimated by measuring the stem length (distance between the soil and the terminal bud).

5.2.3 Bioassays

Two different experiments were carried out: the first to measure the effect of moderate drought on mite performance and leaf damage; the second to obtain plant material for chemical and biochemical analyses. Both experiments were carried out in a growth chamber under the same environmental conditions as those used for the mite rearing.

Experiment 1. Tomato plants from each of the four TR accessions were randomly assigned to control or moderate drought treatments. At about seven-nine days, after stopping irrigation, the drought stress conditions had stabilized, corresponding with the phenological stage of six-seven expanded tomato leaves. Then, plants were infested with *T. evansi* females, of unknown age, collected from the laboratory colony by using a vacuum pump D-95 (Dinko S.A., Barcelona, Spain) with a sucking power of 10-50 mmHg connected to a modified Eppendorf. They were placed on the two sub-terminal leaflets of tomato leaves three, four, and five (eight mites per leaflet, 48 mites per plant). All plants (infested and non-infested) were confined with a ventilated metacrylate cylinder fitting the pot diameter and set up in a growth chamber in a complete randomized block design. A total of nine replicates per treatment were performed. Mite performance was assessed at 4 days post infestation (dpi) to avoid overlapping generations, as eggs need at least 5 days to hatch in our experimental conditions. All leaves were detached from the plants, and the number of eggs and mobile mite stages (larvae, nymphs and adults) were counted under a stereomicroscope M125 (Leica Mycosystem, Wetzlar, Germany). The leaf damaged area (mm² of chlorotic lesions) was determined by scanning the damaged leaflets using hp scanjet (HP Scanjet 5590 Digital Flatbed Scanner series, USA) and analyzing the scanned leaflets with the program GIMP 2.8 (www.gimp.org), as described by Ximénez-Embún et al. (2016).

Experiment 2. Plants of two of the selected varieties (TR126 and TR154) and of the cultivar Moneymaker were assigned to four different groups combining two treatments: drought stress (control or moderate drought) and *T. evansi* infestation (infested or non-infested). When drought stress conditions had stabilized, plants were infested as described above. Four days after infestation plant material was collected. The left leaflets from leaves three, four, and five were pooled and grounded in liquid nitrogen to a fine powder and stored for the analysis of total protein, free amino acids and plant defense proteins. The right leaflets from the same leaves were dried together in an oven at 70 °C for 3 days, weighed before and after drying to assess the percentage of water, and grounded using a mortar and pestle to obtain a fine powder and stored for free sugar analysis. Six replicates per treatment were done.

5.2.4 Chemical and biochemical analysis

Chemicals and Equipment. Unless specified otherwise, all chemical compounds were obtained from Sigma-Aldrich (St Luis, USA). Fluorimetric measurements were made using a Varioskan Flash reader (ThermoFisher Scientific, Willmington, USA), and spectrophotometric

measurements with a VERSAmax microplate reader (Molecular Devices Corp., Sunnyvale, USA).

Free sugars. Samples of 40 mg of leaf powder were homogenized in 650 µl of ethanol 95% (v/v), heated at 80 °C for 20 min, centrifuged at 10,000 rpm for 10 min, and the supernatant collected. The process was repeated two more times and the three supernatants were pooled. A volume of 750 µl of the mixture was dried on a SpeedVac Concentrator Savant SVC-100H (ThermoFisher scientific, Willmington, DE, USA) and redissolved in 500 µl of water. Soluble carbohydrate concentration was estimated by the anthrone method (Maness 2010) using glucose as standard. In brief, 1 ml of anthrone reagent (0.2% v/v anthrone on 95% sulfuric acid) was added to the extract, heated at 90 °C for 15 min, and the absorbance measured at 630 nm

Free amino acids. The extraction of the free amino acids was done as described by Hacham et al. (2002). Samples of 50 mg of frozen leaf powder were homogenized with 600 µl of water:chloroform:methanol (3:5:12 v/v/v). After centrifugation at 12,000 rpm for 2 min, the supernatant was collected and the residue was re-extracted with 600 µl of the same mixture, pooling the two supernatants. A mixture of 300 µl of chloroform and 450 µl of water were added to the supernatants, and after centrifugation the upper water:methanol phase was collected and dried in the SpeedVac. The samples were dissolved on 100 µl of sodium citrate loading buffer pH 2.2 (Biochrom, USA) and 10 µl were injected on a Biochrom 30 Amino Acid Analyser (Biochrom, USA) at the Protein Chemistry Service at CIB (CSIC, Madrid, Spain).

Soluble Protein. Samples of 100 mg of leaf frozen powder were homogenized in 500 µl of 0.15M NaCl and ground with fine sand. The homogenate was centrifuged at 12,000 rpm for 5 min at 4 °C, and the soluble protein quantified by absorbance at 280 nm on a Nanodrop 1000 spectrophotometer (ThermoFisher scientific, Willmington, USA).

Plant defense proteins. Samples of 100 mg of leaf frozen powder were homogenized with 500 µl of extraction buffer (0.15 M NaCl for protease inhibitors, and 0.1 M phosphate buffer, pH 7.0; 5% w:v polyvinylpolypyrrolidone for oxidative enzymes) and soluble protein quantified as explained above.

Protease inhibitors. The inhibitory activity of plant protein extracts was tested against commercial enzymes: papain (EC 3.4.22.2), cathepsin B from bovine spleen (EC 3.4.22.1), trypsin from bovine pancreas (EC 3.4.21.4), α -chymotrypsin from bovine pancreas (EC 3.4.21.1), cathepsin D from bovine spleen (EC 3.4.23.5) and leucine aminopeptidase from porcine pancreas (EC 3.4.11.1); as described by Ximénez-Embún *et al.* (2016). Samples of 20 µg of plant protein extracts (40 µg in case of leucine aminopeptidase inhibition assay) were

preincubated for 10 min with 100 ng of the commercial enzyme. Reaction conditions are summarized in Table S5.1. Results were expressed as a percentage of protease activity inhibited.

Oxidative enzymes. Polyphenol oxidase (PPO) activity was analyzed by incubating 20 μ l of enzyme extract with catechol (40 mM final concentration) in 160 μ l of Tris-HCl pH 8.5 buffer at 30 °C for 1 h. Absorbance was read at 420 nm. Peroxidase (POD) activity was determined incubating 20 μ l of a 1:10 dilution of the enzyme extract with guaiacol (5 mM final concentration) and H₂O₂ (2.5 mM final concentration) in 150 μ l of potassium phosphate pH 6 buffer at 30 °C for 10 min. Absorbance was read at 470 nm. PPO and POD activities were expressed as nmol substrate metabolized relative to time and total protein content.

5.2.5 Statistical analysis

All plant and mite data were checked for the assumptions of normality and heteroscedasticity, and transformed if necessary. Stem length and stomatal conductance were $\log_{10}(x)$ transformed while Fv/Fm was $\log_{10}(x+1)$ transformed, and the three parameters were analyzed at mite infestation and at 4 days post infestation using a two-way ANOVA (using as fixed factors drought-stress treatment and tomato variety) and Bonferroni *post hoc* tests to compare all combinations of treatments and varieties for each time. The number of *T. evansi* eggs, mobile forms and leaf damaged area were $\log_{10}(x)$ transformed and analyzed by a two-way ANOVA (drought stress treatment and tomato variety as fixed factors) and Bonferroni *post hoc* tests to compare drought-stress treatments within each variety. The percentage of water, free sugars, protein, free amino acids, and protease inhibition were arcsine square root transformed. These data and the oxidative enzyme activities were analyzed for each tomato variety separately using a two-way ANOVA (using as fixed factors drought stress and mite infestation treatments) and Newman-Keuls *post hoc* tests to make comparisons among treatments.

5.3 Results

5.3.1 Effect of drought on stomatal conductance, photosynthetic efficiency and tomato plant growth.

The impact of moderate drought stress on stomatal conductance, photosynthetic efficiency, and stem length was similar in Experiment 1 (Figure S5.1) and 2 (Figure S5.2). The infestation by *T. evansi* did not affect any of the physiological parameters analyzed, and thus data from infested and non-infested plants were pooled in Experiment 2. The Fv/Fm, values in

turn, corroborated that severe drought stress conditions were never reached, since values were never below 0.7 (Richie, 2006; Mishra et al., 2012).

In all plant varieties, drought induced a significant reduction of stomatal conductance that was about 3.5 and 4.5 times smaller at mite infestation and 4 dpi, respectively (Figures S5.1A and S5.2A). No significant differences in stomatal conductance were found among the four accessions in control plants. However, the following differences in stomatal conductance were observed when plants were exposed to drought stress: it was significantly higher in TR126 than in TR154 at mite infestation, and in TR58 and TR126 than in TR61 at 4 dpi in Experiment 1 (Figure S5.1A); and higher in Moneymaker than in TR154 and both higher than in TR126 at the time of mite infestation in Experiment 2 (Figure S5.2). The photosynthetic efficiency (Fv/Fm) was not affected by drought on any variety (Figures S5.1B and S5.2B), with the exception of TR126 that has lower Fv/Fm in control than in drought-stressed plants at mite infestation in Experiment 2. However, some differences were observed among varieties, being Fv/Fm lower for TR58 at mite infestation in Experiment 1 (Figure S5.1B) and for TR126 at mite infestation in Experiment 2, (Figure S5.2B), though they were not significant differences at 4 dpi in both cases. Moderate drought stress reduced the plant growth, as stem length was smaller on drought stressed plants in all cases (Figures S5.1C and S5.2C). There were also differences in growing rate between varieties, being TR58 the fastest one with the lowest reduction in growth due to drought (Figure S5.1C). The stem length of Moneymaker was smaller than in TR126 and TR154 (Figure S5.2C).

5.3.2 Effect of drought on *T. evansi* performance.

A differential effect of moderate drought was observed on *T. evansi* performance depending on the tomato TR accessions they were feeding on (Figure 5.1). Females laid 1.3 and 1.4 times more eggs and produced 1.5 and 1.3 more damaged area on drought stressed than on control plants when feed on varieties TR61 and TR154, respectively (Figure 5.1A&B). In contrast, similar numbers of eggs and damaged area were obtained on control and drought stressed plants when fed on varieties TR58 and TR126. The number of mobile forms recovered at 4 dpi (surviving females, since eggs need at least 5 days to hatch in our experimental conditions) was not significantly different between drought stressed and control plants for all TR accessions.

The accessions TR154 and TR126 were selected for further chemical and biochemical analyses. The reasoning for this selection is that they represent the two types of plant-mediated effects of water deficit on mite performance: enhanced (TR154) and no effect (TR126); and that in both cases the biological parameters obtained on control plants are closer to the values obtained on tomato MoneyMaker under identical experimental conditions (Ximénez-Embún et al., 2016) (Figure 5.1).

Figure 5.1 Performance of *T. evansi* in the tomato accessions TR58, TR61, TR126 and TR154. The number of eggs (a) and leaf damaged area (b) on control and moderate drought stressed tomato plants at 4 days post infestation were measured. Data are mean $\pm$ SE of 9 replicates/treatment. * indicates a statistically significant difference between treatments within each variety; n.s. indicates no significant difference (Two-way ANOVA, Bonferroni *post hoc* test, $P < 0.05$). The values obtained by Ximénez-Embún et al. (2016) on tomato MoneyMaker under identical experimental conditions: control (cMM) and moderate drought stress (dMM), have been included for comparison.

5.3.3 Changes in plant nutritional composition induced by drought and *T. evansi*.

The nutritional composition of tomato leaves was determined by analyzing water, free sugars, protein and total free amino acid content (Figure 5.2). The combination of drought stress and *T. evansi* infestation was the most significant factor for Moneymaker, inducing the amount of total free sugars and amino acids, whereas drought stress alone induced an increase on total free sugars and *T. evansi* infestation alone had not significant effects. TR154 was affected by *T. evansi* infestation that induced an increase on free sugars, and all treatments had lower levels of protein than the control. In contrast, drought and/or *T. evansi* infestation didn't caused any significant change on the nutritional composition in TR126. When the three were compared under control conditions, Moneymaker had significantly lower amounts of water (One-way ANOVA, and Newman-Keuls *post hoc* test, $F = 4.737$, $P = 0.0254$) and free sugars ($F = 23.11$,

Figure 5.2 Effect of moderate drought, *T. evansi* infestation and the combination of both on nutritional composition: A) water, B) total free sugars, C) protein and D) total free amino acids of tomato accessions TR126, TR154 and Moneymaker (MM) at 4 days post infestation. Data are mean $\pm$ SE of 6 replicates/treatment. Different lower case letters indicate significant differences among treatments for a given variety (Two-way ANOVA, Newman-Keuls *post hoc* test, $p < 0.05$).

$P < 0.0001$) than TR154 and TR126, but similar amounts of protein ($F = 1.151$, $P = 0.3427$) and free amino acids ($F = 1.391$, $P = 0.2791$).

The levels of specific amino acids, classified as essential or non-essential for *T. urticae* according to Rodriguez and Hampton (1966), were analyzed (Table 5.1). Proline, an indicator of drought stress, was induced in the three varieties by drought alone or in combination with mite infestation. With respect to the rest of amino acids, similar results were obtained with the varieties TR154 and Moneymaker, which responded to the combination of both stresses with an increase of non-essential (serine) and essential amino acids (isoleucine, leucine, tyrosine, histidine, lysine and arginine). The only differences being an increase of glutamine in Moneymaker and of valine in TR154. In contrast, only one essential amino acid (isoleucine) was induced in the variety TR126 by the combination of drought and mite infestation, and the non-essential glutamine was induced by drought.

5.3.4 Effect of drought and *T. evansi* on tomato defense proteins.

Tomato plant defense proteins were affected by drought stress and mite infestation, but different responses were obtained depending on the variety (Table 5.2). Moneymaker was the variety with less defense proteins altered: drought stress induced an increase on peroxidase activity, and *T. evansi* infestation induced the inhibition activity against cathepsin D. The variety TR126 showed the highest changes in the levels of defense proteins as response to stress: drought induced the inhibitory activity against trypsin and aminopeptidase, *T. evansi* increased the inhibition against cathepsin B and chymotrypsin, and the combination of both stresses the inhibitory activity against cathepsins B and D, trypsin and chymotrypsin. In turn, the variety TR154 showed an intermediate response: drought induced a decrease on the cathepsin B inhibitory activity and an increase on peroxidase activity, *T. evansi* induced the inhibition of cathepsins B and D and papain, and the combination of both stresses induced an increase of peroxidase activity.

Table 5.1 Effect of moderate drought stress, *T. evansi* infestation and the combination of both (Dr+Te) on the amino acid composition of the tomato accessions TR126, TR154 and Moneymaker at 4 days post infestation.

	TR126				TR154				Moneymaker			
	Control	Drought	<i>T. evansi</i>	Dr+Te	Control	Drought	<i>T. evansi</i>	Dr+Te	Control	Drought	<i>T. evansi</i>	Dr+Te
Non-essential amino acids												
Asp	2.10±0.50a	2.79±0.91a	2.97±0.52a	1.91±0.18a	4.00±0.89a	2.02±0.17a	2.84±1.12a	3.80±0.83a	1.47±0.11a	2.57±1.28a	1.32±0.08a	3.04±0.80a
Thr	0.40±0.10a	0.98±0.36a	0.58±0.09a	0.71±0.08a	1.10±0.28a	0.89±0.11a	0.78±0.27a	1.48±0.27a	0.54±0.07a	1.70±0.94a	0.57±0.03a	1.83±0.37a
Ser	2.00±0.70a	2.68±1.02a	2.77±0.92a	2.72±0.51a	2.49±0.45a	4.42±0.97ab	2.44±0.62a	7.36±1.53b	1.87±0.43a	8.42±5.52ab	1.80±0.26a	11.8±3.41b
Glu	3.20±0.40a	10.00±3.60b	4.79±0.25ab	5.13±0.90ab	12.00±2.60a	7.33±0.52a	8.82±2.63a	14.09±3.00a	4.21±0.50a	4.34±1.00a	4.34±0.30a	8.38±1.15b
Gly	0.03±0.01a	0.05±0.02a	0.04±0.01a	0.07±0.03a	0.04±0.01a	0.03±0.01a	0.03±0.01a	0.04±0.01a	0.02±0.01a	0.13±0.11a	0.02±0.01a	0.05±0.01a
Ala	0.35±0.09a	0.43±0.16a	0.43±0.08a	0.49±0.21a	0.41±0.12a	0.26±0.04a	0.28±0.09a	0.47±0.10a	0.22±0.06a	0.92±0.73a	0.25±0.06a	0.56±0.13a
Cys	0.03±0.01a	0.05±0.02a	0.04±0.01a	0.05±0.01a	0.06±0.01a	0.05±0.01a	0.05±0.02a	0.10±0.02a	0.02±0.01a	0.12±0.1a	0.02±0.01a	0.09±0.02a
Pro	0.08±0.02a	0.79±0.27b	0.13±0.02a	0.84±0.32b	0.28±0.06a	0.75±0.09b	0.2±0.05 a	1.39±0.44b	0.11±0.01a	1.20±0.79 b	0.11±0.01a	2.42±0.75b
Essential amino acids												
Val	0.16±0.04a	0.36±0.13a	0.22±0.04a	0.51±0.16a	0.36±0.07a	0.65±0.17ab	0.41±0.1 a	1.10±0.22b	0.26±0.08a	1.55±1.16a	0.29±0.04a	1.55±0.41a
Met	0.02±0.01a	0.03±0.02a	0.03±0.01a	0.08±0.05a	0.04±0.01a	0.03±0.01a	0.03±0.01a	0.04±0.01a	0.02±0.01a	0.21±0.18a	0.02±0.01a	0.07±0.02a
Ile	0.10±0.02a	0.20±0.07ab	0.15±0.04ab	0.33±0.10b	0.21±0.04a	0.47±0.14ab	0.28±0.08a	0.80±0.18b	0.18±0.05a	1.01±0.75ab	0.21±0.04a	1.38±0.24b
Leu	0.08±0.02a	0.14±0.05a	0.11±0.03a	0.46±0.25a	0.20±0.04a	0.38±0.13ab	0.25±0.06a	0.65±0.14b	0.15±0.05a	1.45±1.23ab	0.18±0.03a	1.32±0.24b
Tyr	0.07±0.02a	0.1±0.03a	0.11±0.03a	0.27±0.12a	0.13±0.02a	0.36±0.13ab	0.22±0.07ab	0.55±0.12b	0.13±0.05a	0.86±0.66a	0.16±0.03a	1.46±0.22b
Phe	0.13±0.02a	0.26±0.09a	0.19±0.02a	0.42±0.17a	0.34±0.07a	0.35±0.09a	0.34±0.10a	0.62±0.12a	0.19±0.03a	1.04±0.82a	0.2±0.02a	0.84±0.20a
His	0.06±0.02a	0.09±0.03a	0.10±0.04a	0.15±0.04a	0.11±0.02a	0.33±0.08a	0.18±0.04a	0.68±0.19b	0.12±0.04a	0.45±0.28a	0.13±0.03a	0.88±0.16b
Lys	0.09±0.03a	0.12±0.05a	0.15±0.06a	0.37±0.20a	0.17±0.03a	0.36±0.09a	0.19±0.05a	0.63±0.12b	0.11±0.03a	1.17±0.97a	0.13±0.02a	1.09±0.24b
Arg	0.30±0.13a	0.07±0.04a	0.73±0.55a	0.24±0.12a	0.16±0.03a	0.40±0.15ab	0.27±0.10ab	0.63±0.15b	0.15±0.05a	2.10±1.64a	0.18±0.04a	3.93±1.33b

Data are the mean amount of amino acid (mg) per gram of dry weight ± SE. Different lower case letters indicates significant differences among treatments for a given variety (Two-way ANOVA, Newman-Keuls *post hoc* test, $p < 0.05$). The division between essential and no-essential amino acids is based on a study with the closely related species *Tetranychus urticae* (Rodriguez and Hampton, 1966). In bold the amino acids that showed significant difference between treatments for a given accession.

Table 5.2 Effect of moderate drought, *T. evansi* infestation and the combination of both (Dr+Te) on plant defense proteins of the tomato accessions TR126, TR154 and Moneymaker at 4 days post infestation.

	TR126				TR154				Moneymaker			
	Control	Drought	<i>T. evansi</i>	Dr+Te	Control	Drought	<i>T. evansi</i>	Dr+Te	Control	Drought	<i>T. evansi</i>	Dr+Te
Protease inhibitors (% inhibition)												
Cathepsin B	47±3ab	41±2a	55±3b	53±3b	46±4a	36±2b	58±2c	53±3ac	42±4a	38±5a	54±4a	45±5a
Papain	66±5a	46±7a	71±5a	61±11a	49±5ab	40±6a	70±7b	61±11ab	50±8a	45±9a	72±7a	64±5a
Cathepsin D	36±3a	55±4ab	47±3ab	59±5b	47±3a	47±5a	65±4b	59±5ab	45±4a	55±5ab	67±6b	61±5ab
Trypsin	24±1a	32±2b	27±1ab	33±2b	36±5a	31±1a	36±2a	33±2a	32±5a	29±3a	36±4a	29±4a
Chymotrypsin	33±5a	35±3a	52±5b	49±5b	44±3a	38±3a	51±5a	49±5a	40±2ab	29±5a	49±5b	43±3ab
Aminopeptidase	36±2a	42±1b	34±2a	36±1a	31±2a	39±6a	41±1a	36±1a	37±3a	36±3a	43±5a	41±3a
Oxidative enzymes (specific activity)												
Polyphenol oxidases¹	5.8±0.7a	7.4±1.0a	5±1a	5.1±0.7a	4.8±0.6a	5.5±0.7a	6.2±1a	5.1±0.6a	4.8±0.5a	6.2±0.4a	5.4±0.7a	5.0±0.5a
Peroxidases²	3.5±0.4a	3.2±0.5a	3±0a	3.6±0.5a	2.9±0.3a	3.8±0.3b	4±0.6b	4±0.2b	2.4±0.3a	3.6±0.4b	2.9±0.3ab	2.8±0.3ab

Data are mean ± SE. Different lower case letters indicates significant differences among treatments for a given variety (Two-way ANOVA, Newman-Keuls *post hoc* test, $p < 0.05$). In bold the defense proteins that showed significant difference between treatments for a given accession.

(1) PPO: nmol Cathecol metabolized/ mg Protein*min

(2) POD: nmol Guaiacol metabolized/ mg Protein*min.

5.4 Discussion

Our data reveal that drought stress, *T. evansi* infestation and their combined effect have a differential effect on the performance of *T. evansi* on drought-adapted 'Tomàtiga de Ramellet' tomatoes. Thus, the mite performance on two of the TR accessions (TR61 and TR154) had a similar pattern that the one found on Moneymaker, though the actual values of the biological traits (egg laid and leaf damage) were considerably smaller. In contrast, no increase in mite performance was observed on TR58 and TR126, as similar amount of eggs laid and leaf damage were recorded on drought-stressed and not stressed tomato plants. This is an important issue, as it has been reported that the changes induced by drought stress on the tomato cultivar Moneymaker triggered a bottom up effect on *T. evansi*, thus increasing mite performance and the damage to the plant (Ximénez-Embún et al., 2016). These results cannot be explained by morphological adaptation of the TR accession to water deficit conditions, as all TR accessions had a significant reduction of stem length but didn't show a consistent differential response. Other parameters like the stomatal conductance and Fv/Fm did not differ consistently between the accessions. Previous studies of 'Tomàtiga de Ramellet' tomatoes showed that differences in Fv/Fm between varieties were not maintained over time in most accessions (Galmes et al., 2011).

A clear link, in contrast, could be observed between some of the plant nutritional components and *T. evansi* performance on the two accessions analyzed. We have considered those nutrients that could be relevant for mite performance. Among them, the increased levels of relative water content of TR accessions compared with Moneymaker seems to be clearly related to their adaptation to water deficit conditions, but not to the differential performance of *T. evansi*. The accumulation of free sugars has been reported to increase tomato tolerance to drought (Khodakovskaya et al., 2010; Tapia et al., 2016), in line with the high basal levels found in both TR accessions. However, a significant induction of free sugars, as a response to drought stress was only observed in the non drought-adapted Moneymaker variety, which is the one with the best *T. evansi* performance (Ximénez-Embún et al., 2016). Tomato plants have been reported to metabolize protein into amino acids in response to drought stress (Bauer et al., 1997; Ximénez-Embún et al., 2016). We have observed the amino acid composition changed between accessions as well as the induction patterns by drought stress and *T. evansi*. We recorded a similar pattern of increase of essential amino acids for Moneymaker and TR154, particularly by the combined effect of drought stress and mite infestation, which can explain the better performance of *T. evansi* on these plants. In contrast, the performance was not affected in

TR126. Interestingly, in this later case only one essential amino acid was increased significantly under the same stress conditions. The positive effect of amino acid concentration on mite performance has been previously observed in *T. evansi* (Ximénez-Embún et al., 2016) and *T. urticae*, a related species (Tulisalo, 1971; Dabrowski and Bielak, 1978; Ximénez-Embún et al., 2017). A special mention should be made of proline, an osmolite traditionally used as a good indicator of plant response to drought (Claussen, 2005). However, drought-adapted tomato cultivars have been reported to have a smaller increase in proline than sensitive ones (Sanchez-Rodriguez et al., 2010; Tapia et al., 2016). Thus, the levels found in TR126 and TR154 are in agreement with this, and there are much lower than the ones found in Moneymaker, where proline was reported to act as a feeding stimulant of *T. evansi* above certain thresholds (Ximénez-Embún et al., 2016).

Another important aspect that determines plant palatability and thus, mite performance, is the production of plant defense proteins. Interestingly, differences were observed between accessions, with TR126 showing the highest induction of defense proteins in response to drought and/or mite infestation, and Moneymaker the lowest levels of induction. The main differences were observed on protease inhibitors (PIs), which are recognized as key components of the defensive response of tomato to mite infestation (Kant et al., 2004; Santamaria et al., 2012, 2015b; Alba et al., 2015). Serine (trypsin and chymotrypsin) and aminopeptidase PIs were induced in TR126 by drought stress, *T. evansi* or both stresses combined, but not in Moneymaker or TR154 by any treatment. Differences were also observed for cysteine (cathepsin B) PIs that were induced by mite infestation, though reduced by drought stress, in TR154, but were not altered in TR126 or Moneymaker. The only type of PI analyzed that were induced in all plants by mite infestation alone or when combined with drought stress are those targeting aspartyl (cathepsin D) proteases. It has been reported that *T. evansi* suppresses plant defenses in tomato (Kant et al., 2015), including the down-regulation of PIs in two different tomato varieties (Santa Clara and Castlemart) (Sarmiento et al., 2011; Alba et al., 2015). Our results suggest that this suppression ability is maintained on Moneymaker, but lost on TR126 for serine PIs and on TR154 for cysteine PIs. An induction of cysteine (papain and cathepsin B) PIs by drought, *T. evansi* infestation and the combination of both was reported on Moneymaker (Ximénez-Embún et al., 2016), but this induction was only observed at a later response (10 dpi) and with mite infestation levels about 8 times bigger than in the present study. The differential induction of cysteine and serine PIs in the TR accessions, when compared to Moneymaker, might be a factor contributing to the lower performance of the mites on these

plants. *T. evansi* relies mostly on cysteine (cathepsin B, and L- and legumain-like) and aspartyl (cathepsin D-like) proteases and aminopeptidases for proteolytic digestion (Ximénez-Embún et al., 2016). Thus, the ingestion of PIs targeting some of these proteases may be harmful, as already demonstrated for *T. urticae* (Carrillo et al., 2011; Santamaria et al., 2012). Serine proteases do not appear to be directly involved in the hydrolysis of dietary proteins in this species, but tomato serine PIs may target other physiological processes, as has been indicated for *T. urticae* which has a similar digestive proteolytic profile (Santamaria et al., 2012, 2015b).

With regard to oxidative enzymes, peroxidase and polyphenol oxidase activities have been reported to be induced in tomato by drought stress (English-Loeb et al., 1997; Inbar et al., 2001) and mite infestation (Kant et al., 2004; Alba et al., 2015; Ximénez-Embún et al., 2016). We have found different results for peroxidase activity depending on the tomato plants, since it was induced by drought, mite and the combination of both on TR154, only by drought in Moneymaker, and it was not affected in TR126. The production of peroxidases is an adaptive mechanism for the scavenging and detoxification of reactive oxygen species in drought-stressed plants (Rai et al., 2013), but its induction might vary depending on the tomato plant genotype (Sanchez-Rodriguez et al., 2010). This might explain the absence of induction in the case of TR126. We did not observe significant differences in polyphenol oxidase activity, whose expression pattern has been shown to vary depending on both the duration of plant exposure to drought stress and the infestation by different mite species (Ximénez-Embún et al., 2016, 2017).

5.5 Conclusions

All together, our data revealed differential plant-mediated effects of water deficit on the performance of *T. evansi* on drought-adapted 'Tomàtiga de Ramellet' tomatoes. Those accessions where free amino acids were induced in response to drought stress resulted more suitable for the mites, as it has already been reported for the commercial cultivar Moneymaker (Ximénez-Embún et al., 2016). On the contrary, the accessions where defense proteins were induced by drought stress and/or mite infestation contributed to the lower performance of *T. evansi*. These findings have important implications for decision making in the selection of the tomato seed to be planted in a forthcoming climate change scenario, as the nutritional changes observed might speed up or slow down the expansion of this important invasive species in area-wide tomato production.

5.6 Supporting information

Figure S5.1 Effect of moderate drought on A) stomatal conductance (gs), B) maximum quantum yield of PSII photochemistry (Fv/Fm) and C) stem length of the tomato accessions TR58, TR61, TR126 and TR154 at mite infestation and at 4 days post infestation. Data shown are mean $\pm$ SE of 9 replicates/treatment from Experiment 1. Different lower case letters indicates significant differences among all combinations of treatments and varieties for each time (Two-way ANOVA, Bonferroni *post hoc* test, $p < 0.05$).

Figure S5.2 Effect of moderate drought on A) stomatal conductance (gs), B) maximum quantum yield of PSII photochemistry (Fv/Fm) and C) stem length of the tomato accessions Moneymaker (MM), TR126 and TR154 at mite infestation and at 4 days post infestation. Data (mean $\pm$ SE) shown are from infested (6 replicas) plus non-infested (6 replicas) plants on Experiment 2. Different lower case letters indicate significant differences among all combinations of treatments and varieties from each time (Two-way ANOVA, Bonferroni *post hoc* test, $p < 0.05$).

Table S5.1. Summary of analytical methods to assess the inhibitory activity of plant protein extracts ¹

Commercial enzyme ²	Substrate ³	Buffer ⁴	Incubation	Measurement ⁵
Cathepsin B from bovine spleen (EC 3.4.22.1)	Z-RR-AMC	100 mM Na phosphate, pH 6.0 (10 mM L-cysteine, 10 mM EDTA, 0.01% (v/v) Brij 35)	30 min at 28 °C	excitation filter 350 nm emission filter 465 nm
Papain (EC 3.4.22.2)	Z-FR-AMC	100 mM Na phosphate, pH 6.0 (10 mM L-cysteine, 10 mM EDTA, 0.01% (v/v) Brij 35)	30 min at 28 °C	excitation filter 350 nm emission filter 465 nm
Cathepsin D from bovine spleen (EC 3.4.23.5)	MocAc-GKPILFFRLK (Dnp)-D-R- NH ₂	100 mM sodium citrate, pH 3.5 (0.15M NaCl, 5 mM MgCl ₂)	20 min at 30 °C	excitation filter 328 nm emission filter 393 nm
Trypsin from bovine pancreas (EC 3.4.21.4),	Z-LA-AMC	100 mM Tris-HCl, pH 7.5 (0.15M NaCl, 5 mM MgCl ₂)	1 h at 35 °C	excitation filter 350 nm emission filter 465 nm
α -Chymotrypsin from bovine pancreas (EC 3.4.21.1),	SucAAPF-AMC	100 mM Tris-HCl, pH 7.5 (0.15M NaCl, 5 mM MgCl ₂)	30 min at 35 °C	excitation filter 350 nm emission filter 465 nm
Leucine aminopeptidase from porcine pancreas (EC 3.4.11.1).	LpNa	100 mM Tris-HCl, pH 8 (0.15M NaCl, 5 mM MgCl ₂)	30 min at 30 °C	absorbance at 410 nm

¹ Procedures adapted from Ximénez-Embún et al. (2016). Samples of 20 µg of plant protein extracts (40 µg in case of leucine aminopeptidase) were preincubated for 10 min with 100 ng of the commercial enzyme.

² All purchased from Sigma-Aldrich (St Luis, USA).

³ The substrates were added at a final concentration of 20 µM. Z-RR-AMC (N-carbobenzoxycyloxy-Arg-Arg-7-amido-4-methylcoumarin) for cathepsin B, Z-FR-AMC (N-carbobenzoxycyloxy-Phe-Arg-7-amido-4-methylcoumarin) for papain, Z-LA-AMC (Z-L-Arg-7-amido-4-methylcoumarin) for trypsin, SucAAPF-AMC (Suc-Ala-Ala-Pro-Phe-7-amido-4-methylcoumarin) for chymotrypsin, all purchased from Calbiochem (MerkMilipore, Billerica, USA), MocAc-GKPILFFRLK(Dnp)-D-R- NH₂ from Peptanova (Germany) for cathepsin D, and LpNa (L-leucine p-nitroanilide) from Sigma-Aldrich (St Luis, USA) for leucine aminopeptidase.

⁴ Concentrations are expressed at molarity in the reaction mixture.

⁵ AMC (7-amino-4-methylcoumarin) (Bachem, Switzerland) as standard for all fluorescent substrates, except MCA (MoCAC-Pro-Leu-Gly) (Peptanova GmbH, Germany) for cathepsin D. Double blanks were used to account for spontaneous breakdown of substrates and the plant protease activity, and all assays were done in duplicate.

Chapter 6

GENERAL DISCUSSION

The effect of biotic (arthropod pests) and abiotic (drought) stresses on plants has been usually studied separately. However, in nature both stresses often co-occur, interacting with each other modifying the plant response output. Moreover, biotic and abiotic stress interactions will increase under conditions of climate change. Thus, it is becoming increasingly important to understand the ecological consequences of arthropod pests and water deficit interaction, particularly in summer irrigated crops like tomato in Mediterranean areas, where deficit irrigation schedules are predicted. In this thesis, we have investigated the tomato plant response to mites and drought stress and how they interact in order to provide insights on the influence of drought stress on mite's behavior, development and physiology. We have demonstrated that the performance of two different spider mites (*Tetranychus urticae* and *T. evansi*) and one eriophyid mite (*Aculops lycopersici*) was enhanced when feeding on moderate drought-stressed tomato plants (cv. Moneymaker), increasing thus outbreak and yield loss risks in area-wide tomato production. However, when four accessions of the drought-adapted tomatoes, 'Tomàtiga de Ramellet' (TR) were analyzed, it was found an enhancement on mite performance by drought in two accessions (TR61 and TR154), but not in the other two (TR58 and TR126), which could help to support the choice of cultivars by farmers.

In all experiments the moderate drought conditions caused a reduction on tomato plant growth, stomatal conductance and Fv/Fm, as an adaptation to water deficit conditions and in agreement with previous reports (Thompson et al., 2007; Mishra et al., 2012; Nankishore and Farrell, 2016; Tapia et al., 2016). The accumulation of proline, considered as a drought response indicator (Claussen, 2004; Sanchez-Rodriguez et al., 2010), was also observed in Moneymaker and in the four TR accessions tested. However, an increase on free amino acids considered essential for mites (Rodriguez and Hampton, 1966) was observed on Moneymaker and on TR154 but not on TR126. Accumulation of free sugars was also observed in all experiments with Moneymaker. The effect of drought on plant defences was minor and no consistent, especially for the protease inhibitors (PIs) as observed by English-Loeb et al. (1997). All together, our data revealed a positive link between the induction of soluble carbohydrates and amino acids used by the plant for osmotic adjustment and the mite performance.

When feeding on tomato plants, *T. evansi*, *A. lycopersici* and some strains of *T. urticae* have been reported to be able to suppress the induction of plant defences (Glas et al., 2014; Alba et al., 2015; Kant et al., 2015; Wybouw et al., 2015). However, we have showed that while *T. evansi* and *A. lycopersici* suppress mainly serine PIs and induce cysteine PIs and oxidative

enzyme activities, *T. urticae* had the opposite effect suppressing cysteine PI and oxidative enzyme and inducing serine PI, being the induction higher in response to a non-adapted strain than an adapted one. The mechanism of defense suppression, which can explain the differences between mite species, is still unknown (Kant et al., 2015), though mite salivary secretions contain a wide amount of proteins that can potentially interfere with plant defences (Jonckheere et al., 2016; Villarroel et al., 2016). This suppression ability is lost on TR126 for serine PIs and on TR154 for cysteine PIs and oxidative enzymes, which might be a factor contributing to the lower performance of the mites on these plants when compared to Moneymaker. In regard of nutrients, we observed that *T. urticae* and *T. evansi* infestation induced the accumulation of free sugars while *A. lycopersici* induces the increase of essential amino acids. This can explain the facilitation of *T. urticae* development on tomato leaf previously infested with *A. lycopersici* (Glas et al., 2014).

The plant response to the combination of both stresses (drought and mite infestation) determines the performance of mite on drought stressed plants. We observed an additive effect between both stresses on most of the physiological parameters analyzed. Thus, the levels of free sugars and free amino acids in most of the experiments were higher when both stresses were combined than when the stresses were presented separately. In the case of defence compounds, the combination of stresses also induced higher levels of PIs, whereas oxidative enzyme activities were not altered. In any case, tomato plant defences didn't showed a clear effects on *T. evansi* and *T. urticae* digestive and detoxification enzymes, which in the later case depended mostly on their adaptation to tomato. Our results are thus, in agreement with the "plant stress hypothesis" (White, 1984, 2009), as mites benefit from the increases on free sugars and amino acids and are able to overcome the increase defense proteins. The link between plant nutritional value and mite performance was confirmed by the use of tomato drought-adapted TR accessions that differ on its metabolic response to the biotic and abiotic stresses. This represent a new approach on studying the plant mediated herbivore-drought interaction, as most of the previous works focus mainly on how drought induced changes on tomato plants affects differently insects with different feeding strategies (English-Loeb et al., 1997; Inbar et al., 2001).

All together, these data revealed that drought stimulates mite performance and increases plant damage on different mite species. Furthermore, it provides an experimental framework for screening for drought-resistant tomato accessions that will be at the same time resistant to herbivore mites, since we have found that the levels of nutrients will determine that a plant that

is resistant to drought might be also resistant to mites or more susceptible, as it was observed on the TR126 accession. This finding is especially important in the task of adapting area-wide tomato production to mitigate the effects of climate change, and for the management and prediction of herbivore mite proliferation.

Chapter 7

CONCLUSIONS

7. Conclusions

1. The performance of three key mite pests (*Tetranychus urticae*, *T. evansi* and *Aculops lycopersici*) was remarkably improved when feeding on moderate drought-stressed tomato plants (cv. Moneymaker) increasing plant damage and the potential yield losses.
2. Both drought and mite infestation induced changes in the nutritional quality of tomato plants, as more essential amino acids and/or free sugars were available, contributing to improve mite performance of all mite species fed on Moneymaker.
3. An additive effect between both stresses was recorded on most of the physiological parameters analyzed. In general, the levels of free sugars and free essential amino acids were higher when both stresses were combined.
4. It was demonstrated that L-proline, the amino acid highest induced by drought, had a phagostimulant effect on *T. evansi*, increasing both egg laid and leaf damage area in tomato.
5. The link between plant nutritional value and mite performance was confirmed by the use of drought-adapted ‘Tomàtiga de Ramellet’ tomato accessions, since *T. evansi* performance was enhanced only on those accessions in which drought induced the accumulation of essential free amino acids.
6. The three mite species attenuate some of the defense responses of Moneymaker, but while *T. evansi* and *A. lycopersici* suppress mainly serine PIs and induce cysteine PIs and oxidative enzymes, *T. urticae* had the opposite effect suppressing cysteine PIs and oxidative enzymes and inducing serine PIs.
7. The plant defenses suppression ability was lost in *T. evansi* when feeding on some of the ‘Tomàtiga de Ramellet’ accessions, which might be a factor contributing to its lesser performance on these plants as compared to Moneymaker.
8. The digestive and detoxification physiological responses of the mites indicate that *T. evansi* and tomato-adapted *T. urticae* are able to circumvent the potential adverse effects of tomato plant defenses.

9. Our findings support the “Plant stress hypothesis” and suggest that drought-stressed tomato plants, even at a mild level, may be more prone to mite outbreaks in a climate change scenario, which might strongly affect tomato production on area-wide scales.
10. Overall, our data have important implications for decision-making in the choice of cultivars by farmers and provides an experimental framework for screening for drought-adapted tomato accessions, and for the management and prediction of herbivore mite proliferation.

Chapter 8

REFERENCES

- Achuo, E.A., Prinsen, E., Hofte M. 2006. Influence of drought, salt stress and abscisic acid on the resistance of tomato to *Botrytis cinerea* and *Oidium neolycopersici*. *Plant Pathology* 55, 178-186.
- Agrawal, A.A., Vala, F., Sabelis, M.W. 2002. Induction of preference and performance after acclimation to novel hosts in a phytophagous spider mite: adaptive plasticity? *The American Naturalist* 159, 553–565.
- Ahn, S.J., Dermauw, W., Wybouw, N., Heckel, D.G., Van Leeuwen, T. 2014 Bacterial origin of a diverse family of UDP-glycosyltransferase genes in the *Tetranychus urticae* genome. *Insect Biochemistry and Molecular Biology* 50, 43–57.
- Akimov, I.A., Barabanova, V.V. 1997 Morphological and functional characteristics of the digestive system in tetranychid mites (Trombidiformes, Tetranychoidae). *Entomologicheskoe Obozrenie* 56, 912-922.
- Alba, J.M., Schimmel, B.C.J., Glas, J.J., Ataide, L.M., Pappas, M.L., Villarroel, C.A., Schuurink, R.C., Sabelis, M.W., Kant, M.R. 2015 Spider mites suppress tomato defenses downstream of jasmonate and salicylate independently of hormonal crosstalk. *New Phytologist* 205, 828–840.
- Asbahi, A.A.A., Maqtari, M.A.A., Naji, K.M. 2012 ABA biosynthesis defective mutants reduce some free amino acids accumulation under drought stress in tomato leaves in comparison with Arabidopsis plants tissues. *Journal of Stress Physiology & Biochemistry* 8, 179-192.
- Ataide, L.M.S., Pappas, M.L., Schimmel, B.C.J., Alba, J.M., Orenes, A., Janssen, A., Pallini, A., Schuurink, R.C., Kant, M.R. 2016 Induced plant-defenses suppress herbivore reproduction but also constrain predation of their offspring. *Plant Science* 252, 300-310.
- Atamian, H.S., Harmer S.L. 2016 Circadian regulation of hormone signaling and plant physiology. *Plant Molecular Biology* 91, 691–702.
- Atkinson, N.J., Jain, R., Urwin, P.E. 2015. The response of plants to simultaneous biotic and abiotic stress. In: Mahalingam R. (ed.), *Combined Stresses in Plants* pp 181-201, Springer International Publishing, Switzerland.
- Azandémè-Hounmalon, G.Y., Fellous, S., Kreiter, S., Fiaboe, K.K.M., Subramanian, S., Kungu, M., Martin, T. 2014 Dispersal behavior of *Tetranychus evansi* and *T. urticae* on tomato at several spatial scales and densities: Implications for integrated pest management. *PLoS ONE* 9, e95071.
- Bai, Y., Lindhout, P. 2007 Domestication and breeding of tomatoes: what have we gained and what can we gain in the future. *Annals of Botany* 100, 1085-1094.
- Bartels, D., Sunkar, R. 2005 Drought and salt tolerance in plants. *Critical Reviews in Plant Sciences* 24, 23–58.
- Bauer, D., Biehler, K., Fock, H., Carrayol, E., Hirel, B., Migge, A., Becker, T.W. 1997 A role for cytosolic glutamine synthetase in the remobilization of leaf nitrogen during water stress in tomato. *Physiologia Plantarum* 99, 241–242.

- Bebber, D.P. 2015 Range-expanding pests and pathogens in a warming world. *Annual Review of Phytopathology* 53, 335-356.
- Bensoussan, N., Santamaria M.E., Zhurov, V., Diaz, I., Grbic, M., Grbi, V. 2016 Plant-herbivore interaction: dissection of the cellular pattern of *Tetranychus urticae* feeding on the host plant. *Frontiers in Plant Science* 7, 1105.
- Blum, A. 2017 Osmotic adjustment is a prime drought stress adaptive engine in support of plant production. *Plant, Cell & Environment* 40, 4–10.
- Boubou, A., Migeon, A., Roderick, G. K., Navajas, M. 2011 Recent emergence and worldwide spread of the red tomato spider mite, *Tetranychus evansi*: genetic variation and multiple cryptic invasions. *Biological Invasions* 13, 81-92.
- Boubou, A., Migeon, A., Roderick, G.K., Auger, P., Cornuet, J., Magalhaes, S., Navajas, M. 2012. Test of colonization scenarios reveals complex invasion history of the red tomato spider mite *Tetranychus evansi*. *PLoS ONE* 7, e35601.
- Boughton, A.J., Hoover, K., Felton, G.W. 2006. Impact of chemical elicitor applications on greenhouse tomato plants and population growth on the green peach aphid, *Myzus persicae*. *Entomologia Experimentalis et Applicata* 120, 175–188.
- Bradford, M.M., 1976 A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72, 248–254.
- Brodbeck, B., Strong, D. 1987 Amino acid nutrition of herbivorous insects and stress to host plants. In: Barbosa, P., Schultz, J., (eds.), *Insect Outbreaks: Ecological and Evolutionary Perspectives*, pp 347-364. Academic Press, New York, USA.
- Butler, G.D. 1955 The effect of alfalfa irrigation treatments on the two-spotted mite in alfalfa. *Journal of Economic Entomology* 48, 221-222.
- Cantore, V., Lechkar, O., Karabulut, E., Sellami, M.H., Albrizio, R., Boari, F., Stellacci, A.M., Todorovic, M. 2016 Combined effect of deficit irrigation and strobilurin application on yield, fruit quality and water use efficiency of “cherry” tomato (*Solanum lycopersicum* L.). *Agricultural Water Management* 167, 53-61.
- Carrillo, L., Martinez, M., Ramessar, K., Cambra, I., Castañera, P., Ortego, F., Diaz, I. 2011 Expression of a barley cystatin gene in maize enhances resistance against phytophagous mites by altering their cysteine-proteases. *Plant Cell Reports* 30, 101-112.
- Chaves, M.M., Maroco, J.P., Pereira, J.S. 2003 Understanding plant responses to drought- from genes to the whole plant. *Functional Plant Biology* 30, 239-264.
- Chaves, M.M., Oliveira, M.M. 2004 Mechanisms underlying plant resilience to water deficits: prospects for water-saving agriculture. *Journal of Experimental Botany* 55, 2365–2384.
- Chen, H., Wilkerson, CG., Kuchar, J.A., Phinney, B.S., Howe, G.A. 2005 Jasmonate-inducible plant enzymes degrade essential amino acids in the herbivore midgut. *Proceedings of the National Academy of Sciences of the USA* 102, 19237–19242.

- Chen, H., Gonzales-Vigil, E., Wilkerson, Howe, G.A. 2007 Stability of plant defense proteins in the gut of insect herbivore. *Plant Physiology* 143, 1954-1967.
- Claussen, W. 2005 Proline as a measure of stress in tomato plants. *Plant Science* 168, 241-248.
- Clauw, P., Coppens, F., De Beuf, K., Dhodt, S., Van Daele, T., Maleux, K., Storme, V., Clement, L., Gonzalez, N., Inze, D. 2015 Leaf responses to mild drought stress in natural variants of Arabidopsis. *Plant Physiology* 167, 800-816.
- Constabel, C.P., Barbehenn, R. 2008 Defensive roles of polyphenol oxidase in plants. In: Schaller, A. (ed) Induced plant resistance to herbivory, pp 253–269. Springer Science Business Media.
- Cornelissen, T., Wilson-Fernandes, G., Vasconcellos-Neto, J. 2008 Size does matter: Variation in herbivory between and within plants and the plant vigor hypothesis. *Oikos* 117, 1121-1130.
- Cornelissen, T. 2011 Climate change and its effects on terrestrial insects and herbivory patterns. *Neotropical Entomology* 40, 155-163.
- Cruz de Carvalho, M.H. 2008 Drought stress and reactive oxygen species: Production, scavenging and signalling. *Plant Signal Behaviour* 3, 156-165.
- Dabrowski, Z.T., Bielak, B. 1978 Effect of some plant chemical compounds on the behaviour and reproduction of spider mites (Acarina: Tetranychidae). *Entomologia Experimentalis et Applicata* 24, 317–326.
- Daszkowska-Golec, A., Szarejko, I. 2013 Open or close the gate – Stomata action under the control of phytohormones in drought stress conditions. *Frontiers in Plant Science* 4, 138.
- DeLucia, E.H., Nabity, P.D., Zavala, J.A., Berenbaum, M.R. 2012 Climate change: resetting plant-insect interaction. *Plant Physiology* 160, 1677-1685.
- De Micco V., Aronne G. 2012 Morpho-anatomical traits for plant adaptation to drought. In: Aroca R.(Ed.), Plant Responses to Drought Stress, pp 37–61. Springer, Heidelberg, Germany.
- De Ollas, C., Dodd, I.C. 2016 Physiological impacts of ABA-JA interactions under water limitation. *Plant Molecular Biology* 91, 641-650.
- Dermauw, W., Wybouw, N., Rombauts, S., Menten, B., Vontas, J., Grbic, M., Clark, R. M., Feyereisen, R., Van Leeuwen, T., 2013 A link between host plant adaptation and pesticide resistance in the polyphagous spider mite Tetranychus urticae. *Proceedings of the National Academy of Sciences of the USA* 110, 113-122.
- Diaz-Mendoza M., Velasco-Arroyo B., González-Melendi P., Martínez M., Díaz I. 2014 C1A cysteine protease–cystatin interactions in leaf senescence. *Journal of Experimental Botany* 65, 3825–3833.
- Dilley, M., Chen, R. S., Deichmann, U., Lerner-Lam, A. L., Arnold, M., Agwe, J., Buys, P., Kjevstad, O., Lyon, B., Yetmen, G. 2005. Natural Disaster Hotspots: A Global Risk Analysis. Washington, DC: World Bank.

- Doares, S.H., Syrovets, T., Weiler, E.W., Ryan, C.A. 1995 Oligogalacturonides and chitosan activate plant defensive genes. *Proceedings of the National Academy of Sciences of the USA* 92, 4095-4098.
- Doherty, H.M., Selvendran, R.R., Bowles, D.J. 1988 The wound response of tomato plants can be inhibited by aspirin and related hydroxy-benzoic acids. *Physiological and Molecular Plant Pathology* 33, 377-384.
- Duso, C., Castagnoli, M., Simoni, S., Angeli, G. 2010 The impact of eriophyoids on crops: recent issues on *Aculops schlechtendali*, *Calepitrimerus vitis* and *Aculops lycopersici*. *Experimental & Applied Acarology* 51, 151-168.
- Eckardt, N.A. 2015 The plant cell reviews dynamic aspects of plant hormone signaling and crosstalk. *The Plant Cell* 27, 1-2.
- El Oirdi., M., Abd El Rahman, T., Rigano, L., El Hadrami, A., Rodriguez, M.C., Daayf, F., Vojnov, A., Bouarab, K. 2011. *Botrytis cinerea* manipulates the antagonistic effects between immune pathways to promote disease development in tomato. *Plant Cell* 23, 2405-2421.
- English-Loeb, G.M. 1989. Nonlinear responses of spider mites to drought-stressed host plants. *Ecological Entomology* 14, 45-55.
- English-Loeb, G.M., 1990 Plant drought stress and outbreaks of spider mites: a field test. *Ecology* 71, 1401-1411.
- English-Loeb, G., Stout, M.J., Duffey, S.S. 1997 Drought stress in tomatoes: Changes in plant chemistry and potential nonlinear consequences for insect herbivores. *Oikos*, 79, 456-468.
- EPPO 2004 European and Mediterranean Plant Protection Organization. Reporting Service no. 05 – 2004. Num. article: 2004/080. <https://gd.eppo.int/reporting/article-1601>.
- Errard, A., Ulrichs, C., Kühne, S., Mewis, I., Mishing, N., Maul, R., Drungowski, M., Parolin, P., Scheiner, M., Baldermann, S. 2016 Metabolite profiling reveals a specific response in tomato to predaceous *Chrysoperla carnea* larvae and herbivore(s)-predator interactions with the generalist pests *Tetranychus urticae* and *Myzus persicae*. *Frontiers in Plant Science* 7, 1256.
- ESRYCE 2015 Encuesta sobre superficies y rendimientos de cultivos. Ministerio de Agricultura, Alimentación y Medio Ambiente. Secretaría General Técnica. Subdirección General de Estadística. Madrid
- Ferragut, F., Garzon-Luque, E., Pekas, A. 2013 The invasive spider mite *Tetranychus evansi* (Acari: Tetranychidae) alters community composition and host-plant use of native relatives. *Experimental & Applied Acarology* 60, 321-341.
- Ferrero, M., Gigot, C., Tixier, M.S., Van Houten, Y.M., Kreiter, S. 2010 Egg hatching response to a range of air humidities for six species of predatory mites. *Entomologia Experimentalis et Applicata* 135, 237-244.
- Fitter, A.H., Hay, R.K.M. 2002 Environmental physiology of plants. Academic Press, London UK.

- Foolad, M.R. 2007 Current status of breeding tomatoes for salt and drought tolerance. In: Jenks, M.A., Hasegawa, P.M., Jain, S.M. (Eds.) *Advances in molecular breeding toward drought and salt tolerant crops*, pp 679-700. Springer, Netherlands.
- Fry, J.D. 1989 Evolutionary adaptation to host plants in a laboratory population of the phytophagous mite *Tetranychus urticae* Koch. *Oecologia* 81, 559–565.
- Fujita, M., Fujita, M., Noutoshi, Y., Takahashi, F., Narusaka, Y., Yamaguchi-Shinozaki, K., Shinozaki, K. 2006 Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signalling networks. *Current Opinion in Plant Biology* 9, 436-442.
- Fullana-Pericas, M., Conesa, M.A., Soler, S., Ribas-Carbo, M., Granell, A., Galmes, J., 2016 Variations of leaf morphology, photosynthetic traits and water-use efficiency in Western-Mediterranean tomato landraces. *Photosynthetica*, In press.
- Galmes, J., Conesa, M.A., Ochogavia, J.M., Perdomo, J.A., Francis, D.M., Ribas-Carbo, M., Save, R., Flexas, J., Medrano, H., Cifre, J. 2011 Physiological and morphological adaptations in relation to water use efficiency in Mediterranean accessions of *Solanum lycopersicum*. *Plant, Cell & Environment* 34, 245-260.
- Galmes, J., Conesa, M.A., Ochogavia, J.M., Perdomo, J.A., Francis, D.M., Ribas-Carbo, M., Save, R., Flexas, J., Medrano, H., Cifre, J. 2011 Physiological and morphological adaptations in relation to water use efficiency in Mediterranean accessions of *Solanum lycopersicum*. *Plant Cell & Environment* 34, 245-260.
- Galmes, J., Ochogavia, J.M., Gago, J., Roldan, E.J., Cifre, J., Conesa, M.A. 2013 Leaf responses to drought stress in Mediterranean accessions of *Solanum lycopersicum*: anatomical adaptations in relation to gas exchange parameters. *Plant, Cell & Environment* 36, 920-935.
- Gamir, J., Darwiche, R., Van't Hof, P., Choudhary, V., Stumpe, M., Schneiter, R., Mauch, F. 2016 The sterol-binding activity of PATHOGENESIS-RELATED PROTEIN 1 reveals the mode of action of an antimicrobial protein. *The plant journal*, In press.
- Gillman, J. H., Rieger, M. W., Dirr, M. A., Braman, S. K. 1999 Drought stress increases densities but not populations of two-spotted spider mite on *Buddleia davidii* 'Pink Delight'. *HortScience* 34, 280-282.
- Gispert, M.C., Perring, T.M., de Lara, C.Z., Cazares, C.L., 1989 Efecto del riego en la fluctuación poblacional del acaro del tomate (*Aculops lycopersici* [Massee]). *Agrociencia* 76, 153-165.
- Glas, J.J. 2014 Consequences of russet mite-induced tomato defences for community interactions. PhD thesis. University of Amsterdam, The Netherlands.
- Glas, J.J., Schimmel, C.J., Alba, J.M., Escobar-Bravo, R., Schuurink, R.C., Kant, M. 2012 Plant glandular trichomes as target for breeding or engineering resistance to herbivores. *International Journal of Molecular Sciences* 13, 17077-17103.
- Glas, J.J., Alba, J.M., Simoni, S., Villarroel, C.A., Stoops, M., Schimmel, B.C.J., Schuurink, R.C., Sabelis, M.W., Kant, M. 2014 Defense suppression benefits herbivores that have a monopoly on their feeding site but can backfire within natural communities. *BMC Biology* 12, 98.

- Godfray, H.J., Beddington, J.R., Crute, I.R., Haddad, L., Lawrence, D., Muir, J.F., Pretty, J., Robinson, S., Thomas, S.M., Toulmin, C. 2010 Food security: The challenge of feeding 9 billion people. *Science* 327, 812-818.
- Golldack, D., Li C., Mohan, H., Probst, N. 2014 Tolerance to drought and salt stress in plants: Unraveling the signaling networks. *Frontiers in Plant Science* 5, 151.
- Gomez, S., Steinbrenner, A.D., Osorio S., Schueller, M., Ferrieri, R.A., Fernie, A.R., Orians, C.M. 2012 From shoots to roots: transport and metabolic changes in tomato after simulated feeding by a specialist lepidopteran. *Entomologia Experimentalis et Applicata* 144, 101-111.
- Gornall, J., Betts, R., Burke, E., Clark, R., Camp, J., Willett, K., Wiltshire, A. 2010 Implications of climate change for agricultural productivity in the early twenty-first century. *Philosophical Transactions of the Royal Society B* 365, 2973-2989.
- Gotoh, T., Bruin, J., Sabelis, M.W., Menken, S.B.J., 1993 Host race formation in *Tetranychus urticae*: genetic differentiation, host plant preference and mate choice in a tomato and a cucumber strain. *Entomologia Experimentalis et Applicata* 68, 171–178.
- Gotoh, T., Sugimoto, N., Pallini, A., Knapp, M., Hernandez-Suarez, E., Ferragut, F., Ho, C.C., Migeon, A., Navajas, M., Nachman, G. 2010 Reproductive performance of seven strains of the tomato red spider mite *Tetranychus evansi* (Acari: Tetranychidae) at five temperatures. *Experimental & Applied Acarology* 52, 239–259.
- Gould, F. 1978 Resistance of cucumber varieties to *Tetranychus urticae*: Genetic and environmental determinants. *Journal of Economic Entomology* 71, 680-683.
- Grbic, M., Van Leeuwen, T., Clark, R. M., Rombauts, S., Rouze, P., Grbic, V., Osborne, E. J., Dermauw, W., Ngoc, P. C., Ortego, F., Hernandez- Crespo, P., Diaz, I., Martinez, M., Navajas, M., Sucena, E., Magalhaes, S., Nagy, L., Pace, R. M., Djuranovic, S., Smagghe, G., Iga, M., Christiaens, O., Veenstra, J. A., Ewer, J., Villalobos, R. M., Hutter, J. L., Hudson, S. D., Velez, M., Yi, S. V., Zeng, J., Pires-daSilva, A., Roch, F., Cazaux, M., Navarro, M., Zhurov, V., Acevedo, G., Bjelica, A., Fawcett, J. A., Bonnet, E., Martens, C., Baele, G., Wissler, L., Sanchez-Rodriguez, A., Tirry, L., Blais, C., Demeestere, K., Henz, S. R., Gregory, T. R., Mathieu, J., Verdon, L., Farinelli, L., Schmutz, J., Lindquist, E., Feyereisen, R., Van de Peer, Y. 2011 The genome of *Tetranychus urticae* reveals herbivorous pest adaptations. *Nature* 479, 487-492.
- Guo, H., Sun, Y., Li, C., Tong, B., Harris, M., Zhu-Salzman, K., Ge, F. 2013 Pea aphid promotes amino acid metabolism both in *Medicago truncatula* and bacteriocytes to favor aphid population growth under elevated CO₂. *Global Change Biology* 19, 3210-3223.
- Gutbrodt, B., Mody, K., Dorn S. 2011 Drought changes plant chemistry and causes contrasting responses in lepidopteran herbivores. *Oikos*, 120, 1732-1740.
- Hacham, Y., Avraham, T., Amir, R. 2002 The N-terminal region of Arabidopsis cystathionine γ -synthase plays an important regulatory role in methionine metabolism. *Plant Physiology* 128, 454–462.

- Haque, M.M., Kawai, A. 2003 Effect of temperatura on development and reproduction of the tomato russet mite, *Aculops lycopersici* (Massee) (Acari: Eriophyidae). *Applied Entomology & Zoology* 38, 97-101.
- Han, P., Dong, Y.C., Lavoit, A.V., Adamowicz, S., Bearez, P., Wajnberg, E., Desneux, N. 2015 Effect of plant nitrogen and water status on the foraging behavior and fitness of an omnivorous arthropod. *Ecology and Evolution* 5, 5468-5477.
- Han, P., Lavoit, A.V., Le Bot, J., Amiens-Desneux, N. 2014 Nitrogen and water availability to tomato plants triggers bottom-up effect on the leaf miner *Tuta absoluta*. *Scientific Reports* 4, 4455.
- Harb, A., Krishnan, A., Ambavaram, M.M.R., Pereira, A. 2010 Molecular and physiological analysis of drought stress in Arabidopsis reveals early responses leading to acclimation in plant growth. *Plant Physiology* 154, 1254-1271.
- Hayat, S., Hasan, S.A., Fariduddin, Q., Ahmad, A. 2008 Growth of tomato (*Lycopersicon esculentum*) in response to salicylic acid under water stress. *Journal of Plant Interactions* 3, 297-304.
- Hollingsworth, C.S., Berry, R.E. 1982 Twospotted spider mite (Acari: Tetranychidae) in peppermint: population dynamics and influence of cultural practices. *Environmental Entomology* 11, 1280-1284.
- Howe, G.A., Jander, G. 2008 Plant immunity to insect herbivores. *Annual Review of Plant Biology* 59, 41-66.
- Howe, G.A., Herde, M. 2015 Interaction of plant defense compounds with the insect gut: new insights from genomic and molecular analysis. *Current Opinion in Insect Science* 9, 62-68.
- Hu, H., Xiong, L. 2014 Genetic engineering and breeding of drought-resistant crops. *Annual Review of Plant Biology* 65, 715-741.
- Huang, Y., Xiao, B., Xiong, L. 2007 Characterization of a stress responsive proteinase inhibitor gene with positive effect in improving drought resistance in rice. *Planta* 226, 73-85.
- Huberty, A.E., Denno, R.E. 2004 Plant water stress and its consequences for herbivorous insects: a new synthesis. *Ecology* 85, 1383-1398.
- Hummel, I., Pantin, F., Sulpice, R., Piques, M., Rolland, G., Dauzat, M., Christophe, A., Pervent, M., Bouteillé, M., Stitt, M., Gibon, Y., Muller, B. 2010 Arabidopsis plants acclimate to water deficit at low cost through changes of carbon usage: An integrated perspective using growth, metabolite, enzyme, and gene expression analysis. *Plant Physiology* 154, 357-372.
- Inbar, M., Doostdar, H., Mayer, R.T. 2001 Suitability of stressed and vigorous plants to various insect herbivores. *Oikos* 94, 228-235.
- Iovieno, P., Punzo, P., Guida, G., Mistretta, C., Van Oosten, M.J., Nurcato, R., Bostan, H., Colantuono, C., Costa, A., Bagnaresi, P., Chiusano, M.L., Albrizio, R., Giorio, P., Batelli, G., Grillo, S. 2016 Transcriptomic changes drive physiological responses to

- progressive drought stress and rehydration in tomato. *Frontiers in Plant Science* 7, 371.
- IPCC, 2013 Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (eds.). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp.
- Jayasinghe, G.G., Mallik, B., 2010 Growth stage based economic injury levels for two spotted spider mite, *Tetranychus urticae* Koch (Acari, Tetranychidae) on tomato, *Lycopersicon esculentum* Mill. *Tropical Agricultural Research* 22, 54-65.
- Johnson, S.N., Ryalls, J.M.W., Karley, A.J. 2014 Global change and crop resistance to aphids: Contrasting responses of lucerne genotypes to elevated atmospheric carbon dioxide. *Annals of Applied Biology* 165, 62-72.
- Jonckheere, W., Dermauw, W., Zhurov, V., Wybpuw, N., Van den Bulcke, J., Villarroel, C.a., Greenhalgh, R., Grbic, M., Schuurink, R.C., Tirry, L., Baggerman, G., Clark, R.M., Kant, M.R., Vanholme, B., Menschaert, G., Van Leeuwen, T. 2016 The salivary protein repertoire of the polyphagous spider mite *Tetranychus urticae*: a quest for effectors. *Molecular and Cellular Proteomics* 15, 3594-3613.
- Kamau, A.W., Mueke J.M., Khaemba B.M., 1992 Resistance of tomato varieties to the tomato russet mite, *Aculops lycopersici* (Massee) (Acarina: Eriophyidae). *Insect Science and its Application* 13(3), 351-356.
- Kang, S.G., Choi, J.H., Suh, S.G. 2002 A Leaf-specific 27 kDA protein of potato Kunitz-type proteinase inhibitor is induced in response to abscisic acid, ethylene, methyl jasmonate, and water deficit. *Molecules & Cells* 13, 144-147.
- Kant, M.R., Ament, K., Sabelis, M.W., Haring, M.A., Schuurink, R.C. 2004 Differential timing of spider mite-induced direct and indirect defenses in tomato plants. *Plant Physiology* 135, 483-495.
- Kant, M.R., Sabelis, M.W., Haring, M.A., Schuurink, R.C., 2008. Intraspecific variation in a generalist herbivore accounts for induction and impact of host-plant defenses. *Proceedings of the Royal Society B- Biological Science* 275, 443-452.
- Kant, M., Jonckheere, W., Knecht, B., Lemos, F., Liu, J., Schimmel, B.C.J., Villarroel, C.A., Ataide, L.M.S., Dermauw, W., Glas, J.J., Egas, M., Janssen, A., Van Leeuwen, T., Schuurink, R.C., Sabelis, M.W., Alba, J.M. 2015 Mechanisms and ecological consequences of plant defence induction and suppression in herbivore communities. *Annals of Botany* 115, 1015-1051.
- Kasap, I. 2004 Effect of apple cultivar and of temperature on the biology and life table parameters of the two-spotted spider mite *Tetranychus urticae*. *Phytoparasitica*, 32, 73-82.
- Kazan, K., Manners, J.M. 2013 MYC2: The Master in Action. *Molecular Plant* 6, 686-703.
- Kazan, K., Lyons, R. 2014 Intervention of phytohormone pathways by pathogen effectors. *Plant Cell* 26, 2285-2309.

- Keifer, H.H. 1946 A review of North American eriophyid mites. *Journal of Economic Entomology* 39, 563-570.
- Kennedy, G.G. 2003 Tomato, pests, parasitids and predators, tritrophic interactions involving the genus *Lycopersicon*. *Annual Review of Entomology* 48, 51-72.
- Khodakovskaya, M., Sword, C., Wu, Q., Perera I.Y., Boss, W.,F., Brown, C.S., Sederoff, W. 2010 Increasing inositol (1,4,5)-trisphosphate metabolism affects drought tolerance, carbohydrate metabolism and phosphate-sensitive biomass increases in tomato. *Plant Biotechnology Journal* 8, 170-183.
- Koyama, Y., Yao, I., Akimoto, S.A. 2004 Aphid galls accumulate high concentrations of amino acids: a support for the nutrition hypothesis for gall formation. *Entomologia experimentalis et applicata* 113, 35–44.
- Krantz, G.W., Lindquist, E.E. 1979 Evolution of phytophagous mites (Acari). *Annual Review of Entomology* 24, 121-158.
- Lange, W.H., Bronson, L. 1981 Insect pests of tomato. *Annual Review of Entomology* 26, 345-371.
- Levitt, J. 1980 Responses of Plants to Environmental Stresses. New York: Academic Press 607 pp.
- Li, C., Williams, M.M., Loh, Y.T., Lee, G.I., Howe, G.A. 2002 Resistance of cultivated tomato to cell content-feeding herbivores is regulated by the octadecanoid-signaling pathway. *Plant Physiology* 130, 494-503.
- Lin, T., Zhu, G., Zhang, J., Xu, X., Yu, Q., Zheng, Z., Zhang, Z., Lun, Y., Li, S., Wang, X., Huang, Z., Li, J., Zhang, C., Wang, T., Zhang, Y., Wang, A., Zhang, Y., Lin, K., Li, C., Xiong, G., Xue, Y., Mazzucato, A., Causse, M., Fei, Z., Giovannoni, J.J., Chetelat, R.T., Zamir, D., Stadler, T., Li, J., Ye, Z., Du, Y., Huang, S. 2014 Genomic analyses provide insights into the history of tomato breeding. *Nature Genetics* 46, 1220-1226.
- Lindquist, E.E. 1985 Diagnosis and Phylogenetics relationships. In: Helle, W, Sabelis, M.W. (eds.) Spider mites their biology, natural enemies and control Volume 1A, pp 63-75. Elsevier, Amsterdam, The Netherlands.
- Lindquist, E.E. 1996 External anatomy and systematics. In: Lindquist, E.E. Sabelis, M.W. and Bruin, J. (eds.) Eriophyoid mites- Their biology, natural enemies and control, pp 3-31. Elsevier Science Publisher Amsterdam, The Netherlands.
- Luedeling, E., Steinmann, K.P., Zhang, M., Brown, P.H., Grant, J., Girvetz, E.H., 2011. Climate change effects on walnut pests in California. *Global Change Biology* 17, 228–238.
- Macke E, Magalhães S, Khan HDT, Luciano A, Frantz A, Facon B, Olivieri I. 2010. Sex allocation in haplodiploids is mediated by egg size: evidence in the spider mite *Tetranychus urticae* Koch. *Proceedings of the Royal Society B-Biological Science* 278, 1054-1063.
- Magalhaes, S., Blanchet, E., Egas, M., Olivieri, I. 2009 Are adaptation costs necessary to build up a local adaptation pattern?. *BMC Evolutionary Biology* 9,182.

- Maness, N., 2010 Extraction and analysis of soluble carbohydrates. In, Sunkar, R. (ed.). Plant Stress Tolerance, Methods in Molecular Biology, pp. 341–370. Springer Science+Business Media.
- Marinosci, C., Magalhaes, S., Macke, E., Navajas, M., Carbonell, D., Devaux, C., Olivieri, I. 2015 Effects of host plant on life-history traits in the polyphagous spider mite *Tetranychus urticae*. *Ecology and Evolution* 5, 3151-3158.
- Martel, C., Zhurov, V., Navarro, M., Martinez, M., Cazaux, M., Auger, P., Migeon, A., Santamaria, M. E., Wybouw, N., Diaz, I., Van Leeuwen, T., Navajas, M., Grbic, M., Grbic, V. 2015 Tomato whole genome transcriptional response to *Tetranychus urticae* identifies divergence of spider mite-induced responses between tomato and Arabidopsis. *Molecular Plant Microbe Interaction* 28, 343-361.
- Mattson, W., Haack, R. 1987 The role of drought in outbreaks of plant-eating insects. *BioScience* 37, 110–118.
- Maxmen, A. 2013 Under attack. *Nature* 501, S15-S17.
- Mayer, A.M. 2006 Polyphenol oxidases in plants and fungi: Going places? A review. *Phytochemistry* 67, 2318-2331.
- Meck, E.D., Kennedy, G.G., Walgenbach, J.F. 2013 Effect of *Tetranychus urticae* (Acari: Tetranychidae) on yield, quality, and economics of tomato production. *Crop Protection* 52, 84-90.
- Mellors, W.K., Allegro, A., Hsu, A.N. 1984 Effects of carbofuran and water stress on growth of soybean plants and twospotted spider mite (Acari: Tetranychidae) populations under greenhouse conditions. *Environmental Entomology* 13, 561-567.
- Meynard, C.N., Migeon A., Navajas, M. 2013 Uncertainties in predicting species distributions under climate change: A case study using *Tetranychus evansi* (Acari: Tetranychidae), a widespread agricultural pest. *PLoS ONE* 8, e66445.
- Michalska, K., Skoracka, A., Navia, D., Amrine, J.W. 2010 Behavioural studies on eriophyoid mites: an overview. *Experimental & Applied Acarology* 51, 31-59.
- Migeon, A., Dorkeld, F. 2006-2016 Spider Mites Web: a comprehensive database for the Tetranychidae. <http://www.montpellier.inra.fr/CBGP/spmweb>
- Migeon, A., Ferragut, F., Escudero-Colomar, L. A., Fiaboe, K., Knapp, M., de Moraes, G. J., Ueckermann, E., Navajas, M. 2009 Modelling the potential distribution of the invasive tomato red spider mite, *Tetranychus evansi* (Acari: Tetranychidae). *Experimental and Applied Acarology* 48, 199-212.
- Migeon, A., Auger, P., Hufbauer, R., Navajas, M. 2015 Genetic traits leading to invasiveness: plasticity in cold hardiness explains current distribution of an invasive agricultural pest, *Tetranychus evansi* (Acari: Tetranychidae). *Biological Invasions* 17, 2275-2285.
- Miller, G., Suzuki, N., Ciftci-Yilmaz, S., Mittler, R. 2010 Reactive oxygen species homeostasis and signalling during drought and salinity stresses. *Plant Cell & Environment* 33, 453–467.

- Mir, R.R., Zaman-Allah, M., Sreenivasulu, N., Trethowan, R., Varshney, R.K. 2012 Integrated genomics, physiology and breeding approaches for improving drought tolerance in crops. *Theoretical and Applied Genetics* 125, 625-645.
- Mishra, K.B., Iannacone, R., Petrozza, A., Mishra, A., Armentano, N., La Vecchia, G., Trtílek, M., Cellini, F., Nedbal, L. 2012 Engineered drought tolerance in tomato plants is reflected in chlorophyll fluorescence emission. *Plant Science* 182, 79-86.
- Mody, K., Eichenberger, D., Dorn, S. 2009 Stress magnitude matters: Different intensities of pulsed water stress produce non-monotonic resistance responses of host plants to insect herbivores. *Ecological Entomology* 34, 133-143.
- Mosolov, V.V., Valueva, T.A., Inhibitors of proteolytic enzymes under abiotic stress in planta (review). *Applied Biochemistry and Microbiology* 47, 453-459.
- Mueller, N.D., Gerber, J.S., Johnston, M., Ray, D.K., Ramankutty, N., Foley, J.A. 2012 Closing yield gaps through nutrient and water management. *Nature* 490, 254-257.
- Muñoz-Espinoza, V.A., López-Climent, M.F., Casaretto, J.A., Gómez-Cadenas, A. 2015 Water stress responses on tomato mutants impaired hormone biosynthesis reveal abscisic acid, jasmonic acid and salicylic acid interactions. *Frontiers in Plant Science* 6, 997.
- Nachappa, P., Margolies, D.C., Nechols, J.R., Whitfield, A.E., Rotenberg, D. 2013 Tomato Spotted Wilt Virus benefits a non-vector arthropod, *Tetranychus urticae* by modulating different plant responses in tomato. *PLoS ONE* 8, e75909.
- Nakashima, K., Yamaguchi-Shinozaki, K. 2013. ABA signaling in stress-response and seed development. *Plant Cell Reports* 32, 959-970.
- Nankishore, A., Farrell, A.D. 2016 The response of contrasting tomato genotypes to combined heat and drought stress. *Journal of Plant Physiology* 202, 75-82.
- Nansen, C., Sidumo, A.J., Capareda, S. 2010 Variogram analysis of hyperspectral data to characterize the impact of biotic and abiotic stress of maize plants and to estimate biofuel potential. *Applied Spectroscopy* 64, 627-636.
- Navajas, M., de Moraes, G.J., Auger, P., Migeon, A. 2013 Review of the invasion of *Tetranychus evansi*: biology, colonization pathways, potential expansion and prospects for biological control. *Experimental Applied Acarology* 59, 43-46.
- Nguyen, D., D'Agostino, N., Tytgat, T.O.G., Sun, P., Lortzing, T., Visser, E.J.W., Cristescu, S.M., Steppuhn, A., Mariani, C., van Dam, N.M., Rieu, I. 2016a Drought and flooding have distinct effects on herbivore induced responses and resistance in *Solanum dulcamara*. *Plant Cell & Environment* 39, 1485-1499.
- Nguyen, D., Rieu, I., Mariani, C., van Dam, N.M. 2016b How plants handle multiple stresses: hormonal interactions underlying responses to abiotic stress and insect herbivory. *Plant Molecular biology* 91, 727-740.
- Nisbet, A. J., Billingsley, P. F. 2000 A comparative survey of the hydrolytic enzymes of ectoparasitic and free-living mites. *International Journal of Parasitology* 30, 19-27.

- Noctor, G., Mhamdi, A., Foyer, C.H. 2014 The roles of reactive oxygen metabolism in drought: not so cut and dried. *Plant Physiology* 164, 1636–1648.
- Nuzzaci, G., Alberti, G., 1996. Internal anatomy and physiology. In: Lindquist, E.E., Sabelis, M.W., Bruin, J. (Eds.), *Eriophyoid mites - Their biology, natural enemies and control*, pp. 101-150. Elsevier Science Publisher, Amsterdam, The Netherlands.
- O'Donnell, P.J., Calvert, C., Atzorn, R., Wasternack, C., Leyser, H.M.O., Bowles, D.J. 1996 Ethylene as a signal mediating the wound response of tomato plants. *Science* 274, 1914–1917.
- Ortego, F. 2012 Physiological adaptations of the insect gut to herbivory. In: Smagghe, G., Diaz, I. (eds.) *Arthropod-Plant Interactions—Novel Insights and Approaches for IPM*. Series: Progress in Biological Control. Volume 14, pp 75–88. Springer.
- Parmesan, C. 2006 Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution and Systematics* 37, 637-639.
- Passardi, F., Cosio, C., Penel, C., Dunald, C. 2005 Peroxidases have more functions than Swiss army knife. *Plant Cell Reports* 24, 255-265.
- Petanovic, R., Kielkiewicz, M. 2010 Plant-erriophyoid mite interactions: cellular biochemistry and metabolic responses induced in mite-injured plants. Part I. *Experimental and Applied Acarology* 51, 61-80.
- Petrozza A., Santaniello, A., Summerer, S., Di Tommaso, G., Di Tommaso, D., Paparelli, E., Piaggese, A., Perata, P., Cellini, F. 2014 Physiological responses to Megafol® treatments in tomato plants under drought stress: A phenomic and molecular approach. *Scientia Horticulturae* 174, 185-192.
- Pieterse, C.M.J., VenderDoes, D., Zamioudis, C., Leon-Reyes, A., VanWees, S.C.M. 2012 Hormonal modulation of plant immunity. *Annual Review of Cell and Developmental Biology* 28, 489-521.
- Pimentel, D. 2009 Pesticides and pest control. In, Peshin, R., Dhawan A.K. (Eds.), *Integrated pest management: innovation-development process*, pp. 83-87. Springer international publishing, Switzerland.
- Rai, K.G., Rai, N.P., Rathaur, S. Kumar, S., Singh M. 2013 Expression of rd29A:AtDREB1A/CBF3 in tomato alleviates drought-induced oxidative stress by regulating key enzymatic and non-enzymatic antioxidants. *Plant Physiology and Biochemistry* 69, 90-100.
- Riahi, E., P. Shishehbor, A. R. Nemati, Z. Saeidi. 2013 Temperature effects on development and life table parameters of *Tetranychus urticae* (Acari: Tetranychidae). *Journal of Agricultural Science & Technology* 15, 661–672.
- Rigano, M.M., Arena, C., Di Matteo, A., Sellitto, S., Frusciante L., Barone, A. 2016 Eco-physiological response to wáter stress of drought-tolerant and drought-sensitive tomato genotypes. *Plant Biosystems* 150, 682-9.
- Ripoll, J., Urban, L., Staudt, M., Lopez-Lauri, F., Bidel, L.P.R., Bertin, N. 2014 Water shortage and quality of fresh fruits-making the most of the unavoidable. *Journal of Experimental Botany* 65, 4097-4117.

- Ritchie, G.A. 2006 Chlorophyll fluorescence: What is it and what do the numbers mean? in: Riley, L.E., Dumroese, R.K., Landis, T.D. (Eds.). Tech. coords. 2006. National Proceedings: Forest and Conservation Nursery Associations—2005. Proc. RMRS-P-43. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. pp: 34-43. Available at: <http://www.rmnr.net/nurseries/publications/proceedings>
- Rivelli, A.R., Trotta, V., Toma, I., Fanti, P., Battaglia, D. 2013 Relation between plant water status and *Macrosiphum euphorbiae* (Hemiptera: Aphididae) population dynamics on three cultivars of tomato. *European Journal of Entomology* 110, 617-625.
- Rodriguez, J.G., Hampton, R.E. 1966 Essential amino acids determined in the two-spotted spider mite, *Tetranychus urticae* Koch (Acarina, Tetranychidae) with glucose-U-C14. *Journal of Insect Physiology* 12, 1209-1216.
- Royalty, R.N., Perring, T.M. 1988 Morphological analysis of damage to tomato leaflets by tomato russet mite (Acari: Eriophyidae) *Journal of Economic Entomology* 8, 816-820.
- Sabelis, M.W., Bruin, J. 1996 Evolutionary Ecology: Life history patterns, food plant choice and dispersal. In *Eriophyoid Mites – Their Biology, Natural Enemies and Control* (eds. Lindquist E.E., Sabelis M.W. & Bruin J.) World Crop Pest Series Vol. 6. pp. 329-366. Elsevier Science Publishers, Amsterdam, The Netherlands.
- Sadras, V.O., Wilson, L.J., Lally, D.A. 1998 Water deficit enhanced cotton resistance to spider mite herbivory. *Annals of Botany* 81, 273-286.
- Sanchez-Rodriguez, E., Rubio-Wilhelmi, M., Cervilla, L.M., Blasco, B., Rios, J.J., Rosales, M.A., Romero, L., Ruiz, J.M. 2010 Genotypic differences in some physiological parameters symptomatic for oxidative stress under moderate drought in tomato plants. *Plant Science* 178, 30-40.
- Sandström, J., Telang, A., Moran, N.A. 2000 Nutritional enhancement of host plants by aphids - a comparison of three aphid species on grasses. *Journal Insect Physiology* 46, 33–40.
- Santamaria, M.E., Cambra, I., Martinez, M., Pozancos, C., González-Melendi, P., Grbic, V., Castañera, P., Ortego, F., Diaz, I. 2012 Gene pyramiding of peptidase inhibitors enhances plant resistance to the spider mite *Tetranychus urticae*. *Plos ONE* 7, e43011.
- Santamaria, M.E., González-Cabrera, J., Martínez, M., Grbic, V., Castañera, P., Díaz, I., Ortego, F. 2015a Digestive proteases in bodies and faeces of the two-spotted spider mite, *Tetranychus urticae*. *Journal of Insect Physiology* 78, 69-77.
- Santamaria, M.E., Arnaiz, A., Diaz-Mendoza, M., Martinez, M., Diaz, I. 2015b Inhibitory properties of cysteine protease pro-peptides from barley confer resistance to spider mite feeding. *PLoS ONE* 10, e0128323.
- Sarmiento, R.A., Lemos, F., Dias, C.R., Kikuchi, W.T., Rodrigues, J.C, Pallini, A., Sabelis, M.W., Janssen A. 2011a A herbivorous mite down-regulates plant defence and produces web to exclude competitors. *PLoS ONE* 6, e23757.

- Sarmiento, R.A., Lemos, F., Bleeker, P.M., Schuurink, R.C., Pallini, A., Oliveira, M.G., Lima, E.R., Kant, M., Sabelis, M.W., Janssen, A. 2011b A herbivore that manipulates plant defence. *Ecology Letters* 14, 229-236.
- Saunyama, I.G.M., Knapp, M. 2003 Effects of pruning and trellising of tomatoes on red spider mite incidence and crop yield in Zimbabwe. *African Crop Science Journal* 11, 269–277.
- Scaraffia, P.Y., Wells, M.A. 2003 Proline can be utilized as an energy substrate during flight of *Aedes aegypti* females. *Journal of Insect Physiology* 49, 591-601.
- Schmidt, L., Schurr, U., Rose, U.S.R. 2009 Local and systemic effects of two herbivores with different feeding mechanisms on primary metabolism of cotton leaves. *Plant Cell & Environment* 32, 893-903.
- Schuman, M.C., Baldwin, I.T. 2016 The layers of plant response to herbivores. *Annual Review of Entomology* 61, 373-394.
- Seagraves, M.P., Riedell, W.E., Lundgren, J.G. 2011 Oviposition preference for water-stressed plants in *Orius insidiosus* (Hemiptera: Anthocoridae). *Journal of Insect Behavior* 24, 132-143.
- Sharma, P., Jha, A.B., Dubey, R.S., Pessarakli, M. 2012 Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *Journal of Botany* 2012, Article ID 217037, 26 pp.
- Showler, A.T. 2013 Water deficit stress-host plant nutrient accumulations and associations with phytophagous arthropods. In, Vahdati, K. (ed.), *Abiotic Stress-Plant Responses and Applications in Agriculture*, pp. 387-410. InTech, Rijeka, Croatia.
- Sinclair, T.R. 2011 Challenges in breeding for yield increase for drought. *Trends in Plant Science* 15, 289-293.
- Singh, A., Singh, S., Singh, I.K. 2016 Recent insights into the molecular mechanism of jasmonate signaling during insect-plant interaction. *Australian Plant Pathology* 45, 123-133.
- Simmons, A.T., Gurr, G.M. 2005 Trichomes of *Lycopersicon* species and their hybrids: effects on pests and natural enemies. *Agricultural and Forest Entomology* 7, 265-276.
- Skirycz, A., Inze, D. 2010 More from less: plant growth under limited water. *Current Opinion in Biotechnology* 21, 1–7.
- Stavrinides, M.C., Daane, K.M., Lampinen, B.D., Mills, N.J. 2010 Plant water stress, leaf temperature, and spider mite (Acari: Tetranychidae) outbreaks in California vineyards. *Environmental Entomology* 39, 1232-1241.
- Steduto, P., Hsiao, T. C., Fereres, E., Raes, D. 2012 Crop yield response to water, FAO Irrigation and drainage paper 66. Food and Agriculture Organization of the United Nations. Rome, Italy.
- Steinbrenner, A.D., Gomez, S., Osorio, S., Fernie, A.R., Orians, C.M. 2011 Herbivore-induced changes in tomato (*Solanum lycopersicum*) primary metabolism: a whole plant perspective. *Journal of Chemical Ecology* 37, 1294-1303.

- Stout, M.J., Workman, J., Duffey, S.S. 1994 Differential induction of tomato foliar proteins by arthropod herbivores. *Journal of Chemical Ecology* 20, 2575-2594
- Stout, M.J., Workman, K.V. Duffey, S.S. 1996 Identity, spatial distribution, and variability of induced chemical responses in tomato plants. *Entomologia Experimentalis et Applicata* 79, 255-271.
- Suzuki, N., Rivero, R.M., Shulaev, V., Blumwald, E., Mittler, R. 2014 Abiotic and biotic stress combinations. *New Phytologist* 203, 32-43.
- Szabados, L., Savouré, A. 2010 Proline: a multifunctional amino acid. *Trends Plant Science* 15, 89-97.
- Tapia, G., Mendez, J., Inostroza, L. 2016 Different combinations of morpho-physiological traits are responsible for tolerance to drought in wild tomatoes *Solanum chilense* and *Solanum peruvianum*. *Plant Biology* 18, 406-416.
- Thaler, J.S., Fidantsef, A.L., Duffey, S.S., Bostock, R.M. 1999 Trade-offs in plant defense against pathogens and herbivores: A field demonstration of chemical elicitors of induced resistance. *Journal of Chemical Ecology* 25, 1597-609.
- Thaler, J.S., Bostock, R.M. 2004 Interaction between abscisic-acid-mediated responses and plant resistance to pathogens and insects. *Ecology* 85, 48-58.
- Thipyapong, P., Melkonian, J., Wolfe, D.W., Steffens, J.C. 2004 Suppression of polyphenol oxidases increases stress tolerance in tomato. *Plant Science* 167, 693-703.
- Thompson, A.J., Andrews, J., Mulholland, B.J., McKee, J.M.T., Hilton, H.W., Horridge, J.S., Farquhar, G.D., Smeeton, R.C., Smillie, I.R.a., Black, C.R., Taylor, I.B. 2007 Overproduction of abscisic acid in tomato increases transpiration efficiency and root hydraulic conductivity and influences leaf expansion. *Plant Physiology* 143, 1095-1917.
- Tulisalo, U. 1971 Free and bound amino acids of three host plant species and various fertilizer treatments affecting the fecundity of the two-spotted spider mite, *Tetranychus urticae* Koch (Acarina, Tetranychidae). *Annales Entomologici Fennici*, 37, 155-163.
- Tylianakis, J.M., Didham, R.K., Bascompte, J., Wardle, D.A. 2008 Global change and species interactions in terrestrial ecosystems. *Ecology letters* 11, 1351 – 1363.
- Vacante, V. 2016 The Handbook of Mites of Economic Plants: Identification, Bio-Ecology and Control. CABI, Wallingford, UK. 872 pp.
- Valencia, A., Bustillo, A.E., Ossa, G.E., Chrispeels, M.J. 2000 α -Amilases of the coffee berry borer (*Hypothenemus hampei*) and their inhibition by two amylase inhibitors. *Insect Biochemistry and Molecular Biology* 30, 207-213.
- Van der Geest, L.P.S. 1985 Physiology and Genetics. In: Helle, W and Sabelis, M.W. (eds.) Spider mites their biology, natural enemies and control Volume 1A, pp 63-75. Elsevier, Amsterdam, The Netherlands.
- Van Houten, Y.M., Glas, J.J., Hoogerbrugge, H., Rothe, J., Blockmans, K.J., Simoni, S., Van Arkel, J., Alba, J.M., Kant, M.R., Sabelis, M.W. 2013 Herbivory-associated

- degradation of tomato trichomes and its impact on biological control of *Aculops lycopersici*. *Experimental and Applied Acarology* 60, 127-138.
- Van Leeuwen, T., Dermauw, W. 2016 The molecular evolution of xenobiotic metabolism and resistance in Chelicerate mites. *Annual Review of Entomology* 61, 475-498.
- Verma, V., Ravindran, P., Kumar, P.P. 2016 Plant hormone-mediated regulation of stress responses. *BMC Plant Biology* 16, 86.
- Verwoerd, T.C., Dekker, B., Hoekema, A. 1989 A small-scale procedure for the rapid isolation of plant RNAs. *Nucleic Acids Research* 17, 2362.
- Verslues, P.E., Agarwal, M., Katiyar-Agarwal, S., Zhu, J., Zhu, J.K. 2006 Methods and concepts in quantifying resistance to drought, salt and freezing abiotic stresses that affects plant water status. *The Plant Journal* 45, 532-539.
- Verslues P.E., Juenger, T.E. 2011 Drought, metabolites, and Arabidopsis natural variation: a promising combination for understanding adaption to water-limited environments. *Current Opinion in Plant Biology* 14, 240-245.
- Villarroel, C.A., Jonckheere, W., Alba, J.M., Glas, J.J., Dermauw, W., Haring, M.A., Van Leeuwen, T., Schuurink, R.C., Kant, M.R. 2016. Salivary proteins of spider mites suppress defenses in *Nicotiana benthamiana* and promote mite reproduction. *Plant Journal* 86, 119-131.
- War, A. R., Paulraj, M. G., Ahmad, T., Buhroo, A. A., Hussain, B., Ignacimuthu, S., Sharma, H.C. 2012 Mechanisms of plant defense against insect herbivores. *Plant Signal Behavior* 7, 1306–1320.
- Wermelinger, B., Oertli, J.J., Baumgartner, J. 1991 Environmental factors affecting the life-tables of *Tetranychus urticae* (Acari: Tetranychidae). III. Host-plant nutrition. *Experimental & Applied Acarology* 12, 259–274.
- White, T.C.R. 1984 The abundance of invertebrate herbivores in relation to the availability of nitrogen in stressed food plants. *Oecologia*, 63, 90-105.
- White, T.C.R. 2009 Plant vigor versus plant stress: A false dichotomy. *Oikos* 118, 807-808.
- World Economic Forum Water Initiative 2009 Managing our future water needs for agriculture, industry, human health and the environment. In Draft for Discussion—Meeting of the World Economic Forum.
- Wybouw, N., Zhurov, V., Martel, C., Bruinsma, K. A., Hendrickx, F., Grbić, V., van Leeuwen, T. 2015 Adaptation of a polyphagous herbivore to a novel host plant extensively shapes the transcriptome of herbivore and host. *Molecular Ecology* 24, 4647-4663.
- Ximénez-Embún, M.G., Ortego, F., Castañera, P. 2016 Drought stressed tomato plants triggers bottom-up effects on the invasive *Tetranychus evansi*. *PLoS ONE* 11, e0145275.
- Ximénez-Embún, M.G., Castañera, P., Ortego, F. 2017 Drought stress in tomato increases the performance of adapted and non-adapted strains of *Tetranychus urticae*. *Journal of Insect Physiology* 96, 73-81.
- Xu Z., Zhou G., Shimizu, H. 2010. Plant response to drought and rewatering. *Plant Signal Behaviour* 5, 649–654.

- Youngman, R.R., Sanderson, J.P., Barnes, M.M. 1988 Life history parameters of *Tetranychus pacificus* McGregor (Acari: Tetranychidae) on almonds under differential water stress. *Environmental Entomology* 17, 488-495.
- Yu, H., Kowalski, S.P., Steffens, J.C. 1992 Comparison of polyphenol oxidase expression in glandular trichomes of *Solanum* and *Lycopersicon* species. *Plant Physiology* 100, 1885–1890.
- Zhang, X., Liu, S., Takano, T. 2008 Two cysteine proteinase inhibitors from *Arabidopsis thaliana*, AtCYSa and AtCYSb, increasing the salt, drought, oxidation and cold tolerance. *Plant Molecular Biology* 68, 131–143.
- Zhou, S., Lou, Y.R., Tzin, V., Jander, G. 2015 Alteration of plant primary metabolism in response to insect herbivory. *Plant Physiology* 169, 1488-1498.
- Zhurov, V., Navarro, M., Bruinsma, K.A., Arbona, V., Santamaria, M.E., Cazaux, M., Wybouw, N., Osborne, E.J., Ens, C., Rioja, C., Vermeirssen, V., Rubio-Somoza, I., Krishna, P., Diaz, I., Schmid, M., Gomez-Cadenas, A., VandePeer, Y., Grbic, M., Clark, R.M., VanLeuween, T., Grbic, V. 2014 Reciprocal responses in the interaction between *Arabidopsis* and the cell-content-feeding chelicerate herbivore spider mite. *Plant physiology* 164, 384-399.
- Zhu-Salzman, K., Luthe, D.S., Felton, G.W. 2008 Arthropod-inducible proteins: broad spectrum defenses against multiple herbivores. *Plant Physiology* 146, 852–858.
- Zhu-Salzman, K., Zeng, R. 2015 Insect response to plant defensive protease inhibitors. *Annual Review of Entomology* 60, 233-252.