

A PREVIOUS HAMSTRING INJURY AFFECTS LIMB DOMINANCE IN SOCCER KICKING

Archit Navandar¹, Santiago Veiga¹, Carlos Garcia¹, Javier Rueda¹, Gonzalo Torres¹, David Chorro¹ and Enrique Navarro¹
Faculty of Sports Sciences, Technical University of Madrid, Madrid, Spain¹

The objective of this study was to study if the influence of limb dominance on kicking is affected by a previous hamstring injury to either limb. Forty-five players (twenty-six females and nineteen males) belonging to elite soccer teams volunteered to take part in the study. They performed 5 instep soccer kicks with either limb into a target 7m away. Differences were found across limb dominance for the uninjured group in peak linear velocities, hip flexion velocity, knee angle at ball impact, and hip moments. However, normal limb dominance characteristics were affected in the presence of a previous hamstring injury. A previous injury prevented the rapid extension of the hamstring muscle fibres, being more predominant in the non-dominant limb. Thus the effects of a previous hamstring injury must be taken into account while dealing with the kicking technique, and the location of the injury on the dominant or non-dominant limb is equally important.

KEY WORDS: INVERSE DYNAMICS, MOTION ANALYSIS, KINEMATICS, KINETICS.

INTRODUCTION: Studies on European soccer accounted injuries to the hamstrings contributed to 37% of all non-contact injuries (Ekstrand, Hägglund, & Waldén, 2011). 22% of players sustained at least one hamstring injury during a season and the number of hamstring strain injuries increasing annually by 4% (Ekstrand, Waldén, & Hägglund, 2016). The dominant leg accounts for a greater frequency of lesions – 54.4% vs 36.5% in the non-dominant leg (Noya, 2015), but no significant differences were observed in the number of hamstring injuries between the dominant and the non-dominant leg (Freckleton & Pizzari, 2013). The hamstring muscle strain-type injury is considered to be a multifactorial problem, and the factors of age and previous hamstring injury must be considered alongside any other risk factors being examined (Freckleton & Pizzari, 2013). The hamstring muscle complex is known to be affected by the kicking skill (Navandar et al., 2015; Navandar, Gulino, Antonio, & Navarro, 2013), and various investigations have shown that limb dominance also influence the kicking skill (Navandar et al., 2016; Nunome, Ikegami, Kozakai, Apriantono, & Sano, 2006). However, no study has looked at the combined effect of limb dominance and previous hamstring injury on the kicking skill. Therefore, the objective of this study was to ascertain if the influence of limb dominance on kicking is affected by a previous hamstring injury to either limb.

METHODS: Participants: Forty-five elite soccer players (females: $n = 26$, age = 22.15 ± 4.50 years, weight = 60.71 ± 9.48 kg; males: $n = 19$, age = 21.16 ± 2.00 years, weight = 71.46 ± 6.22 kg) participated in the study. Players were asked about their preferred leg, playing position and previous injury history in a questionnaire prior to the data capture session. Only players having suffered a grade 1, 2 or 3 hamstring injury which caused them to miss the subsequent match in the last two years were considered as “previously injured”. All players who were injured had suffered a unilateral injury (females: injured in dominant limb = 6, injured in non-dominant limb = 4, uninjured = 16; males: injured in dominant limb = 3, injured in non-dominant limb = 1, uninjured = 15).

Data capture: Twenty-four retro-reflective markers (diameter = 14 mm) were attached to the anatomical landmarks of each participant’s body, and 4 markers being attached to the ball. In a procedure similar to Navandar et al. (2016), a 6 camera Vicon Motion Capture System at 250Hz and a Kistler Force Platform at 1000 Hz were used to capture static and dynamic data in laboratory conditions, on FIFA approved artificial turf. In the dynamic capture, the players

were instructed to kick the stationary ball with their dominant and non-dominant legs at a 1m x 1m target 7m away, as hard as possible using the instep soccer kick, while placing their support leg on a force platform. Kicks were recorded until 5 kicks were performed with each leg meeting the above criteria. Ball velocities were calculated and the trial containing the median value of ball velocity for each limb was selected for further analysis.

Data Processing: A standard inverse dynamics procedure was used to calculate the kinetic and kinematic variables of the kicking and support limb, with the kicking motion being divided into the backswing, leg cocking, leg acceleration and follow through phases (Lees, 2013).

Data Analysis: Kinematic and kinetic variables in the sagittal plane, and normalized phases of kicking were compared in a 3-way ANOVA with the independent factors being limb dominance, previous injury and sex. When a significant effect was observed between the ANOVA levels, multiple comparisons were performed with the Bonferroni correction, and the results were interpreted with 95% confidence intervals (CIs) and effect sizes (as partial η^2 values). The threshold values for small, medium and large effect sizes were set as 0.01, 0.06 and 0.14 respectively (Cohen, 1992; Richardson, 2011).

RESULTS AND DISCUSSION: Differences in the peak linear velocities between the dominant and non-dominant limb were found only for the uninjured group, and not for the injured group (Table 1). Previous studies have consistently stated the superiority of the dominant limb in linear velocities, (Nunome et al., 2006; van den Tillaar & Ulvik, 2014). However, a previous injury prevented large differences from occurring. These differences can be explained on the sequence of velocities, which follow a proximal to distal sequence (Lees & Nolan, 1998), and the differences that occur at the different phases of the kick.

Table 1
Comparison of peak linear velocities

		Ball Velocity (m/s)	Toe velocity (m/s)	Ankle velocity (m/s)	Knee Velocity (m/s)
Uninjured group	ND	23.97 ± 3.40**	19.15 ± 2.48**	15.85 ± 1.77**	8.74 ± 1.04**
	D	26.75 ± 2.74**	21.08 ± 2.29**	17.19 ± 1.67**	9.53 ± 0.91**
Injured group	ND	24.00 ± 2.00	19.00 ± 2.55	16.00 ± 1.41	8.80 ± 0.84
	D	25.44 ± 2.74	20.11 ± 1.76	16.44 ± 1.33	9.00 ± 0.87

*ND: Non-dominant limb kick, D: dominant limb kick. * Significant difference between the two groups with $p < 0.05$, ** Significant difference between the two groups with $p < 0.001$.*

The end of the backswing phase is defined as the instance when the hip reaches peak extension angle (Lees, 2013), and a peak flexion moment (defined in literature as the soccer paradox (Lees, Asai, Bull-Andersen, Nunome, & Sterzing, 2010)). At this instance, in the uninjured group, the dominant limb kicks have a greater hip flexion moment as compared to non-dominant limb kicks (Table 2). However, this difference is not seen in the previously injured group, with the values being slightly higher (small and medium differences in values between injured (partial $\eta^2 = 0.01$) and uninjured group (partial $\eta^2 = 0.06$)). Thus, previously injured players applied a greater moment in order to restrict the hip extension, when the hamstring muscle fibres were, theoretically, the shortest in length.


This was followed by a greater hip flexion velocity in the dominant limbs for the uninjured group at the end of the leg cocking phase (Figure 1). It is in this phase when the knee continues to flex while the hip begins to flex. Theoretically, the muscle fibres in the hamstring muscle group begin to lengthen, and this lengthening is slower in the previously injured group in the non-dominant limb (partial $\eta^2 = 0.06$, $p = 0.03$). It is in this phase when the knee reaches a peak velocity, and due to this inhibition of rapid lengthening in the leg cocking phase, this difference across limb dominance is seen only in the uninjured group.

Table 2
Peak moments of the hip

		Peak Flexion moment (Nm/kg)	Peak Extension Moment(Nm/kg)
Uninjured group	Non-Dominant	3.62 ± 1.02*	-2.95 ± 1.12*
	Dominant	4.03 ± 0.74*	-3.69 ± 0.92*
Injured group	Non-Dominant	3.80 ± 0.45	-2.80 ± 0.84
	Dominant	4.44 ± 1.13	-3.78 ± 1.20

* Significant difference between the two groups with $p < 0.05$.


Figure 1
Hip flexion velocity (rad/s) at the instance of peak knee flexion



ND: Non-dominant limb kick, D: dominant limb kick. * Significant difference between the two groups with $p < 0.05$.

Looking at female kicks alone at the instance of ball impact, a greater knee flexion angle was observed in the non-dominant limb kicks (Figure 2), which facilitated a better transfer of momentum leading to higher ball velocities (Navandar et al., 2016). However, if one were to group them further based on previous injury, this difference is seen only in the uninjured group. Previously injured female soccer players have a lower knee flexion angle at ball impact in the non-dominant limb, restricting the extension of hamstring muscle fibres (the knee extends and the hip flexes in this phase). However, this does not translate to higher ball velocities in the injured, non-dominant limb kicks for females, and this could be due to the angle of the ankle at ball impact which determines its quality.

Figure 2
Comparison of knee flexion angle (°) at ball impact in female soccer players)



ND: Non-dominant limb kick, D: dominant limb kick. * Significant difference between the two groups with $p < 0.05$.

Finally, in the follow through phase a difference across limb dominance is seen for the uninjured group in the peak hip extension moment (Table 2), which is reached when the hip reaches a maximum flexion angle. This phase corresponds to one where the fibres are lengthened to a peak, and this is affected in the previously injured limbs.

Conclusions: Similar to previous studies, differences were found between the dominant and non-dominant limb kicks. However, these differences were affected by the presence of a previous injury to the hamstring muscle group. Injured dominant and non-dominant limb kicks showed no differences in phases where the hamstring muscles were lengthening. Coaches, rehabilitators and biomechanists must take this into account while dealing with a player who has previously injured his hamstrings, as it appears to affect the non-dominant limb more than the dominant limb.

REFERENCES:

- Cohen, J. (1992). A Power Primer. *Psychological Bulletin*, 112(1), 155-159.
- Ekstrand, J., Häggglund, M., & Waldén, M. (2011). Epidemiology of Muscle Injuries in Professional Football. *American Journal of Sports Medicine*, 39(6), 1226-1232.
- Ekstrand, J., Waldén, M., & Häggglund, M. (2016). Hamstring injuries have increased by 4% annually in men's professional football, since 2001: a 13-year longitudinal analysis of the UEFA Elite Club injury study. *British Journal of Sports Medicine*. doi: 10.1136/bjsports-2015-095359
- Freckleton, G., & Pizzari, T. (2013). Risk factors for hamstring muscle strain injury in sport: a systematic review and meta-analysis. *British Journal of Sports Medicine*, 47, 351-358. doi: 10.1136/bjsports-2011-090664
- Lees, A. (2013). Biomechanics applied to soccer skills. In A. M. Williams (Ed.), *Science and Soccer* (3 ed., pp. 218-223). London: Routledge.
- Lees, A., Asai, T., Bull-Andersen, T., Nunome, H., & Sterzing, T. (2010). The biomechanics of kicking in soccer: A review. *Journal of Sports Sciences*, 28(8), 805-817. doi: 10.1080/02640414.2010.481305
- Lees, A., & Nolan, L. (1998). The biomechanics of soccer: A review. *Journal of Sports Sciences*, 16, 211-234.
- Navandar, A., Garcia, C., Veiga, S., Navarro, E., Torres, G., & Chorro, D. (2015). *Effect of previous hamstring injury and limb dominance on kicking biomechanics in elite female soccer players*. Paper presented at the 33rd International Conference on Biomechanics in Sport, Poitiers.
- Navandar, A., Gulino, M., Antonio, R., & Navarro, E. (2013). Effect of hamstring injuries on kicking in soccer using inverse dynamics. *Biomecánica*, 21, 7-19.
- Navandar, A., Veiga, S., Garcia, C., Rueda, J., Torres, G., Chorro, D., & Navarro, E. (2016). *Gender effects on limb dominance in kicking biomechanics in elite soccer players* Paper presented at the 34th International Conference on the Biomechanics of Sport, Tsukuba.
- Noya, J. (2015). *Análisis de la incidencia lesional en el fútbol profesional español en la temporada 2008-2009*. (Doctor of Philosophy), Universidad Politecnica de Madrid, Madrid.
- Nunome, H., Ikegami, Y., Kozakai, R., Apriantono, T., & Sano, S. (2006). Segmental dynamics of soccer instep kicking with the preferred and non-preferred leg. *Journal of Sports Sciences*, 24(5), 529-541.
- Richardson, J. T. E. (2011). Eta squared and partial eta squared as measures of effect size in educational research. *Educational Research Review*, 6(2), 135-147.
- van den Tillaar, R., & Ulvik, A. (2014). Influence of Instruction on Velocity and Accuracy in Soccer Kicking of Experienced Soccer Players. *Journal of Motor Behavior*, 46(5), 287-291. doi: 10.1080/00222895.2014.898609