POLITÉCNICA

Universidad Politécnica de Madrid

Escuela Técnica Superior de Ingeniería Agronómica,
Alimentaria y de Biosistemas

Departamento de Biotecnología-Biología Vegetal

**Characterisation of Non-symbiotic Rhizobial and
Serratia Populations from Soil**

PhD Thesis

Amalia Soenens Martínez de Murguía

BSc in Biotechnology

MSc in Agricultural and
Forestry Biotechnology

Madrid, 2018

**Programa Oficial de Doctorado en Biotecnología y
Recursos Genéticos de Plantas y Microorganismos
Asociados**

Departamento de Biotecnología-Biología Vegetal
Escuela Técnica Superior de Ingeniería Agronómica,
Alimentaria y de Biosistemas

Universidad Politécnica de Madrid

PhD Thesis:

Characterisation of Non-symbiotic Rhizobial and *Serratia*
Populations from Soil

Author:

Amalia Soenens Martinez de Murguia

BSc in Biotechnology

MSc in Agricultural and
Forestry Biotechnology

Director:

Juan Imperial Ródenas

BSc in Biology

PhD in Biology

A mi madre

*La naturaleza es grande en las cosas grandes, mas es grandísima en las cosas
diminutas.*

Jacques Henri Bernardin de Saint-Pierre

El azar afortunado suele ser casi siempre el premio del esfuerzo perseverante.

Santiago Ramón y Cajal

*And once the storm is over, you won't remember how you made it through, how
you managed to survive. You won't even be sure, whether the storm is really
over. But one thing is certain. When you come out of the storm, you won't be the
same person who walked in. That's what this storm's all about.*

Haruki Murakami

Agradecimientos

En primer lugar, quisiera agradecer de manera especial a mi director Dr. Juan Imperial Ródenas el haberme dado la oportunidad de formar parte de su grupo durante estos últimos años. Gracias Juan por admitirme como estudiante de master en tu laboratorio y poder continuar con mi tesis doctoral bajo tu dirección. Gran parte de mi desarrollo personal y científico durante estos años te lo debo a ti. Has sido el mejor director que podría haber tenido y te estaré eternamente agradecida. Además de ser un brillante científico y pensador eres una gran persona.

To Prof Alison Smith for welcoming me in her lab during my final degree project. My experience in the JIC encourage me to begin with my scientific career.

To Prof George Salmond for allowing me to do my PhD stay in his lab and showing me the exciting world of *Serratia*.

To Dr Rita Monson for your infinite help during my stay in Cambridge, from the beginning to the end. It was definitely not an easy task showing me everything from the start, thank you very much for your patience! And thanks to the rest of the lab members that made my stay in Cambridge so great; Sarah, Megan, Tracy, Andy, Bihe, Alex..

A la Dra. Azucena González por acogerme en el ICA (CSIC) y adentrarme en el mundo de los productos naturales. Tampoco me puedo olvidar de todos los compis del ICA (Fer, Felipe, Lucía, Viri, Paula y Kodjo) que facilitasteis enormemente mi vida en el centro (¡sois unos “cracks”!).

A los miembros del laboratorio (271) pasados y presentes; Bea, Rosabel y Nuria. Gracias Bea por todo el apoyo durante las etapas iniciales en el laboratorio, por dirigirme mi trabajo fin de máster y ¡por enseñarme a hacer los aislamientos de suelo! A Rosabel por ayudarme siempre, mostrar una sonrisa y aclararme tantas dudas en el laboratorio; y a Ferrán, por ser mi único estudiante de verano, fue un placer tenerte y aprendí mucho enseñándote. Al 281, con quien he compartido laboratorio durante todos estos años. Quiero agradecer a Manuel González-Guerrero toda su ayuda en mi inicio en el laboratorio y especialmente, durante mi trabajo fin de master. A todos los miembros presentes y pasados del 281: María, Isidro, Vivi, Manolo, Ángela, Jesús, Rosi, Marta, Patri y Javi. A Marta, por el apoyo, viajes y las cervezas, y a los peques, Patri y Javi, por aguantarnos tantas risas (y ¡mucho ánimo a los dos que esto se acaba

en nada!). A los estudiantes tan simpáticos: Edu, Eric, Esther, Mario y Alba, ¡gracias por darle tanta alegría al labo!

El CBGP es un ecosistema diverso donde muchas personas y grupos contribuyen a que se respire un ambiente agradable, divertido y de apoyo. Por esto no puedo olvidarme de nuestros primos hermanos del 251, donde nunca he recibido un no por respuesta cuando he necesitado ayuda (¡y han sido muchas veces!). Muchas gracias a todos los que ya no están (Ana, David, Álvaro..) y a los que siguen (Marta, Luis..) y, por supuesto, a Pepe Palacios y Tomás Ruiz-Argüeso. También agradecer a nuestros vecinos de enfrente, los “Rubio” y los “silencing”, por vuestra ayuda, por aguantar todas nuestras risas (sobre todo la mía...) y charlas, y hacer del “pasillo” un ambiente siempre agradable, ¡da gusto tener unos vecinos así! Sois el pilar del CBGP. A María de esterilización y lavado, por ser tan simpática, agradable y recibirme siempre con una sonrisa aun cuando te tocaba esperarme para poner la autoclave, ¡es una lujo tenerte en el CBGP!

Las cañas de apoyo y sobre todo de diversión (y salidas, que para qué nos vamos a engañar, dan mucha vida...) durante la tesis han sido fundamentales. Estos momentos no hubieran sido posibles sin esas grandes amistades que han ido surgiendo y que sé que se quedan para siempre. A Moni, por nuestras aventuras (¡que son muchas..!) y viajes juntas (¡parece mentira que hayamos llegado hasta la India juntas!), por hacer de niñera, por todo el apoyo que me has dado en los momentos más difíciles y por un gran año de master que pasamos juntas (y todo lo que vino después, claro). Eres una de las personas más trabajadoras que conozco, ojalá hubiéramos acabado a la vez, pero ¡ya no te queda nada! Y yo estaré ahí (aunque sea en la distancia) para apoyarte siempre. Rosi, eres mi otra pieza fundamental del puzzle, compañera de laboratorio y compañera de vida. Sin ti como compañera nada hubiera sido lo mismo (y mucho menos divertido seguro). Los momentos son incontables, has sido y eres mi principal apoyo en el laboratorio, me has aguantado, ayudado y apoyado siempre incondicionalmente, aun cuando no sabías muy bien cómo hacerlo, ¡eres una amiga y compañera 10! Chicas, soy consciente de la suerte que he tenido en coincidir con vosotras, y que continúen nuestras aventuras. A LA NUEVA INCORPORACIÓN, ¡María! Cuanto me alegra haberte conocido, aunque haya sido poco tiempo lo hemos exprimido al máximo, cuantas aventuras hemos escrito en nuestro libro ya... ¡y cuantas nos quedan! Gracias por ser como eres, eres única, ¡nunca cambies por favor! Por siempre dar ese ambiente tan positivo y motivante en el laboratorio y por ayudarme siempre. He aprendido muchas cosas contigo y desde luego esta última

etapa hubiera sido mucho más agridulce sin ti, además estos agradecimientos no tendrían ningún sentido (literalmente). Te mereces lo mejor y estoy segura de que ese momento va a llegar muy pronto. A las mamis: a Gema por las todas las charlas, consejos, ayuda y momentos ¡te mereces lo mejor y estoy segura de que por fin llega lo bueno! Y a Ángela por cuidarnos tanto y por todas esas risas que pasamos dentro y fuera del laboratorio. Tampoco me puedo olvidar de todos los compis de CBeerGP, en especial de Rocío, Ana, y Laura.

Sin duda, he sido y soy muy afortunada por poder contar con el apoyo de MIS AMIGAS que no están conmigo diariamente en el labo pero que seguís siendo de lo más importante, ahora y siempre. Teneros es un tesoro. Aunque en esta última etapa haya estado más distante tengo que agradeceros que siempre estáis ahí. GRACIAS a mis amigas de siempre, “las golfas” y las de Sanse, por vuestro apoyo incondicional y vuestra comprensión, muchas gracias de verdad, solo puedo decir ¡qué gran suerte teneros! También a Crissa y Bea, ¡por todos nuestros fines de semanas gastronómicos llenos de charlas hasta horas infinitas! A Blanqui por todos nuestros años de carrera juntas que nunca olvidaré ¡mi mejor compi de carrera sin duda! A Andrea e Iris, porque, aunque nos vemos menos de lo que me gustaría, nos queremos mucho. ¡Parece que este año es nuestro año! To Loren, for so many years together and still counting, thank you for always being there, even in the distance and giving me so many great advices. I think long-distance friendship defines US, but far doesn't mean less.

A mi familia “las Murgui”; a Laura y Borja, a Natalia, y a los Amunarriz. A mi tía Leti, ¡por todas esas charlas científicas tan interesantes y apoyo siempre! A Sveta por cuidarme tanto y, por supuesto, a mi abuela, LA SÚPER ABUELA, por tanto cariño y risas siempre. Aunque me sigas preguntando por mis estudios y tengas tres hijas doctoras, ¡eres la mejor abuela del mundo!

Juan, eres, sin lugar a dudas, el mejor compañero de vida que puedo tener (soy una afortunada). Conocerle ha sido una de las mejores cosas que me ha pasado (Gracias Crissa, ¡te estaré eternamente agradecida!). Aunque son muchas las cosas que te agradecería, me quedo con todas las risas, los buenos momentos y tu cariño. Has sido clave en esta última etapa, sé que ha sido un tiempo difícil, pero te has superado con creces. ¡Qué ganas de empezar nuestra siguiente etapa llena de aventuras! aunque a veces me da vértigo, sé que no hay nadie mejor con quien comenzarla. A efectos prácticos, quiero darte las gracias por ¡toda tu ayuda con la edición!

Por último, y como no podría ser de otra manera, quiero dar gracias infinitas a mi Madre. Eres mi ejemplo a seguir, y lo que soy te lo debo a ti. Por todo tu sacrificio, generosidad, consejos, valores, apoyo incondicional y comprensión, GRACIAS. Soy una gran afortunada por tenerte y poder disfrutar tanto de nuestro tiempo juntas. También por guiarme siempre por el buen camino en mis momentos más flojos, y por supuesto, poner las cosas en su lugar. ¡No solo tengo la mejor abuela del mundo sino también la mejor madre! Además, he tenido la gran suerte de tener una madre doctora y no podría dejar de agradecerte todo lo que me has enseñado, todas las charlas científicas y toda tu inmensa ayuda con el manuscrito y las referencias, ¡eres una máquina de las correcciones! Desde luego, nada de esto hubiera sido posible sin ti.

TABLE OF CONTENTS

List of Tables.....	V
List of Figures.....	VII
Abbreviations.....	XI
Abstract.....	XV
Resumen.....	XVII
1 INTRODUCTION	1
1.1 Soil microbiota.....	3
1.2 Rhizobia and biological nitrogen fixation (BNF).....	4
1.2.1 <i>Neorhizobium</i>	10
1.2.2 <i>Ensifer</i>	11
1.2.3 <i>Bradyrhizobium</i>	12
1.3 <i>Serratia</i>	13
1.4 Plant growth promoting rhizobacteria (PGPR)	18
2 OBJECTIVES	25
3 MATERIALS AND METHODS	29
3.1 Sampling sites and soil characteristics	31
3.2 Bacterial, fungal and phage strains and culture conditions	33
3.3 Culture media, reagents and solutions	38
3.4 Plant material and techniques.....	42
3.4.1 Seed sterilisation and germination	42
3.4.2 Trap plant assay	43
3.4.3 Most probable number (MPN)	43
3.4.4 Isolation of Rhizobia from nodules	43
3.4.5 Inoculation assay to test for nodulation capacity	43
3.5 Bacterial isolation and techniques	44
3.5.1 Total aerobic counts in soil	44
3.5.2 Phenotypic characterisation	44
3.5.2.1 Carbon and Nitrogen sources	44
3.5.2.2 Fatty acids identification.....	44
3.5.2.3 MALDI-TOF analysis	44
3.5.3 Isolation of <i>Neorhizobium</i> from soil	45
3.5.4 Isolation of <i>Ensifer</i> from soil	45
3.5.5 Isolation of <i>Bradyrhizobium</i> from soil.....	46
3.5.6 Isolation of <i>Serratia</i> from soil	46
3.5.7 Phenotypic <i>Serratia</i> assays	48
3.5.7.1 Swarming and swimming.....	48
3.5.7.2 Protease activity and plant cell wall-degrading enzymes: cellulase and pectate lyase.....	48
3.5.7.3 Quorum sensing.....	49

3.5.7.4 Siderophore activity	49
3.5.7.5 Antibiotic production activity	50
3.6 Bacteriophage techniques and assays with <i>Serratia</i>	50
3.6.1 Bacteriophage stock and titrate preparation	50
3.6.2 Phage sensitivity assay	51
3.7 Fungal techniques and assays with <i>Serratia</i>	51
3.7.1 Yeast growth inhibition	51
3.7.2 Fungal activity against <i>Verticillium dahliae</i>	52
3.8 Recombinant DNA techniques	52
3.8.1 DNA purification and visualisation	52
3.8.1.1 Genomic DNA extractions	52
3.8.1.2 DNA quantification	52
3.8.1.3 Agarose gel electrophoresis	52
3.8.1.4 DNA clean-up after enzymatic reactions	53
3.8.2 DNA manipulations	53
3.8.2.1 Polymerase chain reaction (PCR)	53
3.8.2.2 Oligonucleotide primers	54
3.8.3 DNA sequencing methods	55
3.8.3.1 Sanger sequencing	55
3.8.3.2 Whole genome sequencing	55
3.9 Bioinformatics	55
3.9.1 Phylogenetic analysis	55
3.9.2 Genome assembly and annotation	56
3.9.3 Genomic comparisons	56
3.10 Data analyses	56
4 RESULTS	57
4.1 <i>Neorhizobium</i>	59
4.1.1 Direct isolation of <i>Neorhizobium</i> from soil	59
4.1.2 Genotypic and genomic characterisation of <i>Neorhizobium</i> isolates	61
4.1.3 Description of a new species, <i>N. tomejilense</i> sp. nov.	68
4.1.4 Phylogenetic analysis	69
4.1.5 Genome properties and ANI values	71
4.1.6 Phenotypic analysis	73
4.2 <i>Ensifer</i>	79
4.2.1 Development of an <i>Ensifer</i> genetic marker	79
4.2.2 <i>Ensifer</i> direct isolation	79
4.2.3 Characterisation of <i>Ensifer</i> populations in Tomejil and Motril soils	82
4.2.4 Genome comparison between a soil non-symbiotic isolate and a nodule symbiotic isolate from the same soil	86
4.3 <i>Bradyrhizobium</i>	93
4.3.1 <i>Bradyrhizobium</i> semi-selective medium BSJM optimisation and marker development	93
4.3.2 <i>Bradyrhizobium</i> direct isolation method strategy	96
4.3.3 Characterisation of <i>Bradyrhizobium</i> populations	99
4.4 <i>Serratia</i>	109
4.4.1 <i>Serratia</i> isolation using the BSJM semi-selective medium	109
4.4.2 Characterisation of <i>Serratia</i> populations	110

4.4.3	Characterisation of secondary metabolism and phage sensitivity phenotypes in the <i>Serratia</i> soil isolates	115
4.4.4	Correlation between <i>Serratia</i> phylogeny and the phenotypes obtained	121
5	DISCUSSION	127
5.1	Non-symbiotic rhizobial populations in soil	130
5.2	<i>Serratia</i> populations in soil.....	135
5.3	Rhizobia and <i>Serratia</i>, two different strategies for life in the soils and potential PGPRs.....	138
6	CONCLUSIONS	141
7	BIBLIOGRAPHY	145
8	SUPPLEMENTAL TABLES AND FIGURES	177

List of Tables

Table 1. Non-symbiotic <i>Ensifer</i> and <i>Bradyrhizobium</i> isolates described in the literature.	9
Table 2. PGPR activities described in <i>Neorhizobium</i> , <i>Ensifer</i> , <i>Bradyrhizobium</i> and <i>Serratia</i> .	21
Table 3. Physicochemical properties of soils from different locations.	32
Table 4. Bacterial, fungal and phage strains used in this study.	33
Table 5. Media used in this study.	38
Table 6. Antibiotics, antifungals and supplements used in this study.	40
Table 7. Solutions used in this study.	41
Table 8. PCR components.	53
Table 9. PCR standard conditions.	54
Table 10. Primers used in this study.	54
Table 11. Summary of results obtained for <i>Neorhizobium</i> sp. isolation.	59
Table 12. Genomic features of <i>Neorhizobium</i> genomes sequenced in this work.	62
Table 13. Presence of <i>repABC</i> regions in the genomes of Tomejil strains.	64
Table 14. Pairwise average nucleotide identities (ANIb) between representative <i>Neorhizobium</i> sp. Tomejil isolates and <i>Neorhizobium</i> sp. type strains.	67
Table 15. Genomic features of <i>Neorhizobium</i> sp. T20_22, T17_20 and T11_12 genomes sequenced in this work.	69
Table 16. Pairwise average nucleotide identities (ANIb) among T20_22, T7_12, T25_27 and type strains from the different <i>Neorhizobium</i> species.	72
Table 17. Differential characteristics among <i>N. tomejilense</i> strains T11_12, T17_20, T20_22 and type strains from <i>Neorhizobium</i> species.	74
Table 18. Fatty acid patterns of <i>N. tomejilense</i> strains T11_12, T17_20, T20_22 and type strains from <i>Neorhizobium</i> species.	76
Table 19. Summary of results obtained with the <i>asmA</i> <i>Ensifer</i> genus marker.	80
Table 20. <i>asmA</i> marker gene (456 bp) conservation in <i>Ensifer</i> sp. strains.	80
Table 21. Total viable aerobic bacteria count (CFU/g), <i>Ensifer</i> CFU/g and nodule MPN/g counts in Tomejil and Motril soils.	82
Table 22. <i>Ensifer</i> isolates obtained from Tomejil and Motril soils.	82
Table 23. Genomic features of <i>Ensifer</i> sp. MMN_5 and <i>Ensifer</i> sp. M4_45 genomes sequenced in this work.	86
Table 24. Pairwise average nucleotide identities (ANIb) between <i>Ensifer</i> sp. M4_45, <i>Ensifer</i> sp. MMN_5 and <i>Ensifer meliloti</i> genomes.	87

Table 25. Size and G+C (%) comparison of the chromosome and pSyms of the reference genome <i>E. meliloti</i> 1021, <i>Ensifer</i> sp. MMN_5 and <i>Ensifer</i> sp. M4_45.	91
Table 26. Pairwise average nucleotide identities (ANIb) calculated for individual replicons from <i>Ensifer</i> sp. MMN_5, <i>Ensifer</i> sp. M4_45, and the reference genome <i>E. meliloti</i> 1021.	91
Table 27. <i>Bradyrhizobium</i> strains tested for growth on semi-selective medium BSJM.	93
Table 28. <i>Bradyrhizobium</i> genomes used in the pipeline for identification of possible genus-specific markers.	95
Table 29. Summary of direct <i>Bradyrhizobium</i> isolation results.	97
Table 30. Sequence conservation of the <i>Bradyrhizobium</i> genus-specific <i>thrC</i> marker (215 bp) in <i>Bradyrhizobium</i> sp. isolates.	98
Table 31. Total viable aerobic bacteria count (CFU/g), <i>Bradyrhizobium</i> CFU/g and nodule MPN/g counts in Burguillos soil.	100
Table 32. Symbiotic phenotypes Burguillos <i>Bradyrhizobium</i> isolates.	100
Table 33. Genomic features of <i>Bradyrhizobium</i> sp. B2_9, B3_1 and B3_2 genomes sequenced in this work.	107
Table 34. Pairwise average nucleotide identities (ANIb) between sequenced <i>Bradyrhizobium</i> sp. isolates and <i>Bradyrhizobium</i> genomes.	108
Table 35. <i>Serratia</i> strains isolated from Tomejil, Motril and Burguillos soils.	110
Table 36. Estimation of <i>Serratia</i> soil populations.	110
Table 37. Genomic features of <i>Serratia</i> sp. B1_6, B1_23, B1_42, T1_10, T1_12, T4_5, T3_17, M3_1 genomes sequenced in this work.	113
Table 38. Pairwise average nucleotide identities (ANIb) between sequenced representative <i>Serratia</i> sp. isolates and <i>Serratia</i> genomes.	114
Table 39. Strain composition of <i>Serratia</i> groups identified by the PCA analysis.	116
Table 40. Secondary metabolism and phage sensitivity phenotypes of representative <i>serratia</i> strains from the groups of the PCA analyses.	119
Table 41. Presence of antibiotic, antifungal and mVOCs gene clusters in <i>Serratia</i> genomes and comparison with phenotypic screening results.	121
 Supplemental Table S1. <i>Neorhizobium</i> , <i>Ensifer</i> , <i>Bradyrhizobium</i> and <i>Serratia</i> Genbank and JGI (bold) accession numbers.	 179
Supplemental Table S2. <i>Neorhizobium</i> , <i>Ensifer</i> , <i>Bradyrhizobium</i> and <i>Serratia</i> genotypes.	189

List of Figures

Figure 1. Rhizobia (green) and <i>Serratia</i> (red) representation as possible PGPRs.	23
Figure 2. Site map of Tomejil, Motril and Burguillos soil sampling.	32
Figure 3. <i>Neorhizobium</i> , <i>Ensifer</i> , <i>Bradyrhizobium</i> and <i>Serratia</i> direct isolation methods.	47
Figure 4. Phylogenetic tree of representative Tomejil soil isolates based on PCR amplified near-complete 16S rDNA (1,234 bp) sequences.	60
Figure 5. Phylogenetic tree of representative Tomejil soil isolates based on PCR amplified partial <i>rpoB</i> (356 bp) sequences.	61
Figure 6. PCR amplification of a genomic DNA band from Tomejil <i>Neorhizobium</i> sp. strains with <i>fnrN</i> primers.	63
Figure 7. Phylogenetic tree based on <i>repABC</i> sequences from Tomejil sequences and <i>Neorhizobium</i> megaplastids.	65
Figure 8. Phylogenetic tree of sequenced Tomejil strains and of <i>Neorhizobium</i> type strains based on a concatenation of complete <i>atpD</i> , <i>glnII</i> , <i>recA</i> , <i>rpoB</i> , and <i>thrC</i> genes (8,949 bp).	66
Figure 9. Dendrogram representation of a Euclidean distance matrix derived from pairwise average nucleotide identities (ANi) between Tomejil and <i>Neorhizobium</i> type strain genomes.	68
Figure 10. Maximum likelihood phylogenetic tree based on 16S rDNA sequences (1,476 bp) of <i>Neorhizobium tomejilense</i> sp. nov., <i>Neorhizobium</i> type strains and representative <i>Rhizobium</i> type strains.	70
Figure 11. Maximum likelihood phylogenetic tree based on full <i>atpD</i> , <i>glnII</i> , <i>recA</i> , <i>rpoB</i> and <i>thrC</i> concatenated sequences (8,386 nt) of <i>Neorhizobium tomejilense</i> sp. nov., and sequences from <i>Neorhizobium</i> type strains and type strains from representative <i>Rhizobium</i> species.	71
Figure 12. M13-RAPD-PCR band pattern fingerprints of <i>N. tomejilense</i> and type strains of <i>Neorhizobium</i> species.	73
Figure 13. Dendrogram constructed by applying average linkage clustering and Pearson's distance correlation from a similarity matrix of identical mass peaks, computed from whole cell mass spectra of all <i>Neorhizobium</i> strains analysed.	75
Figure 14. PCR amplification of the <i>Ensifer</i> marker gene <i>asmA</i> (522 pb) from <i>Ensifer</i> laboratory strain collection and from other laboratory strains.	79
Figure 15. Phylogenetic tree of representative Tomejil and Motril soil and nodule isolates based on PCR amplified near complete 16S rDNA sequences (1,277 bp) and type strains from <i>Ensifer</i> species.	84

Figure 16. Phylogenetic tree of representative Tomejil and Motril soil and nodule isolates based on PCR amplified partial concatenated <i>recA</i> (372 bp) and <i>glnII</i> (471 bp) sequences and type strains from <i>Ensifer</i> species.	85
Figure 17. Progressive Mauve genome alignments after contig reordering with the reference genome <i>Ensifer meliloti</i> 1021.	89
Figure 18. Progressive Mauve genome alignment between <i>Ensifer</i> sp. M4_45 (Motril non-symbiotic isolate) and <i>Ensifer</i> sp. MMN_5 (Motril symbiotic isolate).	90
Figure 19. PCR amplification of the <i>Bradyrhizobium</i> marker <i>thrC</i> (296 bp).	96
Figure 20. Phylogenetic tree of representative Burguillos soil and nodule isolates based on PCR amplified near complete 16S rDNA sequences (1,216 bp) and type strains from <i>Bradyrhizobium</i> species.	102
Figure 21. Phylogenetic tree of representative Burguillos soil and nodule isolates based on PCR amplified partial concatenated <i>recA</i> (375 bp) and <i>glnII</i> (470 bp) sequences and type strains from <i>Bradyrhizobium</i> species.	103
Figure 22. Phylogenetic tree of representative Burguillos nodule isolates based on PCR amplified partial <i>nodC</i> (441 bp) sequences and type strains from <i>Bradyrhizobium</i> species.	105
Figure 23. Phylogenetic tree of representative Burguillos soil and nodule isolates based on PCR amplified partial <i>nifH</i> (147 bp) sequences and type strains from <i>Bradyrhizobium</i> species.	106
Figure 24. Phylogenetic tree of representative Tomejil, Motril and Burguillos soil isolates based on PCR amplified near complete 16S rDNA sequences (1,296 bp) and type strains from <i>Serratia</i> species.	112
Figure 25. Principal component analysis of different phenotypes (a) and of <i>Serratia</i> strains against phenotypes (b).	116
Figure 26. Representative secondary metabolism plate assays and phage sensitivity assays in <i>Serratia</i> .	120
Figure 27. Correlation between <i>Serratia</i> 16S rDNA phylogenetic distribution (left) and PCA analysis of <i>Serratia</i> strains against phenotypes (right) for Tomejil soil strains, boxed in yellow.	123
Figure 28. Correlation between <i>Serratia</i> 16S rDNA phylogenetic distribution (left) and PCA analysis of <i>Serratia</i> strains against phenotypes (right) of Motril soil strains, boxed in red.	124
Figure 29. Correlation between <i>Serratia</i> 16S rDNA phylogenetic distribution (left) and PCA analysis of <i>Serratia</i> strains against phenotypes (right) of Burguillos soil strains boxed in blue.	125

Supplemental Figure S1. Multiple alignment of <i>asmA</i> marker gene (465 pb) in <i>Ensifer</i> sp.	196
Supplemental Figure S2. Multiple alignment of <i>thrC</i> marker gene (215 bp) in <i>Bradyrhizobium</i> sp.	199

Abbreviations

aa	amino acid(s)
ABA	Absciscic Acid
AHL	Acyl-homoserine lactone
AMF	Arbuscular mycorrhiza
ANI	Average Nucleotide Identity
ANIm	ANI-MUMmer
ANlb	ANI-Blast
Atm	atmosphere
atpD	ATP synthase subunit beta
<i>asmA</i>	Outer membrane assembly factor
a.s.l.	above sea level
B	Burguillos soil isolates
BG	Brilliant Green
BHL	N-butanoyl-homoserine lactone
BLAST	Basic Local Alignment Search Tool
BLN	Burguillos soil <i>Lupinus albus</i> Nodule isolate(s)
BNF	Biological Nitrogen Fixation
bp	Base pair(s)
BSJM	<i>Bradyrhizobium japonicum</i> semi-selective media
BSN	Burguillos soil <i>Macroptilium atropurpureum</i> isolate(s)
bv.	Biovar
ca.	<i>circa</i> (approximately)
CAS-HDTMA	chrome azurol S-hexadecyltrimethylammonium bromide
Cd	Cadmium
CFU	Colony Forming Unit(s)
Cu	Copper
CTAB	Cetrimonium bromide
°X	Degrees (Celcius)
DMSO	<i>Dimethyl sulfoxide</i>
DNA	Deoxyribonucleic Acid
dNTPs	Deoxyribonucleotide triphosphates
dpi	days post inoculation
16S rDNA	DNA sequencing of the 16S <i>rRNA</i> or <i>rrs</i>
DTPA	diethylenetriaminepentaacetic acid
EOP	Efficiency of Plating
ESS	<i>E. coli</i> β -lactam super sensitive mutant
EtOH	ethanol
FAME	<i>Fatty Acid Methyl Esters</i>

Fe	Iron
<i>fnrN</i>	microaerobiosis transcriptional activator
GA	Gibberellic acid
GLC	Gas-Liquid Chromatography
glnII	glutamine synthetase II
h	hour
ha.	hectare
HEPES	N-(2-hydroxyethyl)-piperazine-N'-(2-ethanesulfonic acid)
IAA	indole-3-acetic acid
ISR	Induced Systemic Resistance
kb	kilobase pair(s)
LB	Luria-Bertani broth
LPS	<i>lipopolysaccharide</i>
M	Motril soil isolates
Mb	megabases
MES	2-(N-morpholino) ethanesulfonic acid
MIDI	<i>Microbial Identification System</i> for identification
ML	<i>Maximum-likelihood</i>
MMN	Motril soil Medicago sativa Nodule isolates
MNBP	<i>Rhizobium leguminosarum</i> semi-selective medium
MP	Maximum parsimony
MPN	Most Probable Number
mVOCs	microbial <i>volatile organic compounds</i>
N	Nitrogen
ND	not determined
nifH ⁺	Nitrogenase iron protein positive
nifH ⁻	Nitrogenase iron protein negative
NJ	Neighbour-joining
nodC ⁺	<i>N-acetylglucosaminyltransferase</i> positive
nodC ⁻	<i>N-acetylglucosaminyltransferase</i> negative
nt	Nucleotide(s)
NT	not relevant
OD600 _{nm}	Optical Density at 600nm
OM	Organic Matter
P	<i>Phosphorus</i>
Pb	Lead
PCA	Principal Coordinates Analysis
PCNB	P- Chloronitrobenzene
PCR	<i>Polymerase chain reaction</i>

PDA	<i>Potato dextrose agar</i>
Pfu	<i>Plaque forming unit(s)</i>
PGPR	plant growth-promoting rhizobacteria
Phage(s)	<i>Bacteriophage(s)</i>
PIPES	piperazine-N,N'-bis(2-ethanesulfonic acid)
ppm	<i>parts per million</i>
pSyma	Symbiotic plasmid a
pSymb	Symbiotic plasmid b
pSyms	Symbiotic plasmid(s)
Pst DC300	<i>Pseudomonas syringae</i> pv. tomato <i>DC3000</i>
PWDE	Plant Wall Degrading Enzymes
φ	Phage / Bacteriophage
QS	quorum sensing
RAPD	Random Amplification of Polymorphic DNA
RAxML	Randomized Axelerated Maximum Likelihood
recA	DNA <i>recombinase A</i>
Rlv3841	<i>Rhizobium leguminosarum</i> bv. viciae strain 3841
RNA	Ribonucleic acid
rpm	revolutions per minute
SDS	sodium dodecyl sulfate
Se	Selenium
Siratro	<i>Macroptilium atropurpureum</i>
SNF	Symbiotic nitrogen fixation
Sp.	species (singular)
Spp.	species (plural)
T	Tomejil soil isolates
Tb	terabases
TBE	<i>Tris-Borate-EDTA</i>
Te	<i>Tellurium</i>
thrC	Threonine synthase
tm	melting temperature
TMN	Tomejil soil <i>Medicago sativa</i> Nodule isolates
TN	Total Nitrogen
TSB	Tryptic soy <i>broth</i>
TY	Tryptone Yeast
T6SS	Type VI secretion system
UPGMA	Unweighted Pair Group Method with Arithmetic Mean
VOCs	<i>Volatile organic compounds</i>
v/v	volume/volume

Abbreviations

w/v	Weight/volume
YE	Yeast Extract
YMB	Yeast Mannitol Broth

Abstract

Soil represents one of the most complex environments and a large bacterial reservoir. In this work, two groups of Gram-negative soil bacteria were studied, Rhizobia and *Serratia*. Rhizobia establish endosymbioses mainly with plants belonging to the Fabaceae (legumes) group. During this symbiosis, legume plants are able to feed from the atmospheric nitrogen fixed to ammonia by the Rhizobia, which makes them independent from any external fixed nitrogen supply. *Serratia* belong to a genus within the Enterobacteriaceae family; characteristic of these bacteria is their secondary metabolism, producing compounds such as prodigiosin or carbapenem that have been extensively studied. Semi-selective enrichment, followed by PCR screening with genus-specific markers developed in this work and species markers, resulted in the successful direct isolation of fast and slow-growing Rhizobia of the genera *Neorhizobium*, *Ensifer* and *Bradyrhizobium*, and of the enterobacterium *Serratia*, from different soils where no legumes had been cultivated. This methodology allowed the isolation and characterisation of Rhizobia, particularly of those lacking either nodulation or fixation genes, which, before these work, had never studied at this level and of *Serratia*, whose populations have also been poorly studied, in both cases due to the lack of appropriate methodology. Abundance and presence of Rhizobia varied among soils, but the rhizobial non-symbiotic populations were, in all cases, larger and more diverse than the symbiotic ones that were obtained using the trap plant method. This led to the identification of novel non-symbiotic Rhizobia populations and to the first non-symbiotic *Neorhizobium* species description. It also allowed the genomic comparison of two *Ensifer* isolates from the same soil, highly-related phylogenetically, one containing symbiotic genes and another one missing them altogether. Whole genome analyses and alignment showed pSyms reorganisation –particularly of pSymA that harbours the nodulation and fixation genes– between these strains. Slow-growing *Bradyrhizobium* were also isolated; our methodology uncovered a very diverse population consisting of symbiotic, non-symbiotic and nitrogen-fixing isolates. Our *Serratia* soil isolates reproduced the diversity of the whole genus. Phenotypic screening of secondary metabolism related compounds, showed production of a full range of potentially interesting compounds, among them some with antibiotic or antifungal activities.

Resumen

El suelo representa unos de los ambientes más complejos a nivel microbiano y un gran reservorio de éstos. En este trabajo se han estudiado dos grupos diferentes de bacterias gram-negativas, los Rizobios y *Serratia*. Los Rizobios establecen endosimbiosis con plantas, fundamentalmente del grupo de las Fabaceae (leguminosas). En esta simbiosis, las leguminosas se alimentan del nitrógeno atmosférico fijado a amonio por los Rizobios. Como consecuencia las leguminosas se independizan de la necesidad de un suministro externo de nitrógeno fijado. *Serratia* es un género de bacterias de la familia Enterobacteriaceae que destaca por su metabolismo secundario, y algunos compuestos derivados de este metabolismo, como la prodigiosina o los carbapenemas, han sido extensamente estudiados. La utilización de medios semi-selectivos de crecimiento, junto con un cribado por PCR utilizando marcadores de género desarrollados en este trabajo o de especie, resultó en el aislamiento de rizobios de crecimiento rápido y lento de los géneros *Neorhizobium*, *Ensifer* y *Bradyrhizobium*, así como de enterobacterias del género *Serratia*, a partir de suelos donde nunca se habían cultivado leguminosas. Esta metodología, permitió el aislamiento y caracterización de Rizobios, en particular, de Rizobios que no contienen ni genes de nodulación ni de fijación de nitrógeno, denominados no simbióticos, que hasta este trabajo no habían sido descritos detalladamente, también de *Serratia*, cuyas poblaciones tampoco habían sido bien estudiadas, en ambos casos debido a la falta de metodología apropiada. La presencia y abundancia de Rizobios varió en los diferentes suelos estudiados. Las poblaciones de Rizobios no simbióticos fueron en todos los casos, mayores y más diversas que las de los simbióticos obtenidos por el método de planta trampa. Como resultado, se identificaron nuevas poblaciones de Rizobios no-simbióticos y se describió la primera especie de *Neorhizobium* no simbiótico. La comparación a nivel genómico de dos aislados de *Ensifer* de un mismo suelo, uno de ellos simbiótico y otro no simbiótico, muy parecidos filogenéticamente, puso de manifiesto la reorganización de los plásmidos simbióticos (pSyms), particularmente del pSymA que contiene los genes de nodulación y fijación. También se aislaron Rizobios del género de crecimiento lento *Bradyrhizobium*; esta población resultó ser muy diversa, conteniendo aislados tanto simbióticos, como no simbióticos o solamente fijadores de nitrógeno. Por otro lado, los aislamientos de *Serratia* obtenidos reproducen la diversidad de la totalidad del género *Serratia*. El cribado fenotípico de compuestos relacionados con el metabolismo secundario mostró un rango de producción muy interesante de compuestos, como por ejemplo antibióticos y antifúngicos.

INTRODUCTION

1.1 Soil microbiota

Soil constitutes one of the richest microbial ecosystems on Earth (Gans, Wolinsky and Dunbar, 2005) and represents a large reservoir of microbial diversity. It comprises a highly dynamic and complex environment and thus soil bacteria play an important role in key biogeochemical cycles such as those of nitrogen (Mooshammer *et al.*, 2014) and carbon (van der Heijden, Bardgett and van Straalen, 2008), and in nutrient acquisition by decomposition (Stursová *et al.*, 2012). It is also a large bacterial reservoir, with estimations, using ribosome content, of around 10^4 bacterial species and 10^9 bacterial cells per gram of soil, whereas estimations using colony-forming units lower this figure down to 10^6 bacterial cells per gram of soil (Torsvik and Øvreås, 2002; Wust *et al.*, 2016). The pioneering work by Fierer *et al.* (2006) and Lauber *et al.* (2009) explored the diversity of bacterial communities in soils across South and North America and across different biomes and showed that these correlated significantly with soil pH. Since then, it was clear that soil ecosystems are still, to a large extent, uncharacterised, and various metagenomics projects have been undertaken to analyse the diversity of the bacterial communities such as the TerraGenome consortium in 2009 (Vogel *et al.*, 2009), the Earth Microbiome Project in 2010 (Gilbert, Jansson and Knight, 2014), the Brazilian and Chinese Microbiome Projects in 2014 and in 2015, and the call by a group of researchers for a unified international microbiome initiative (Alivisatos *et al.*, 2015). At high taxonomic levels, the Acidobacteria, Actinobacteria, Bacteroidetes, Chloroflexi, Firmicutes, and Proteobacteria phyla generally comprise most of soil diversity (Janssen, 2006; Fierer *et al.*, 2009; Fierer *et al.*, 2012; Lauber *et al.*, 2009; Chu *et al.*, 2010; Tecon and Or, 2017). It is customary to distribute bacteria in soil in two main nutritional groups, oligotrophs and copiotrophs. Oligotrophs are bacteria that can live in environments with low levels of nutrients and are characterised by slow growth rates coupled with low rates of metabolism and therefore low population densities, in contrast to copiotrophs (Meyer, 1994; Tate, 1995). Bacterial soil reservoirs can be, and have been, exploited for the characterisation of many bacteria and natural compounds such as new enzymes (Iqbal, Feng and Brady, 2012) and secondary metabolites (Daniel, 2004; Banik and Brady, 2010; Miao and Davies, 2010), for plant growth improvement (Lucy, Reed and Glick, 2004), biological control of plant pathogens (Raaijmakers, Vlami and de Souza, 2002; van Elsas *et al.*, 2008; Thuerig *et al.*, 2009) or bioremediation (Furukawa, 2003; Fagervold, May and Sowers, 2007; Weyens *et al.*, 2009).

Bacterial genomics has revealed that the genomes of soil bacteria are relatively larger and contain more genes than those of bacteria from aquatic or clinical environments (van Elsas *et al.*, 2007; Land *et al.*, 2015). The largest bacterial genome assembled to date belongs to the myxobacterium *Sorangium cellulosum* (14.8 Mb, containing 11,599 genes) (Han *et al.*, 2013). Soil bacteria tend to have higher GC content and larger genome sizes (Wassenaar *et al.*, 2009), this give them a more extensive adaptation potential to cope with the changing soil environment (Konstantinidis and Tiedje, 2004).

Different types of symbiotic interactions can be established between bacteria and plants. These include mutualism, where two organisms act as mutually beneficial; parasitism, when one organism benefits and the other one gets harmed; commensalisms, when one organism benefits from the relationship without the other one getting harmed; and symbiosis *sensu stricto*, when a close, mutually beneficial biological relationship is formed among two or more individuals of different species. With the advent of metagenomics, the full extent of plant-associated microorganisms, whether exo- or endophytically, has been accessible for the first time. The first, pioneering work on plant metagenomics was the one by Lundberg *et al.* (2012) in the model plant *Arabidopsis thaliana*. In this work, it became clear that soil is not just a source of nutrients, but rather a rich and complex environment of bacteria and other microorganisms that interact with the plant and affect its nutrition, growth and even health. One of the best-known examples of natural microbiological interactions within natural environments against pathogens are suppressive soils. These are soils that, due to their microbiota, can limit the activity of plant pathogens so that they determine little or no disease at all, and their existence has drawn attention in view of their potential for sustainable agriculture (Raaijmakers and Weller, 1998; Weller *et al.*, 2002; Sanguin *et al.*, 2009; Kwak and Weller, 2013).

1.2 Rhizobia and biological nitrogen fixation (BNF)

Nitrogen (N) is an essential nutrient for living beings, constituting part of fundamental biomolecules, such as nucleic and amino acids. Nitrogen comprises 1.5-2% of plant dry matter and approximately 16% of total plant protein fraction (Woodmansee *et al.*, 1978; Frink, Waggoner and Ausubel, 1999; Hirsch and Mauchline, 2015). In plants, especially in agriculture, nitrogen availability limits growth and plant development (Gutiérrez, 2012). Despite being one of the most abundant elements on Earth, where it constitutes 78% of the atmosphere, most of the nitrogen exists in the biologically inaccessible form of dinitrogen

(nitrogen gas, N_2) (Canfield, Glazer and Falkowski, 2010). Furthermore, forms of biologically accessible nitrogen, nitrate (NO_3^-) and ammonia (NH_4^+) are found in limited quantities in soil. This limitation has been overcome by the use of nitrogen fertilizers in agricultural soils to achieve greater yields (White and Brown, 2010). Industrial nitrogen fertilizers were developed at the beginning of the 20th century by Fritz Haber and Karl Bosch (Frink, Waggoner and Ausubel, 1999). Through the Haber-Bosch process, atmospheric N_2 is reduced by hydrogen gas to ammonia (NH_3) at high pressures (150-300 atm) and temperatures (400-500 °C). The availability of chemical N fertilizers has been instrumental in the large increases in agricultural productivity attained in the past century. Despite these advantages, N fertilizers have serious limitations. As a consequence of their synthesis being such an energy-intensive process, it is also economically very expensive, which, together with transportation costs, determines high costs for N fertilizers, a major consideration for vast, less developed agricultural areas. In addition, their incorrect, excessive use has caused additional economic and environmental costs. In intensive agricultural crops, such as maize, ca. 50-75% of the nitrogen fertilizer applied is not used by the plant, but leached into soil or lost into the atmosphere in the form of gas through denitrification (Francis, Schepers and Vigil, 1993; Kant, Bi and Rothstein, 2011). Furthermore, the high solubility of N fertilizers, formulated so as to facilitate their uptake and assimilation by the plant, eases their outreach to surface or ground water (Hirel *et al.*, 2011). As a consequence, water can become contaminated by nitrate, making it not potable and potentially harmful (Gruber and Galloway, 2008; Gutiérrez, 2012). Important eutrophication problems can also arise (Robertson and Vitousek, 2009), as well as pollution due to the production of greenhouse gases during their synthesis (Gruber and Galloway, 2008). It has, therefore, become imperative to reduce the use of nitrogen fertilizers and to aim for sustainable agricultural management practices.

Some prokaryotic microorganisms (bacteria and archaea) collectively known as diazotrophs or nitrogen fixers, are able to catalyse the reduction of dinitrogen to ammonia under biologically-compatible conditions. This process, known as Biological Nitrogen Fixation (BNF) is catalysed by a unique nitrogenase enzyme complex. In order to overcome the widespread nitrogen limitation in soil, certain plants, notably legumes, have evolved a symbiosis with nitrogen-fixing bacteria that allows them to become independent of the need for exogenous fertilisation. Due to its importance as an agricultural process, the nitrogen-fixing legume symbiosis has been intensively studied. Only a distinct group of bacteria, collectively known as Rhizobia, is able to establish endosymbioses with plants belonging to the Fabaceae (legumes) group. Although other non-canonical interactions have been

described, such as the one between the non-legume tree *Parasponia* and Rhizobia, of non-legume plants like *Casuarina*, *Datisca* and *Alnus* with the actinomycete soil bacteria of the genus *Frankia*, and *Cycas revoluta* or *Azolla* with Cyanobacteria (Lindblad, Rai and Bergman, 1987; Peters, 1991; Wall, 2000; Pawlowski, 2008; Op den Camp *et al.*, 2011). These symbioses, usually very specific, involve the interaction of signalling molecules from the plant (flavonoids) and the Rhizobia (Nod factors), although symbioses independent of Nod factors have also been described (Giraud *et al.*, 2007; Oldroyd *et al.*, 2011). This early interaction leads to the development of a specialised organ, the root nodule, where nitrogen fixation takes place within a differentiated form of the nitrogen-fixing Rhizobia, the bacteroids (Oldroyd *et al.*, 2011). Nitrogenase-mediated fixation of atmospheric nitrogen (N₂) into ammonia by bacteroids is a highly energy-demanding process. Nitrogenase is also very oxygen-labile and needs low oxygen concentration atmospheres for activity; such an atmosphere is provided by the nodule. Within it, the plant provides bacteroids with carbon and energy sources, and, in turn, bacteroids provide ammonium to the plant (Mulley *et al.*, 2011; Geddes and Oresnik, 2014; Mitsch *et al.*, 2017). Legume plants are therefore able to feed from this nitrogen fixed by the Rhizobia, which makes them independent from any external fixed nitrogen supply, such as fertilizers, making them a primary basis for sustainable agriculture.

Dutch microbiologist and botanist, Martinus Beijerinck in 1888 first isolated a bacterium from plant root nodules, which he named *Bacillus radicicola* (Beijerinck, 1888). A year later, Frank (1889) defined the genus *Rhizobium* (from Latin meaning 'root living') for these bacteria and identified *Rhizobium leguminosarum* and further species belonging to the same group. A group of diverse Gram-negative soil α - and β -proteobacterial diazotrophs, especially within the Rhizobiaceae and Bradyrhizobiaceae families, are collectively known as Rhizobia. Nowadays, common Rhizobia belong to the genera *Rhizobium*, (*Ensifer*) *Sinorhizobium*, *Neorhizobium*, *Mesorhizobium* and *Bradyrhizobium*. Rhizobia are usually minor components of the soil microflora, and estimations have reported low but stable numbers of Rhizobia populations in soil, 10¹-10⁴ per gram of soil (Singleton and Tavares, 1986). It is believed that, as saprophytes, Rhizobia survive within a complex microbial community by adopting an oligotrophic lifestyle, although much of the Rhizobia research has focused on specific aspects of the rhizobial-legume symbioses without considering Rhizobia as members of complex soil communities. The genetic determinants for a successful symbiosis with a host legume are but a small fraction of the genetic complement of Rhizobia. These determinants are often present in plasmids or mobile genomic islands that can be transferred, exchanged

or lost (López-Guerrero *et al.*, 2012; Andrews and Andrews, 2017). Furthermore, symbiosis-related genes have previously been shown to be highly variable among rhizobial species and strains (Sugawara *et al.*, 2013) and acquired by via horizontal gene- and plasmid-transfer events. Therefore, it is possible that a large, non-symbiotic subpopulation co-exists with the symbiotic subpopulation, and that this represents an ill-studied reservoir of genetic diversity. In a pioneering study, Sullivan *et al.* (2002) were able to show that *Mesorhizobium loti*, the microsymbiont of *Lotus* spp. harbours its symbiotic determinants within a symbiotic genetic island integrated in its chromosome, and that this island can excise and be transferred to other cells. Furthermore, after inoculating a soil with a strain containing a marked symbiotic island, they were able to recover it in different, novel genomic backgrounds, thus proving that the island undergoes cell-to-cell transfer in the soil, and that non-symbiotic *M. loti* strains are present in the soil that can receive the marked symbiotic island and thus acquire the ability to nodulate *Lotus*. It is clear that non-symbiotic Rhizobia persist in soils and can freely acquire and transfer symbiotic genes in the absence of legumes (Sullivan *et al.*, 1995) and therefore, the ability to establish a root-nodule symbiosis with legumes is therefore an adaptation of these bacteria to life in the soil.

Common rhizobial markers used to assess symbiotic capacity are the genes *nodC* and *nifH*. In the case of *nodC*, this is a common *nod* (nodulation) gene essential for nodulation in all rhizobial species investigated so far. This gene encodes an *N*-acetylglucosaminyltransferase which is involved in the first step of Nod factor assembly, and it is also a determinant of host range (Schultze and Kondorosi, 1998; Perret, Staehelin and Broughton, 2000). In the case of *nifH*, this gene encodes the nitrogenase reductase component of nitrogenase, and has become the preferred marker gene for studying the phylogeny, diversity, and abundance of nitrogen-fixing microorganisms. Thus, *nifH* is the most widely sequenced marker gene used to identify nitrogen-fixing bacteria and archaea (Raymond *et al.*, 2004; Ando *et al.*, 2005). On the other hand, the ability of legumes to select compatible Rhizobia from soil, acting as “plant traps” has traditionally allowed an adequate estimation and assessment of the diversity of symbiotically competent Rhizobia. This is very convenient, especially because rhizobial soil populations are low, as commented above. However, the use of legume trap plants allows only a very limited glimpse at rhizobial populations in soil, for two reasons. Firstly, legume-rhizobial symbioses are usually very specific, and a specific legume can only be nodulated by a specific type of Rhizobium, and secondly because Rhizobia lacking symbiotic genes cannot be assessed by this assay. The limitations of this conventional methodology and the restricted knowledge of the unexplored rhizobial communities are

widely known; however only a few attempts have been made to overcome this problem, and even in those few cases, the interest was placed on symbiotically-competent Rhizobia. In the 1970s, the group of Johanna Döbereiner started using semi-solid N-free culture media to isolate diazotrophic bacteria from non-leguminous plants (Döbereiner, 1989). Later on, two media were described to directly isolate *Bradyrhizobium* from soil by Gault and Schwinghamer (1993) and Tong and Sadowsky (1994). In the first case, the semi-selective medium contained antibiotics and zinc. In the second case, the BSJM semi-selective medium was a non-antibiotic and heavy-metal based. Media were however focused on the recovery of symbiotic *Bradyrhizobium* inoculants from soybean-cultivated soils and were mainly tested with non-legume cultivated soils. Bromfield, Wheatcroft and Barran (1994) devised a nutrient poor-medium for the isolation of *Ensifer meliloti* from soil, however they differentiated *E. meliloti* from other soil microorganism with a specific DNA probe derived from *nodH*, a genetic determinant of host range. Louvrier, Laguerre, and Amarger (1995), in ground-breaking work, devised a semi-selective culture medium to enrich *Rhizobium leguminosarum* from soil. They were interested in isolating symbiotic strains that had not been selected by the plant host, in order to test the hypothesis that the different plant host species this bacterium colonizes, select specific genotypes among those present in the soil (Louvrier, Laguerre and Amarger, 1996). Novel techniques were also developed for real Rhizobia estimations in soil: Schmidt, Bakole and Bohlool (1968) devised an fluorescent-antibody approach to study rhizobial populations, however this could only be used for quantification, and Macdonald *et al.* (2011) developed a real-time PCR assay for the detection and quantification of *Rhizobium leguminosarum* from soil. Nowadays, the increased availability of bacterial genome sequences has enabled the design of a pipeline for the development of PhyEco marker genes of bacteria and archaea by Wu, Jospin and Eisen (2013). In this work, they developed a systematic approach, where they took a set complete genome sequences and selected a subset of gene families, referred as markers. These marker genes had to be able to conduct phylogeny-driven ecological studies at species level or for taxonomic assignment of metagenomics sequence reads. Therefore, they assigned them as PhyEco (for phylogenetic and phylogenetic ecology) markers. The PhyEco markers, had to satisfy these criteria: universality across the taxa of interest, evenness, low variation in copy number across taxa, monophyly, to see if the gene was monophyletic for the whole family and unique, to distinguish members of the PhyEco marker family of interest from other families.

Recent 16S rDNA amplicon sequencing and metagenomics projects have reported widespread rhizobial communities in non-legume microbiomes (Lundberg *et al.*, 2012; Shakya *et al.*, 2013; Chaparro, Badri and Vivanco, 2014; Ofek-Lalzar *et al.*, 2014; de Souza *et al.*, 2016; Castañeda and Barbosa, 2017). Although short-fragment 16S rDNA amplification and sequencing has classification limits that often fail to reach genus level, these results indicate that Rhizobia could be universally present in legume and non-legume microbiomes. Table 1 below includes a listing of non-symbiotic Rhizobia isolates of the rhizobial genera studied in this work. Only *Bradyrhizobium* and *Ensifer* isolates are included since no non-symbiotic *Neorhizobium* isolates had been described up until this work.

Table 1. Non-symbiotic *Ensifer* and *Bradyrhizobium* isolates described in the literature.

Strains	Place isolated	Reference
<i>Ensifer adherens</i>	Isolated as a bacterial predator in soil	(Casida, 1982)
<i>Ensifer collicola</i>	Isolated from a soil in South Korea	(Jang <i>et al.</i> , 2017)
<i>Bradyrhizobium betae</i>	Isolates from <i>Beta vulgaris</i> tumours, however it does not form tumours or nodules in any of a number of legumes	(Rivas <i>et al.</i> , 2004)
<i>B. elkanii</i> and <i>B. japonicum</i>	<i>Bradyrhizobium</i> directly isolated from inoculated soybean and uninoculated legume-free field soils in Thailand	(Pongsilp <i>et al.</i> , 2002)
<i>Bradyrhizobium</i> sp. S23321	Free-living ancestral <i>Bradyrhizobium</i> , isolated from paddy field soil	(Gorlach <i>et al.</i> , 1994)
<i>Bradyrhizobium</i> sp. IRBG271	<i>Oryza sativa</i> endophyte	(Tan <i>et al.</i> , 2001)
<i>Bradyrhizobium</i> sp. AG41, AG48, M12, M21	Non-symbiotic isolates from sugarcane root extract	(Rouws <i>et al.</i> , 2014)

Strains	Place isolated	Reference
<i>Bradyrhizobium</i> sp. LTSP 885, LTSP M299	These <i>Bradyrhizobium</i> from North America forest soils dominated the microbial communities	(VanInsberghe <i>et al.</i> , 2015)
<i>Bradyrhizobium</i> sp. BF49, G22	First non-symbiotic <i>Bradyrhizobium</i> from European soils, they contain genes for denitrification and are likely playing an important role in denitrification	(Jones <i>et al.</i> , 2016)

1.2.1 *Neorhizobium*

The genus *Neorhizobium* is a recently described genus formed by three species previously assigned to *Rhizobium*: *N. alkanisoli*, *N. galegae* and *N. huautlense*, all highly related to the *Rhizobium* and *Agrobacterium* genus (Mousavi *et al.*, 2014). Bacteria belonging to *Neorhizobium* genus are dispersed worldwide and have symbiotic capabilities, forming nodules with their plant hosts. *N. galegae* was isolated from root nodules of the cold-climate legume *Galega orientalis*, but it is also able to form nodules with *G. officinalis* (Lindström, 1989). In fact, two different symbiovars (sv), highly specific to each of the two *Galega* host species have been described. *G. orientalis* originates in the Caucasus, it is a disease-resistant plant used as a forage legume in Finland and Estonia (Radeva *et al.*, 2001). *G. officinalis* grows as a wild plant from the Caucasus to Southern Europe (Radeva *et al.*, 2001). The type strain of the genus *Neorhizobium* is *N. galegae* HAMBI 540 and was isolated from *G. orientalis* (Lindström, 1989). On the other hand, *N. huautlense* was isolated from the root nodules of *Sesbania herbacea* in Mexico (Wang *et al.*, 1998) and *N. alkanisoli* was isolated from root nodules of *Caragana intermedia* grown in saline-alkaline soils in the north of China (Li Lu *et al.*, 2009). Most studies with *Neorhizobium* have focused on the molecular and genomic determinants of the symbiovars *N. galegae* sv. *officinalis* and *N. galegae* sv. *orientalis* and, correspondingly, *Neorhizobium* sp. environmental isolates have only vaguely been identified. *N. galegae*, for example, has been found as an endophytic bacterium in nodules of *Lotus arabicus*, *L. creticus* and *L. pusillus* (Rejili *et al.*, 2009) and *Sulla pallida* (North African endemic species) (Beghalem *et al.*, 2017). *Nod* and *nif* genes are harboured on large megaplasms in *N. galegae* (Österman *et al.*, 2014), although, *N. galegae* sv. *officinalis* HAMBI 1140, the most studied representative of the sv. *officinalis*, contains both a megaplasmid and a smaller plasmid (Österman *et al.*, 2015).

1.2.2 *Ensifer*

In 1982 Casida proposed the name *Ensifer adherens* for a bacterial predator of bacteria in soil (Casida, 1982). In 1988 the genus *Sinorhizobium* was created with the species *Sinorhizobium fredii* (Chen and Li, 1988) to later include *Sinorhizobium meliloti* (De Lajudie *et al.*, 1994). However, in 2003, the inclusion of *E. adhaerens* and *Sinorhizobium* species within a single genus led to the *Ensifer* taking preference and to the transfer of *Sinorhizobium* species to *Ensifer* (Willems *et al.*, 2003). *Sinorhizobium* contained the root “Rhizobium” that described a relevant property of the organisms, while *Ensifer* –meaning “sword-bearer”– does not have the same relevance for symbiotic, nitrogen-fixing bacteria. However, as of today, only the use of the genus name *Ensifer* is considered valid, and *E. adherens* represents the type species. Currently, the genus comprises twenty-one species, predominantly isolated from plant nodules. Only two species or isolates have been described to be non-symbiotic (Table 1). Members of the genus *Ensifer* were the first to be sequenced and are among the most studied Rhizobia. *E. meliloti* and its close relative *E. medicae* induce the formation of root nodules on *Medicago* species, including *Medicago sativa* (alfalfa) and the model plant *Medicago truncatula* (Young, 2003). It also includes some species capable of nodulating *Glycine max*: *E. fredii*, *E. glycinis*, *E. shofinae*, *E. xinjiangense* and *E. sojae* (this last one isolated from root nodules of *Glycine max* grown in saline-alkaline soils). *E. fredii* for example, has a very wide host range, nodulating more than 79 plant genera representing all three subfamilies of the family Leguminosae (Pueppke and Broughton, 1999). Other species, such as *E. arboris* and *E. kostiense* were isolated from nodules of leguminous trees (Nick *et al.*, 1999), and *E. mexicanus* was isolated from the woody American legume *Acacia angustissima* growing in tropical forests of Mexico (Lloret *et al.*, 2007). Species from extreme climates have also been described, such as *E. alkalisoli*, isolated from root nodules of *Sesbania cannabina* grown saline-alkaline soils (Li *et al.*, 2016).

Within the *Ensifer* genus, *E. meliloti* (initially *Rhizobium meliloti* and later *Sinorhizobium meliloti*) is a model species to study the legume-Rhizobia symbiosis. *E. meliloti* has been the subject of extensive genetic, biochemical, and metabolic research (Jones *et al.*, 2007). Moreover, the genetic diversity of natural populations of *E. meliloti* has been the subject of extensive analyses (Bailly *et al.*, 2006; Badri *et al.*, 2007; Silva, Kan and Martinez-Romero, 2007; Biondi *et al.*, 2009;). *E. meliloti* genome has a large genome with over 6,000 genes distributed in one chromosome and two megaplasms, called pSymA and pSymB (Galibert *et al.*, 2001). The symbiotic determinants are grouped in symbiotic islands within these large

self-transmissible megaplasms (pSyms) (Lagares, Sanjuán and Pistorio, 2014). Megaplasmid pSymA, which has the largest fraction of symbiosis-related genes in *E. meliloti*, is a more variable replicon than the chromosome or pSymB in this bacterium (Galardini *et al.*, 2011). More recently, genome sequences of *E. medicae* and *E. fredii* have been published (Reeve *et al.*, 2010; Weidner *et al.*, 2012; Tian *et al.*, 2012), and many of their genetic features have been well characterised.

1.2.3 *Bradyrhizobium*

The genus *Bradyrhizobium* (α -proteobacteria) is characterised by slow growth rates and by an alkaline reaction in culture media containing mannitol as the carbon source (Jordan, 1982). *Bradyrhizobium* species are capable of living in soil as free-living bacteria, in symbiosis and/or forming endophytic associations with legumes and non-legumes. *Bradyrhizobium*, *Rhizobium japonicum* at that time, was first discovered due to the symbiotic association with the economically important crop soybean, and most studies have been performed on this species. Currently the genus contains thirty-nine species and seven are soybean symbiont species: *B. daqingense*, *B. diazoefficiens*, *B. elkanii*, *B. huanghuaihaiense*, *B. japonicum*, *B. liaoningense* and *B. ottawaense* (Jordan, 1982; Kuykendall *et al.*, 1992; Xu *et al.*, 1995; Zhang *et al.*, 2012; Delamuta *et al.*, 2013; Wang *et al.*, 2013; Yu *et al.*, 2014). *B. japonicum* USDA 110, reclassified as *B. diazoefficiens* USDA 110 (Delamuta *et al.*, 2013) represents the type strain, and was originally isolated from soybean nodules in Florida, USA (Jordan, 1982). *Bradyrhizobium* species have been characterised as the symbionts of tropical pasture, aquatic, and woody legume species, and are broadly distributed geographically worldwide (van Berkum, Leibold and Eardly, 2006; Lu *et al.*, 2014; Delamuta *et al.*, 2015; Yao *et al.*, 2015). Photosynthetic *Bradyrhizobium* have also been identified. *B. oligotrophicum* is an oligotrophic bacterium originally isolated from rice paddy soils in Japan. It is able to nodulate the aquatic legume plant *Aeschynomene indica* and contains also a photosynthetic gene cluster (Ramírez-Bahena *et al.*, 2013). Other photosynthetic *Bradyrhizobium* are: *Bradyrhizobium* sp. ORS278, *Bradyrhizobium* sp. BTAi1 (Giraud *et al.*, 2007) and another *Bradyrhizobium* isolated from the non-legume *Parasponia andersonii* (Trinick and Hadobas, 1989).

Such a large diversity made it clear that *Bradyrhizobium* has a diverse array of metabolic capacities; the recent genomic sequence availability has made this even more apparent. Besides harbouring the largest genomes among Rhizobia (8-9 Mb) (Kaneko *et al.*, 2011), it has been shown that they perform a wide range of biochemical functions including nitrogen

fixation under free-living conditions, photosynthesis, denitrification and aromatic compound degradation (Evans *et al.*, 1990; Molouba *et al.*, 1999; Bedmar, Robles and Delgado, 2005; Torres *et al.*, 2011; Okubo *et al.*, 2012) and are able to cope with stresses such as heavy metals, extreme pH and antibiotics (Cole and Elkan, 1979; Graham *et al.*, 1994; Tong and Sadowsky, 1994). *B. canariense* for example, was described as an acid-tolerant endosymbiont (Vinuesa *et al.*, 2005a). All of these metabolic capabilities allow the adaptation to the constant changing, oligotrophic soil environment. The *Bradyrhizobium* genome characteristics explain such diverse metabolic capabilities. Two main features are characteristic of *Bradyrhizobium* genomes, the large number of genes and the presence of a single, large chromosome without the need of large plasmids. *B. diazoefficiens* USDA 110 genome is a single circular chromosome of 9 Mb that includes a 681kb symbiosis island with a lower GC content (Kaneko *et al.*, 2002). The need to replicate a single, large chromosome might be responsible for the long generation time's characteristic of these bacteria. Furthermore, keeping a single replicon provides a measure of genomic stability that may be advantageous in complex environments. The symbiotic island contains a high number of transposases, as well as partial duplications of the target tRNA genes for insertion, suggesting that symbiotic islands have a high plasticity, probably due to horizontal transfer, DNA insertions and deletions, as well as homologous recombinations, all resulting in large genome rearrangements (Kaneko *et al.*, 2002). Multiple horizontal gene transfer of symbiosis genes has been largely documented and, as a result, the diversity of symbiotic genes in *Bradyrhizobium* appears to be low (Moulin *et al.*, 2004; Vinuesa *et al.*, 2005b; Menna and Hungria, 2011). Legume nodulation independent of Nod factors has also been reported for some members of *Bradyrhizobium*. This was first demonstrated in some *Aeschynomene* species that are nodulated by photosynthetic *Bradyrhizobia* lacking the canonical *nodABC* genes (Giraud *et al.*, 2007).

1.3 *Serratia*

In 1823, Bartolomeo Bizio, a pharmacist from Padua in Italy, described *Serratia marcescens* (Bizio, 1823) representing the type species of the genus. A remarkable characteristic of this bacterium was the production of a bright red pigment called prodigiosin (Williams, 1973). A miraculous bloody discolouration in polenta (corn mush) and bread drew the attention to Bizio. Bizio identified a microorganism as the cause of this phenomenon that he named *Serratia marcescens*: *Serratia* in honour of an Italian physicist named Serafino Serrati and

marcescens from the Latin word for decaying, because the blood pigment was found to deteriorate quickly.

Bacteria from the genus *Serratia* are Gram-negative bacteria from the Enterobacteriaceae family that are classified, to date, in eighteen described species. It is an extremely diverse group of bacteria capable of occupying many different habitats such as water, soil, plants, vertebrates and humans, that includes biologically and ecologically diverse species, from those that are beneficial to economically important plants, to pathogenic species that are harmful to humans. Some species produce the red-pink, non-diffusible pigment, prodigiosin. Members of the type species *S. marcescens* are prodigiosin producers and opportunistic nosocomial pathogens, and represent the most important human pathogens within the genus (Eisenstein, 1990). Species other than *S. marcescens* have also been implicated in human infections: *S. fonticola* (Bollet *et al.*, 1991); *S. liquefaciens* (Roth *et al.*, 2000; Harnett, Allen and Macmillan, 2001; Dubouix *et al.*, 2005); *S. odorifera* (Cook and Lopez, 1998; Lee, Carey and Perlman, 2006); *S. plymuthica* (Reina, Borrell and Llompарт, 1992; Nouh and Bhandari, 2000); *S. quinivorans* (Bollet *et al.*, 1993); *S. rubidaea* (Ursua *et al.*, 1996). Examples of *Serratia* species isolated from different organisms or habitats are; *S. entomophila* isolated from infected grass grubs (Grimont *et al.*, 1988), *S. symbiotica* isolated from a symbiosis established with *Aphis fabae* (Sabri *et al.*, 2011), *S. myotis* and *S. vespertilionis* from bats with white-nose syndrome (García-Fraile *et al.*, 2015) and *S. aquatilis* (Kämpfer and Glaeser, 2016) isolated from fresh water. *S. plymuthica* and *S. proteamaculans*, are most frequently associated with plants, consequently many isolates come from plant-related atmospheres, for example *S. plymuthica* G3 was isolated as an *Triticum aestivum* endophyte (Liu *et al.*, 2011) and *S. plymuthica* PRI-2C and *S. plymuthica* 3Rp8 and 3Re4-18 were isolated from maize rhizosphere (Garbeva *et al.*, 2006; Adam *et al.*, 2016). *S. proteamaculans* was originally isolated from a leaf-spot disease of the tropical plant *Protea* (Grimont, 1978), later on, *S. proteamaculans* 568 was isolated as a root endophyte from a *Populus trichocarpa* plant (Taghavi *et al.*, 2009) and *S. proteamaculans* S4 from the rhizosphere of wild *Equisetum* sp. (Neupane *et al.*, 2013).

Although only a few *Serratia* have been characterised in detail, a noteworthy characteristic of these bacteria is their secondary metabolism and their ability to produce a wide range of natural bioactive products (Slater *et al.*, 2003; Kai *et al.*, 2007; Kai *et al.*, 2010; Matilla *et al.*, 2012b; Matilla, Leeper and Salmond, 2015; Domik *et al.*, 2016a). Generally, motility, secondary metabolism, and virulence in *Serratia* are under control of quorum sensing (QS)

regulatory systems, and secondary metabolism, especially, has been linked closely with QS. Quorum sensing is based on low molecular mass, diffusible molecules (autoinducers) that act as signalling molecules secreted by single bacteria to increase cell-population densities and regulate expression of certain genes in response to environmental cues. In *Serratia*, the most common QS systems are those that employ *N*-acyl-L-homoserine lactones (AHLs) as the autoinducer molecules, these are normally C4-HSL (BHL) and C6-HSL (OHL) molecules (Thomson *et al.*, 2000). Efforts have been made to develop simple bioassays capable of sensing a wide spectrum of AHL signal molecules; the most commonly used AHL biosensors are the short-chain AHL detector strain *Serratia* sp. ATCC 39006 SP19 (Poulter *et al.*, 2010), the mid-length detector strain *Chromobacterium violaceum* CV026 (McClellan *et al.*, 1997), and the broad-range detector *Agrobacterium tumefaciens* NTL4 (pZLR4), especially capable of detecting longer AHL chains (Shaw *et al.*, 1997; Cha *et al.*, 1998; Farrand, Qin and Oger, 2002; Zhu *et al.*, 2003). In *Serratia*, generally, AHLs are synthesised by members of the LuxI family or homologues of AHL synthases. Quorum sensing has been shown to regulate a remarkably diverse number of phenotypes of biological and ecological significance in *Serratia* such as: prodigiosin, antibiotic and antifungal production, biofilm formation, swimming motility (movement powered by individual flagella rotating in liquid environments) and swarming motility (rapid multicellular bacterial surface movement powered by tufts of rotating flagella) (Kearns, 2010), and other, virulence-related traits, such as plant cell-wall degrading enzymes (PCDWE) and siderophores (Angerer, Klupp and Braun, 1992; Thomson *et al.*, 2000; Slater *et al.*, 2003).

Some examples of secondary metabolism products in *Serratia* are worth noting:

Perhaps one of the best-known characteristics of the genus is the production of prodigiosin (2-methyl-3-pentyl-6-methoxyprodiginine). Prodigiosin is a tripyrrole red-pigment, bioactive molecule from the family of prodiginines. In *Serratia* the production of prodigiosin is restricted to three species: *S. marcescens*, *S. plymuthica* and *S. rubidaea* (Grimont, 1984). It is produced in the late log phase of growth (Slater *et al.*, 2003) and has been shown to be influenced by numerous environmental factors, including phosphate availability, media composition, temperature and pH (Williams, Gott and Qadri, 1971; Williams, 1973; Solé *et al.*, 1994; Slater *et al.*, 2003). Several groups of other bacteria produce prodiginines, such as actinomycetes (*Streptomyces coelicolor* A3(2)), and marine bacteria, such as *Hahella chejuensis* KCTC 2396 and *Pseudoalteromonas denitrificans* (Perry, 1961; Kawauchi *et al.*, 1997; Jeong *et al.*, 2005). This group of compounds has received special attention due to its

numerous bioactivities. Prodigiosin has been shown to have antibacterial, antiprotozoal, antimalarial, anticancer and immunosuppressive activities (Williams and Qadri, 1980; Demain, 1995; Bennett and Bentley, 2000; Pérez-Tomás *et al.*, 2003; Espona-Fiedler *et al.*, 2012; Hsieh *et al.*, 2012; Lu *et al.*, 2012). Furthermore, prior to the development of synthetic colour chemistry, *S. marcescens* prodigiosin was produced commercially, for the dyeing of silk and wool.

Serratia also produce antibiotics of the carbapenem group. Carbapenems constitute a diverse group of β -lactam antibiotics characterised by potent antibacterial and β -lactamase-inhibitory activity. They have a broad spectrum and have important applications as antimicrobial agents, particularly in infections mediated by multidrug-resistance bacteria (Moellering, Eliopoulos and Sentochnik, 1989). *Serratia* make a simple carbapenem, 1-carbapen-2-em-3-carboxylic acid with an operon of nine genes (McGowan *et al.*, 1996). Antifungal products have recently discovered in *Serratia*. Oocydin A, an anti-fungal natural product of the polyketide type –polyketides represent an important class of bioactive natural products with a broad range of biological activities. Oocydin A has been shown to be produced by a large transacyltransferase (AT) polyketide synthase gene cluster (Matilla *et al.*, 2012b; Matilla, Leeper and Salmond, 2015). *S. plymuthica* strains have been shown to have *in vitro* antifungal activity against fungal pathogens (Berg, 2000). Another peculiar compound produced by this genus is the recently described Sodorifen. Sodorifen is an unusual unique volatile hydrocarbon (1,2,4,5,6,7,8-heptamethyl-3-methylenebicyclo[3.2.1]oct-6-ene) (von Reuss *et al.*, 2010). This extraordinary structure is composed of a five-member and six-member ring, where every carbon atom of the skeleton is substituted with either a methyl or a methylene group. Sodorifen was emitted by *Serratia plymuthica* 4Rx13 as a volatile organic compound (VOC) (Kai *et al.*, 2010; Domik *et al.*, 2016a) but is also emitted by other few *S. plymuthica* isolates (Weise *et al.*, 2014). Volatile compounds are characterised by their molecular weights of less than 300 Da, high vapor pressures, low boiling points and low polarities. Microbial volatile organic compounds (VOCs) are usually secondary metabolites produced by bacteria, however, the ecological functions of these microbial VOCs (mVOCs) are largely unknown. It is thought that they can act as signals for communication between various organisms or serve as attractant or defense compounds (Wenke, Kai and Piechulla, 2010; Wenke *et al.*, 2012; Effmert *et al.*, 2012). Recently it has been shown that sodorifen could descend from terpene metabolism and the gene cluster responsible of its synthesis has been identified in *Serratia plymuthica* (Domik *et al.*, 2016a; Domik, Magnus and Piechulla, 2016b), although its biological function

remains elusive. *Serratia plymuthica* PRI-2C for example, produced odoriferous when exposed to VOCs emitted by the fungal pathogen *Fusarium culmorum* (Schmidt *et al.*, 2017).

Bacteriophages or “phages”, literally meaning bacteria eaters, are viruses that exclusively infect bacteria, and that reproduce in concert with their hosts in diverse natural environments (Koskella and Meaden, 2013). Bacteriophages exist in truly astronomical numbers. There is an estimated 10^{31} phage particles on Earth (Wommack and Colwell, 2000; Hendrix, 2003; Chibani-Chennoufi *et al.*, 2004), and infection rates are estimated at 10^{25} infections per second (Hendrix, 2003; Lima-Mendez, Toussaint and Leplae, 2007). This intimate interaction between phages and their hosts results in rapid coevolution, which has been observed under both natural environmental and laboratory conditions (Brockhurst *et al.*, 2007; Gomez and Buckling, 2011). Therefore, they have key roles in the evolution of the ecosystems (Koskella and Brockhurst, 2014). Since their discovery in the early 1900s, phages have been used as therapeutic agents in human infections, as alternative biocontrol agents in agriculture, sewage treatment, and the food industry, and they have also proved to be useful in cancer studies and for the delivery of vaccines. Additionally, transducing phages have been important tools in basic bacterial genetics, functional genomics, and synthetic biology applications (Abedon *et al.*, 2011; Goodridge and Bisha, 2011; Citorik, Mimee and Lu, 2014; Salmond and Fineran, 2015) .

Typical complex phages attach to the host cell surface via their tail fibres. Phages bind to specific receptors on the surface of the host cell such as lipopolysaccharide (LPS), outer membrane proteins on the surface of Gram-negative bacteria, flagellum or pilus (Evans *et al.*, 2010). As contact is made, the tail core moves down and penetrates the host outer membrane. Released lysozyme-like enzymes digest the peptidoglycan layer, allowing the phage tail to then move across the second bacterial membrane. DNA is then released into the host cytoplasm (Moak and Molineux, 2003). Once phage DNA is within the host cytoplasm, phages reproduce via two cycles, the lytic and the lysogenic cycles. In the lytic cycle, phage DNA exists as a separate molecule within the bacterial cell and replicates separately from the host bacterial DNA. Phage will be replicated and structural proteins will be produced. The final step is host cell lysis, releasing the new phage progeny into the extracellular media, from where they can then infect subsequent hosts. Bacterial lysis from lytic phage cycles normally produces clear plaques in bacterial lawns growing on petri dishes. In the lysogenic cycle, phage DNA is incorporated into the bacterial chromosome. Once in the lysogenic state, the phage is referred to as a prophage. The prophage will then

be replicated along with the rest of the bacterial chromosome and therefore will be passed to all daughter cells. Turbid plaques, are usually produced by lysogenic phages since there is a certain probability that the prophage will enter the lytic cycle at each round of replication (Mitarai, Brown and Sneppen, 2016). Factors influencing the lysis-lysogeny decision have been studied mainly with the lambda bacteriophage that infects *Escherichia coli*. It was shown that the decision is probabilistic in nature but influenced by the nutritional state of the infected bacterium as well as by the number of co-infecting phage particles (Herskowitz and Hagen, 1980; Erez *et al.*, 2017).

Among the well-studied *Serratia* phages, the following can be mentioned: ϕ 3M, ϕ 0T8 ϕ MAM1 and ϕ RAY26. ϕ 3M is a generalised transducing phage of *S. marcescens* isolated from the same species (Regue, Fabregat and Vinas, 1991). ϕ 0T8 was isolated from *Serratia* sp. ATCC 39006 and has been shown to be flagellum-dependent and was also able to infect *Pantoea agglomerans* (Evans *et al.*, 2010). ϕ MAM1 is a phage isolated from *S. plymuthica* A153 capable of infecting many environmental isolates of *Serratia* that uses capsular polysaccharides as a receptor (Matilla and Salmond, 2012a).

1.4 Plant growth promoting rhizobacteria (PGPR)

Plant Growth Promoting Rhizobacteria (PGPR) are soil-derived organisms able to colonize plant compartments such as the rhizosphere, rhizoplane or endosphere, and positively influence plant growth under certain stressful environmental and soil conditions, such as drought, osmotic stress, temperature and nutrient deficiency. Rhizobacteria can influence plant growth via different direct or indirect molecular mechanisms. Classical PGPR belong to the genus *Bacillus* and *Pseudomonas* (Beneduzi, Ambrosini and Passaglia, 2012).

As previously introduced, biological nitrogen fixation (BNF) can be carried out by both free-living and symbiotic diazotrophs. Plants can only use fixed forms of nitrogen, and associating with diazotrophs can make them, in many cases, less dependent on exogenous, fixed-nitrogen fertilizers. Besides the aforementioned rhizobia-legume symbiosis, rhizospheric and endophytic diazotrophs have been described to fix nitrogen under field conditions with sugarcane and rice for example (Chakraborty and Sen Gupta, 1959). Brazilian sugarcane varieties have been described to incorporate at least 40 kg fixed nitrogen $\text{Ha}^{-1} \text{ year}^{-1}$ (Urquiaga *et al.*, 2011).

Phosphorus (P) is also a major limiting nutrient necessary for plant development and growth (Marschner, 1995). The amount of P available in soils is very low, ca. 10-100 μM in fertile arable soils (Mengel and Kirkby, 1978), as it is mostly present in insoluble forms. In addition, P uptake is also limited by the soil pH, with high pH limiting P acquisition by plants (Schachtman, Reid and Ayling, 1998). In traditional agriculture, this challenge, as with nitrogen, has been overcome by the use of P fertilizers. Many soil microorganisms, however, have developed strategies to improve P availability / uptake, including solubilizing and mineralizing insoluble soil P. These activities can result in increases in plant growth. Mineralization / solubilisation is achieved by the production of organic acids (such as acetate, succinate, citrate, and gluconate) or phosphatases, liberating orthophosphate from inorganic and organic phosphorus pools. In this case, for example, arbuscular mycorrhizal (AM) symbioses with plants are known to have a key role in P nutrition (Smith *et al.*, 2011).

Iron (Fe) is a crucial element for plant development since it is cofactor in many metabolic pathways, notably in respiration and photosynthesis (Marschner, 1995). Similar to P, Fe is abundant in soil, although availability to plants is low due to the low solubility of Fe^{3+} oxides. Because of the low bioavailability of Fe in the environment, microorganisms and plants have developed specific uptake strategies that usually include the production of siderophores. Siderophores are metal-chelating agents with low molecular masses (200–2,000 Da) that are produced especially under Fe-limiting conditions (Schwyn and Neilands, 1987). The formation of Fe^{3+} –siderophore complexes is also affected by pH because of the competition for the free siderophore ligands between free protons and Fe (Albrecht-Gary and Crumbliss, 1998). Microbial siderophores enhance iron solubility in the plant rhizosphere allowing its transfer to the plant, and also lower Fe availability to plant pathogens, limiting their infection capacity (Bakker, Pieterse and van Loon, 2007).

Phytohormones are effector molecules produced in very low concentrations but able to regulate a variety of cellular processes in plants. They are often divided into five major classes; auxins, cytokinins (CK), gibberellins (GA), abscisic acid (ABA), and ethylene, although more recently other compounds with hormonal activity have been identified, such as strigolactones and brassinosteroids (Santner and Estelle, 2009). Plant phytohormones have many different roles in plant growth and developmental events, such as: cell division, elongation, and differentiation, cellular proliferation maintenance, prevention of senescence, improvement of stress responses and adaptation, seed dormancy formation of floral organs, and lateral shoot growth, fruit-ripening hormone, abscission, and pathogen-defence

signalling (Ecker, 1995; Schmulling, 2002; Wilkinson *et al.* 2012; Kumar, Khurana and Sharma, 2014). Many soil microbes synthesize low amounts of phytohormones that can help improve stress tolerance and plant growth under various stress conditions, including salinity, heat, drought and metal toxicity, as reported in many studies (Costacurta and Vanderleyden, 1995; Dimkpa, Weinand and Asch, 2009; Verma, Ravindran and Kumar, 2016; Egamberdieva *et al.* 2017; Yang *et al.* 2017). The best studied example is the production of the auxin indole-3-acetic acid (IAA) and of ACC deaminase in PGPR. IAA is directly biosynthesised by bacteria (Mazzola and White, 1994; Spaepen, Vanderleyden and Remans, 2007) and directly affects plant development. ACC deaminase produced by bacteria, on the other hand, can decrease the ethylene levels to facilitate plant growth and development, especially following a variety of environmental stresses (Honma and Shimomura, 1978; Glick 2014).

Some living organisms have the capacity to limit the pathogenic effects of pathogens. This suppression of pathogenesis is known as biological control, or biocontrol. Many soil bacteria can produce antibacterial or antifungal compounds able to inhibit pathogenic bacteria or fungi. Thus, biological control promises to be a useful alternative approach in the control of plant pathogens in the sustainable agriculture system. *Pseudomonas chlororaphis* MA 342, for example, is widely commercialised in agriculture as a biological control agent against fungal pathogens (Hökeberg, Gerhardson and Johnsson, 1997; Tombolini *et al.*, 1999).

Inoculation of plants with non-pathogenic bacteria can also induce resistance against a broad spectrum of pathogenic organisms. This induced systemic resistance (ISR) depends mainly on jasmonate (JA) and ethylene signalling. Well-characterized, ISR-inducing bacteria include several *Pseudomonas* (Bakker, Pieterse and van Loon, 2007), *Bacillus* (Kloepper, Ryu and Zhang, 2004) and *Serratia* species (Singh and Jha, 2016).

Finally, direct contact between microorganisms and the plant is not always necessary for growth promotion and biocontrol. Volatile organic compounds (VOCs) are receiving recent attention, as commented previously. The biological functions of microbial VOCs (mVOCs) are largely unknown, however a few reports have characterised their positive effects. mVOCs produced by plant growth promoting rhizobacteria (PGPR) have been shown to have the potential to control plant pathogens, to stimulate plant growth, and to induce systemic disease resistance (Ryu *et al.*, 2003; Lee *et al.*, 2012; Park *et al.*, 2015). *Bacillus subtilis* SYST2 VOCs, for example, are responsible for differential expression of genes involved in

auxin (IAA), gibberellin (GA), cytokinin (CK), expansin, and ethylene biosynthesis or metabolism in tomato roots and leaves (Tahir *et al.*, 2017).

Table 2 summarises examples of *Neorhizobium*, *Ensifer*, *Bradyrhizobium* and *Serratia* isolates with demonstrated PGPR properties and Figure 1 shows a schematic representation of Rhizobia and *Serratia* as possible PGPRs.

Table 2. PGPR activities described in *Neorhizobium*, *Ensifer*, *Bradyrhizobium* and *Serratia*.

Strains	PGPR activity	Symbiotic (if relevant)	Reference
<i>N. huautlense</i> T1-17	Produces IAA, siderophores, and ACC deaminase, and has high Cd, Pb, Cu, and Zn resistance. Immobilizes Cd and Pb in solution and tissue Cd accumulation in rice (<i>Oryza sativa</i> wuyun-23) and hot pepper	ND	(Chen <i>et al.</i> , 2016; Li <i>et al.</i> , 2017)
<i>E. meliloti</i> TAL 380, TAL 1372 and <i>E. fredii</i> USDA 201	High resistance to Te and Se metals	symbiotic	(Kinkle <i>et al.</i> , 1994)
<i>E. meliloti</i> OS6, S26	Exhibited different efficiency concerning plant growth promotion on nitrogen turnover processes in the rhizosphere during the growth of alfalfa	symbiotic	(Babic <i>et al.</i> , 2008)
<i>E. meliloti</i> 1021	Activation of defence mechanisms in different tissues, enhancement of anabolism (e.g. Photosynthesis) to increase biomass and regulated the auxin level or status to promote growth	symbiotic	(Chi <i>et al.</i> , 2010)
<i>E. meliloti</i> CCNWSX0020	Displayed both copper resistance and growth promotion of <i>M. lupulina</i> in Cu-contaminated soil	symbiotic	(Li <i>et al.</i> , 2014)
<i>E. meliloti</i> TSA3, TSA41	Strains associated to spores of the AMF <i>Rhizophagus intraradices</i> produce IAA and siderophores, and solubilize P from mineral phosphate and phytate	non-symbiotic	(Battini <i>et al.</i> , 2016)
<i>Ensifer</i> sp. CNI, RN2, UNI, RNI	IAA production, chitinase activity, phytase activity and alkaline phosphatase activity	symbiotic	(Iyer, Rajput and Rajkumar, 2017)
<i>Ensifer</i> sp.	Sugarcane isolate selected to form a synthetic community that promotes plant growth in maize	non-symbiotic	(Armanhi <i>et al.</i> , 2017)

Strains	PGPR activity	Symbiotic (if relevant)	Reference
<i>B. japonicum</i> 61A152	Production of siderophores such as citrate	symbiotic	(Guerinot, Meidl and Plessner, 1990)
<i>B. japonicum</i> E109, USDA110, and SEMIA5080	They produce the five major phytohormones, IAA, GA, zeatin, ethylene, and ABA in pure cultures	symbiotic	(Boiero <i>et al.</i> , 2007)
<i>Bradyrhizobium</i> sp. NC 92, 5a/70, NC 6, NC 43.3, NC 70.1, 47 A1, IC 6001	Production of catechol-type siderophores	symbiotic	(Nambiar and Sivaramakrishna, 1987)
<i>Bradyrhizobium</i> sp. TAL 310	Te and Se high resistance	symbiotic	(Kinkle <i>et al.</i> , 1994)
<i>Bradyrhizobium</i> sp. YL-6	IAA, siderophore, ACC deaminase production and solubilisation of tricalcium phosphate and cadmium resistance	symbiotic	(Guo and Chi, 2014)
<i>S. marcescens</i> NBRI 1213	Enhanced plant growth promotion and biologic control of foot and root rot of <i>Piper betle</i> caused by the Oomycete <i>Phytophthora nicotianae</i>	NT	(Lavania <i>et al.</i> , 2006)
<i>S. marcescens</i> CDP-13	Enhanced salinity tolerance, induced systemic resistance and suppressed fungal pathogens in wheat	NT	(Singh and Jha, 2016)
<i>S. plymuthica</i> HRO-C48	Suppression of <i>V. dahliae</i> in strawberry, oilseed rape and olive, and pathogen <i>R. solani</i> on lettuce. It also produced IAA. Production mVOCs negatively influence the mycelial growth of the soil-borne phytopathogenic fungus <i>Rhizoctonia solani</i>	NT	(Kurze <i>et al.</i> , 2001; Grosch <i>et al.</i> , 2005; Kai <i>et al.</i> , 2007; Müller <i>et al.</i> , 2009)
<i>S. plymuthica</i> S13	Antagonist toward <i>Didymella bryoniae</i> , the causal agent of black rot in pumpkins under field conditions	NT	(Förnkrantz <i>et al.</i> , 2012; Müller <i>et al.</i> , 2013)
<i>S. plymuthica</i> 4Rx13	Antifungal activities, mVOCs (sodorifen)	NT	(Kai <i>et al.</i> , 2007; Weise <i>et al.</i> , 2014; Domik <i>et al.</i> , 2016a)
<i>S. proteamaculans</i> 1-102 (1-102)	Biocontrol against the fungal pathogen <i>V. dahliae</i>	NT	(Alström, 2001)

Strains	PGPR activity	Symbiotic (if relevant)	Reference
<i>S. proteamaculans</i> 1-102 (1-102)	Promoted soybean–Bradyrhizobia nodulation and growth	NT	(Bai, Souleimanov and Smith, 2002)
<i>S. proteamaculans</i> 568	Plant growth promotion in poplar, <i>Populus trichocarpa</i> DN-34	NT	(Taghavi <i>et al.</i> , 2009)

ND: not determined, NT: not relevant, AMF: Arbuscular mycorrhiza, IAA: indole-3-acetic acid, GA: Gibberellic acid, ABA: Absciscic Acid, Cd: Cadmium, Pb: Lead, Cu: Copper, Se: Selenium, Te: Tellurium, Zn: Zinc, mVOCs: microbial volatile organic compounds.

Figure 1. Rhizobia (green) and *Serratia* (red) representation as possible PGPRs.

OBJECTIVES

Soil constitutes one of the richest microbial ecosystems on Earth and represents a large reservoir of microbial diversity. Rhizobia are common soil bacteria whose outstanding characteristic is their ability to establish endosymbiosis majorly with leguminous plants. Non-symbiotic Rhizobia lacking nodulation or nitrogen fixation genes have vaguely been described, partly, due to the absence of an appropriate methodology. However, it is assumed that these populations remain in soil, although they have never been described comprehensively and their biological role remains undetermined. In contrast, *Serratia*, a genus within the Enterobacteriaceae family, contain human opportunistic pathogen representatives and their study has mainly focus on this pathogenicity and their second metabolism, also characteristic of *Serratia*. Despite of their presence in many different habitats, *Serratia* populations in natural habitats have vaguely been described. These populations represent a great reservoir of novel secondary metabolism compounds.

The work presented aims towards the characterisation of *Neorhizobium*, *Ensifer*, *Bradyrhizobium* and *Serratia* populations in soil. To achieve this goal, the following objectives were proposed:

1. Development and optimisation of direct isolations methods for *Neorhizobium*, *Ensifer*, *Bradyrhizobium* and *Serratia* from soil.
2. Isolation of *Neorhizobium*, *Ensifer*, *Bradyrhizobium* and *Serratia* from different soils and phylogenetic and phenotypic analysis of the isolates obtained.
3. Genome sequencing of representative isolates and further phylogenomic and genomic analyses. Search for relevant genetic determinants related to their soil lifestyle.

MATERIALS AND METHODS

3.1 Sampling sites and soil characteristics

Tomejil, Motril and Burguillos soils were chosen to characterise the possible rhizobial populations. All soils were from the Andalucía region in Southern Spain (Figure 2). Tomejil soil is a rainfed agricultural soil where mainly cereals are grown and no record exists of legumes having been grown on it. Tomejil soil comes from an Agricultural Experimental Station run by the IFAPA (Investigación y Formación Agraria y Pesquera) Las Torres-Tomejil Centre (<http://www.juntadeandalucia.es/agriculturaypesca/ifapa/web/centro/958d230f-22d4-11df-9a5f-c5d9efb4b7b6>) in Seville. A 10 m x 10 m plot (37°24'33.10" N, 5°34'51.91" W; 77 m above sea level (a.s.l.)) was selected and fenced for our research. Winter wheat is usually grown, with occasional cultivation of cardoon. Motril soil is a sugarcane cultivated soil in Motril, Granada (36°44'44.08" N, 3°33'42.23" W; 19 m a.s.l.) and Burguillos soil comes from a meadow (a pristine "dehesa") in Burguillos, Seville (37°35'9.82" N, 5°57'22.83" W; 60 m a.s.l.). A site map of Tomejil, Motril and Burguillos sampling is shown in Figure 2. For soil collection, portions of soil down to 30-50 cm depth were collected with a shovel from several points and mixed. Soil was transported in sealed plastic bags at room temperature or in dry ice. Once in the laboratory, soil samples were maintained at 4°C, -20°C and -80°C. For physicochemical characterisation of soils, samples were externally analysed in Laboratorio de Edafología y Técnicas Analíticas Instrumentales, EUIT Agrícola, Universidad Politécnica de Madrid, UPM, Madrid, except for Fe contents, that were analysed by Departamento de Ciencias y Recursos Agrícolas y Forestales, Universidad de Córdoba. Physicochemical properties of the soils are described in Table 3.

Figure 2. Site map of Tomejil, Motril and Burguillos soil sampling.

Table 3. Physicochemical properties of soils from different locations.

Property	Tomejil soil	Motril soil	Burguillos soil
pH (Ext.1:2.5) ¹	8.1	8.45	6.2
Pretest salinity ¹ ($\mu\text{S}/\text{cm}$) (Ext.1:5)	239	1,830	65
OM (%) ¹	2.48	2.09	4.8
TN (%) ¹	0.192	0.166	0.230
Carbonates (% CaCO_3) ²	34.5	8.5	-
Exchangeable cation (ppm) ¹	Na	80	600
	K	2,279	1,500
	Ca	7,138	4,400
	Mg	247	830

Property		Tomejil soil	Motril soil	Burguillos soil
	P (Olsen)	40.7	154.6	9.2
Fe (mg/kg)	Oxalate	406	755	643
	DTPA	2.34	4.27	39.70
	Clay	17.2	14.4	8.3
Texture (Bouyoucos method) ¹	Silt	38.2	36.6	34.1
	Sand	44.6	49	57.7

OM, organic matter; TN, total nitrogen. ¹Official method according to MAPA 1994. ² Official method according to UNE 103-200-93.

3.2 Bacterial, fungal and phage strains and culture conditions

Bacterial strains used in this work are listed in Table 4. Rhizobial strains were grown in Yeast Mannitol Broth (YMB) at 28°C. Depending on the species in question, Rhizobia were grown from 48h up to 1 week. *Serratia* sp., *Chromobacterium violaceum* and *Dickeya solani* strains were grown in Yeast Mannitol Broth (YMB) and Luria-broth (LB) at 30°C 24h. *Escherichia coli* strains were grown in Luria-broth (LB) at 37°C 24h. The composition of the culture media used is detailed in Table 5. Long-term stocks of bacterial strains were made by mixing 0.750 ml of grown culture with 0.250 ml of 80% glycerol in a 1.5 ml tube or in a 96-well plate, followed by storage in a -80°C freezer. Fungal strains were grown in Potato dextrose agar (PDA) plates incubated at 25°C for 7 days and preserved on PDA plates at 4°C. Phage stocks were kept over chloroform at 4°C.

Table 4. Bacterial, fungal and phage strains used in this study.

Strain	Relevant characteristics	Reference or Source
Bacterial strains		
Rhizobial strains		
<i>Bradyrhizobium</i> sp.		
<i>B. betae</i> LMG 21987	Type strain, Nod ⁺ Fix ⁺	(Rivas <i>et al.</i> , 2004)
<i>B. canariense</i> BTA-1	Type strain, Nod ⁺ Fix ⁺	(Vinuesa <i>et al.</i> , 2005a)
<i>B. cytisi</i> CTAW11	Type strain, Nod ⁺ Fix ⁺	(Chahboune <i>et al.</i> , 2011)
<i>B. diazoefficiens</i> USDA 110	Type strain, Nod ⁺ Fix ⁺	(Delamuta <i>et al.</i> , 2013)

Strain	Relevant characteristics	Reference or Source
Bacterial strains		
<i>B. elkanii</i> USDA 76	Type strain, Nod ⁺ Fix ⁺	(Kuykendall <i>et al.</i> , 1992)
<i>B. icense</i> LMTR 13	Type strain, Nod ⁺ Fix ⁺	(Durán <i>et al.</i> , 2014a)
<i>B. iriomotense</i> LMG 24129	Type strain, Nod ⁺ Fix ⁺	(Islam <i>et al.</i> , 2008)
<i>B. japonicum</i> USDA 6	Type strain, Nod ⁺ Fix ⁺	(Jordan, 1982)
<i>B. jicamae</i> PAC68	Type strain, Nod ⁺ Fix ⁺	(Ramírez-Bahena <i>et al.</i> , 2009)
<i>B. lablabi</i> CEBAU 223086	Type strain, Nod ⁺ Fix ⁺	(Chang <i>et al.</i> , 2011)
<i>B. liaoningense</i> LMG 18230	Type strain, Nod ⁺ Fix ⁺	(Xu <i>et al.</i> , 1995)
<i>B. lupini</i> ISLU 101	<i>L. angustifolius</i> isolate, Nod ⁺ Fix ⁺	IFAPA
<i>B. pachyrhizi</i> PAC 48	Type strain, Nod ⁺ Fix ⁺	(Ramírez-Bahena <i>et al.</i> , 2009)
<i>B. paxllaeri</i> LMTR 21	Type strain, Nod ⁺ Fix ⁺	(Durán <i>et al.</i> , 2014a)
<i>B. retamae</i> LMG 27393	Type strain, Nod ⁺ Fix ⁺	(Guerrouj <i>et al.</i> , 2013)
<i>B. rifense</i> CTAW71	Type strain, Nod ⁺ Fix ⁺	(Chahboune <i>et al.</i> , 2012)
<i>Bradyrhizobium</i> sp. Btai-1	Photosynthetic strain, Nod ⁺ Fix ⁺	(Evans <i>et al.</i> , 1990)
<i>Bradyrhizobium</i> sp. Lmj C	<i>L. mariae-josephae</i> isolate, Spain, Nod ⁺ Fix ⁺	Laboratory collection
<i>Bradyrhizobium</i> sp. ORS 278	Photosynthetic strain, Nod ⁺ Fix ⁺	(Molouba <i>et al.</i> , 1999)
<i>Bradyrhizobium</i> sp. P. 22A	<i>Lupinus mariae-josephae</i> isolate, Mexico, Nod ⁺ Fix ⁺	Laboratory collection
<i>Bradyrhizobium</i> sp. RSA 104	<i>R. sphaerocarpa</i> isolate, Nod ⁺ Fix ⁺	Laboratory collection
<i>B. valentinum</i> LmjM3	Type strain, Nod ⁺ Fix ⁺	(Duran <i>et al.</i> , 2014b)
<i>B. yuanmingense</i> LMG 21827	Type strain, Nod ⁺ Fix ⁺	(Yao <i>et al.</i> , 2002)
BSN_1 through BSN_12	<i>M. atropurpureum</i> nodule isolate, Burguillos soil, Nod ⁺ Fix ⁺	This study
BLN_1 through BLN_11	<i>L. albus</i> nodule isolates, Burguillos soil, Nod ⁺ Fix ⁺	This study
B1_13, B2_2, B3_1, B3_2, B10_118	Soil isolates, Burguillos, Nod ⁻ Fix ⁻	This study
B2_4, B2_9	Soil isolates, Burguillos, Nod ⁻ Fix ⁺	This study
Ensifer sp.		
<i>E. americanum</i> CFNBI 156	Type strain, Nod ⁺ Fix ⁺	(Wang <i>et al.</i> , 2013)

Strain	Relevant characteristics	Reference or Source
Bacterial strains		
<i>E. arboris</i> LMG 14919	Type strain, Nod ⁺ Fix ⁺	(Young, 2003)
<i>E. fredii</i> USDA 191	Type strain, Nod ⁺ Fix ⁺	(Young, 2003)
<i>E. meliloti</i> 1021	Type strain, Nod ⁺ Fix ⁺	(Young, 2003)
<i>E. xinjiangense</i> LMG 17930	Type strain, Nod ⁺ Fix ⁺	(Young, 2003)
TMN_1 through TMN_31	<i>M. sativa</i> nodule isolates, Tomejil soil, Nod ⁺ Fix ⁺	This study
T1_22, T1_27, T1_34, T1_45, T1_71, T1_77, T1_81, T1_100, T1_107, T2_4, T2_6, T2_8, T2_37, T2_50, T2_53, T3_11, T3_13, T3_47	Soil isolates, Tomejil, Nod ⁻ Fix ⁻	This study
MMN_1 through MMN_10	<i>M. sativa</i> nodule isolates, Motril soil, Nod ⁺ Fix ⁺	This study
M4_5, M4_25, M4_29, M4_43, M4_45, M4_46, M4_50, M5_5, M5_14, M5_22, M5_23, M5_38, M5_45, M5_47, M5_50, M5_53, M5_60, M6_1, M6_18, M7_1, M7_2, M7_3, M7_9, M7_26, M7_27, M8_16, M8_18, M8_19	Soil isolates, Motril, Nod ⁻ Fix ⁻	This study
<i>Neorhizobium</i> sp.		
<i>N. alkalisoli</i> DSM 21826	Type strain, Nod ⁺ Fix ⁺ , LB ⁻	(Mousavi <i>et al.</i> , 2014)
<i>N. galegae</i> HAMBI 540	Type strain, Nod ⁺ Fix ⁺ , LB ⁻	(Mousavi <i>et al.</i> , 2014)
<i>N. huautlense</i> DSM 21817	Type strain, Nod ⁺ Fix ⁺ , LB ⁻	(Mousavi <i>et al.</i> , 2014)
T4_1, T4_8, T5_2, T5_26, T5_27, T6_1, T6_21, T6_23, T6_25, T7_1, T7_7, T7_8, T7_9, T7_11, T7_12, T17_20, T8_5, T9_24, T11_12, T13_2, T16_1, T16_2, T16_4, T16_9,	Soil isolates, Tomejil, Nod ⁻ Fix ⁻ , LB ⁻	This study

Strain	Relevant characteristics	Reference or Source
Bacterial strains		
T16_12, T17_4, T17_6, T17_14, T17_15, T17_26, T18_15, T20_10, T20_15, T20_22, T20_25, T21_1, T21_15, T21_19, T22_7, T22_11, T22_47, T23_12, T23_26, T24_19, T24_25, T25_4, T25_5, T25_7, T25_13, T25_19, T25_20, T25_27, T25_28, T25_30, T28_6		
Other rhizobial strains		
<i>Mesorhizobium loti</i>		
MAF3030	Nod ⁺ Fix ⁺	(Kaneko <i>et al.</i> , 2000)
<i>Rhizobium leguminosarum</i>		
bv. viciae 3841	Nod ⁺ Fix ⁺ , LB ⁻	(Johnston and Beringer, 1975)
Non-rhizobial strains		
<i>Agrobacterium tumefaciens</i>		
GM19023 rifr	Plant pathogen	Laboratory collection
<i>Bacillus</i> sp.		
BW25113/pKD46	Soil Firmicute	Laboratory collection
<i>Chromobacterium violaceum</i> CV026	<i>cvil::Tn5</i> , aHSL indicator strain	(McClellan <i>et al.</i> , 1997)
<i>Dickeya solani</i> MK10	Wild type, pathogen of <i>Solanum tuberosum</i>	(Pritchard <i>et al.</i> , 2013)
<i>Escherichia coli</i>		
K-12 MG1655	Wild type	(Bachmann, 1996)
ESS	β -lactam super sensitive indicator strain	(Nishida <i>et al.</i> , 1977)
<i>Pseudomonas syringae</i> pv. tomato DC3000	Plant pathogen	(Morris <i>et al.</i> , 1995)
<i>Serratia</i> sp.		
<i>Serratia</i> sp. ATCC 39006	Wild type, carbapenem+, prodigiosin+	(Parker <i>et al.</i> , 1982)
<i>Serratia</i> sp. ATCC 39006 LacA	<i>lacZ</i> derivative of 39006	(Slater <i>et al.</i> , 2003)
<i>Serratia</i> sp. ATCC 39006 SP19	<i>smal</i> derivative of S 39006 (<i>smal::mini-Tn5 Sm/Sp</i>), aHSL indicator strain	(Poulter <i>et al.</i> , 2010)
<i>S. plymuthica</i> S13	Plant endophyte and PGPR of <i>Cucurbita pepo</i> L. subsp. <i>pepo</i> var. <i>styriaca</i>	(Müller <i>et al.</i> , 2013)

Strain	Relevant characteristics	Reference or Source
Bacterial strains		
T1_1 through T1_18, T2_2, T2_5, T2_6, T2_9, T2_10, T2_12, T3_1, T3_2, T3_4, T3_5, T3_7, T3_9, T3_10 through T3_23, T4_5, T4_8, T4_12, T4_13, T4_15, T4_16, T4_22, T4_27, T4_28, T4_34, T4_35, T4_36, T4_39, T4_41, T5_1, T5_3, T5_7	Soil isolates, Tomejil	This study
M1_1, M2_3(1), M2_3 (2), M3_1, M3_2, M3_16(R)	Soil isolates, Motril, prodigiosin ⁺	This study
M3_16(W)	Soil isolate, Motril	This study
B1_6, B1_10, B1_23, B1_28, B1_38, B1_40, B1_42 through B1_45, B1_59, B1_62, B1_64, B1_71, B1_72, B1_89, B1_94, B1_98, B1_113, B1_114, B1_115, B1_121	Soil isolates, Burguillos	This study
Fungal strains		
<i>Verticillium dahliae</i> 5368	Wild type, plant pathogen	R. Cooper
<i>Saccharomyces cerevisiae</i>	Wild type	GPCS library
Phages		
φ MAM1	Generalized transducing phage for <i>S. plymuthica</i> A153	(Matilla and Salmond, 2014)
φ 3M	Generalized transducing phage for <i>S. marcescens</i>	(Regue, Fabregat and Vinas, 1991)
φ RAY26	Generalized transducing phage <i>E. coli</i> MG1655	GPCS phage library
φ OT8	Generalized transducing phage for <i>Serratia</i> sp. 39006	(Evans <i>et al.</i> , 2010)
<i>M. sativa</i> , <i>Medicago sativa</i> ; <i>L. albus</i> , <i>Lupinus albus</i> ; <i>M. atropopurem</i> , <i>Macroptolium atropopurem</i> , <i>L. angustifolius</i> , <i>Lupinus angustifolius</i> ; prodigiosin ⁺ , prodigiosin production; LB ⁻ , no growth on LB; GPCS, Prof. George Salmond library collection. IFAPA, Instituto de Investigación y Formación Agraria y Pesquera (Centro Las Torres-Tomejil, Junta de Andalucía), Spain.		

3.3 Culture media, reagents and solutions

Media, antibiotics (including other supplements) and solutions used in this work are listed in Table 5, Table 6 and Table 7, respectively. Media, antibiotics (including other supplements) and solutions, were sterilised by either autoclaving at 121°C for 20min or sterile filtration through 0.22 µm filters.

Table 5. Media used in this study.

Medium	Ingredients per litre	Reference
<i>Bradyrhizobium</i> recovery buffer	13.2 g $\text{NH}_4\text{H}_2\text{PO}_4$ 5.2 ml Tween 80 1.05 mg silicon antifoam 10 g gelatine	(Kingsley and Bohlool, 1981)
BSJM	125 mg Na_2HPO_4 250 mg Na_2SO_4 320 mg NH_4Cl 180 mg $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 10 mg CaCl_2 4 mg FeCl_3 1.3 g HEPES 1.1 g MES 1 g Yeast extract 1 g L-arabinose 1 g D-gluconic acid sodium salt 1 mg Brilliant green (BG) 83 mg ZnCl_2 88 mg CoCl_2 105 mg PCNB 300 mg ciclohexymide pH 6.8	(Tong and Sadowsky, 1994)
CAS agar	6 g NaOH 30.24 g PIPES 15 g agar pH 7	(Schwyn and Neilands, 1987)
CMC- sucrose	16 g agar 0.5 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.5 g KH_2PO_4 2 g K_2HPO_4 2.5 g asparagine	(Pemberton <i>et al.</i> , 2005)

	10 g sucrose	
	500 ml 1% (w/v) carboxymethyl cellulose	
	10 g Tryptone	
Luria-broth (LB)	5 g Yeast extract	(Bertani, 1951)
	5 g NaCl	
Luria-broth agar (LB agar)	As Luria-broth, additional 15 g agar	(Bertani, 1951)
Luria-broth top agar (top LB agar)	As Luria-broth, additional 3.5 g or 7.5 g agar	(Bertani, 1951)
	1 g mannitol	
	178.5 mg Na ₂ PO ₄	
	100 mg MgSO ₄ • 7H ₂ O	
	33.25 mg FeCl ₃ • 6H ₂ O	
	53 mg CaCl • 2H ₂ O	
	500 mg NH ₄ NO ₃	
	100 mg ciclohexymide	
MNBP	25 mg bacitracine	(Louvrier, Laguerre and Amarger, 1995)
	3 mg penicillin G	
	3.5 mg PCNB (P- Chloronitrobenzene)	
	500 µg biotin	
	500 µg thiamine	
	500 µg Ca pantothenate	
	5 mg benomyl	
	25 ppm Congo red	
	pH 6.8	
	16 g Bacto agar	
	5 ml 20% (w/v) Bacto Yeast extract	
	10 ml 10% (w/v) (NH ₄) ₂ SO ₄	
Pel agar	1 ml 1M MgSO ₄ • 7 H ₂ O	(Pemberton <i>et al.</i> , 2005)
	10 ml 50% (w/v) glycerol	
	250 ml 2% (w/v) polygalacturonic acid	
	200 ml Pel phosphate Buffer	
	13 g Oxoid nutrient broth	
Prt medium	30 g gelatine	(Cubitt <i>et al.</i> , 2013)
	16 g agar	
	15 g agar	
Potato dextrose agar (PDA)	4 g potato extract	(Atlas, 1996)
	20 g dextrose	
	16 g PDA	
	1 g Tryptone	
Potato dextrose agar (PDA) +10 % LB	0.5 g Yeast extract	
	0.55 g NaCl	
	4 g agar	
Swarming media	7.5 g Eiken agar	(Harshey and

Swimming media	10 g Tryptone	(Matsuyama, 1994)
	5 g Yeast extract	
	5 g NaCl	
	3 g agar	
Tryptic soy broth (TSB)	4 g Tryptone	(Monson <i>et al.</i> , 2013)
	2 g NaCl	
	17 g pancreatic digest of casein	
Tryptone yeast extract (TY)	5 g NaCl	(McCullough, 1949)
	3 g papaic digest of soy bean	
	2.5 g K ₂ HPO ₄	
	2.5 g glucose	
Yeast extract (YE)	6 g Tryptone	(Beringer, 1974)
	3 g Yeast extract	
	0.5 g CaCl ₂	
Yeast extract agar (YE agar)	5 g Yeast extract	(Grant and Pramer, 1962)
	30 g glucose	
Yeast mannitol (YMB)	As Yeast extract, additional, 20 g or 8 g agar	(Grant and Pramer, 1962)
	1 g mannitol	
	0.4 Yeast extract	
	0.1 g NaCl	
	0.5 g K ₂ HPO ₄	
	0.2 g MgSO ₄	
Yeast mannitol agar (YMB agar)	pH 6.8	(Vincent, 1970)
	As Yeast mannitol, additional 15 g agar	

Table 6. Antibiotics, antifungals and supplements used in this study.

Chemical (abbreviation)	Stock solutions preparation	Working concentration
Antibiotics and antifungal		
Bacitracine	5x stock, 125 mg/ml in dH ₂ O, stored at -20°C	25 mg/L
Benomyl	10x stock, 50 mg/ ml in 100% EtOH v/v, stored at -20°C	5 mg/L
Brilliant green (BG)	1x stock, 0.1mg/ml in dH ₂ O, stored at 4°C	1 mg /ml
Ciclohexymide	500x stock, 50mg/ml in 50% Et-OH v/v, stored 20°C	- 100 mg/L
PCNB	10x stock, 35 mg/ml in 100% acetone, stored at -20°C	3.5 mg/L
Penicillin G	10x stock, 30 mg/ml in dH ₂ O, stored at -20°C	3 mg/L
Supplement		
Biotin	100x stock, 5 mg/ml in dH ₂ O, stored at -20°C	500 µg/L

Pantothenate- Ca	100x stock, 5 mg/ml in dH ₂ O, stored at -20°C	500 µg/L
Thiamine	100x stock, 5 mg/ml in dH ₂ O, stored at -20°C	500 µg/L

Table 7. Solutions used in this study.

Solution	Components or ingredients per litre	Reference
Agarose gel mix	0.5-1.2% agarose in TBE 0.8 µl per 100 ml of SYBR® safe (Invitrogen) or 500 ng ethidium bromide per ml gel	
Alkaline lysis solution	2 N NaOH 10% SDS	
CAS-HDTMA	Mix 605 mg Chrome Azurol S with 500 ml dH ₂ O make 100 ml of (1mM FeCl ₃ •6 H ₂ O in 10mM HCl), mix with CAS solution mix 729 mg HDTMA with 400 ml dH ₂ O	(Schwyn and Neilands, 1987)
CTAB / NaCl (hexadecyltrimethyl ammonium bromide)	Dissolve 4.1 g NaCl in 80 ml of dH ₂ O and add 10 g of CTAB while heating (65°C) and stirring. Takes 3 hours to dissolve CTAB. Adjust final volume to 100 ml of H ₂ O and sterilize	(Wilson, 2001)
Deferrated Casamino Acids	Dissolve 10 g of Casamino Acids in 100 ml dH ₂ O extract with 100 ml of 3% 8-hydroxyquinolone in chloroform mix by shaking in a sealed container, leave settle for 20min remove aqueous layer and repeat extraction with equal volume (100 ml) of chloroform alone and autoclave	(Schwyn and Neilands, 1987)
Fahraeus media	0.1 g CaCl ₂ 0.12 g MgSO ₄ • 7H ₂ O 0.1 g KH ₂ PO ₄ 0.15 g Na ₂ HPO ₄ 0.005 g FeCl ₃ Micronutrients mix: 2.03 g MnSO ₄ • 7 H ₂ O 0.22 g ZnSO ₄ • 7 H ₂ O 0.08 g CuSO ₄ • 5 H ₂ O 0.08 g H ₂ MoO ₄ • H ₂ O 0.57 g H ₃ BO ₃ 10 g agar pH 6.5	(Fahraeus, 1957)
Jensen solution	74.5 mg KCl 174 mg K ₂ HPO ₄ 246.5 mg MgSO ₄ 1.8 mg iron citrate	(Somasegaran and Hoben, 1995)

Solution	Components or ingredients per litre	Reference
	Oligonutrients: 0.078 g $\text{CuSO}_4 \cdot 5 \text{H}_2\text{O}$ 0.22 g $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ 2.03 g $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$ 0.01 g $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$ 1.43 g H_3BO_3 pH 6.7	
Loading buffer	30% glycerol 0.25% Orange G (sigma)	
10 x MM9	3 g KH_2PO_4 5 g NaCl 10 g NH_4Cl pH 7	
Salt buffer	0.1 g NaCl 0.5 g K_2HPO_4 0.2 g MgSO_4 pH 6.8	(Vincent, 1970)
Pel phosphate Buffer	15 g Na_2HPO_4 (anhydrous) 0.7 g $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ pH 8.0	
Phage buffer	10 mM Tris-HCl pH 7.4 10 mM MgSO_4 0.01 % gelatin	
TBE	0.1 M Tris-HCl Boric acid 0.1 M 2 mM EDTA	
TE buffer	10 mM Tris-HCl 1 mM EDTA pH 8	

3.4 Plant material and techniques

Commercial seeds from *Medicago sativa* (alfalfa), *Lupinus albus* and *Macroptilium atropurpureum* were used in the present study. All seeds were stored at 4°C.

3.4.1 Seed sterilisation and germination

1. *M. sativa*: seeds were scarified in the presence of concentrated H_2SO_4 for 7 min followed by several washes with cold sterile water. Then seedlings were surface sterilised with

50% (v/v) NaClO for 90s followed again by several water washes and left overnight in sterile water. After 48h at 4°C in darkness, seeds were germinated in 0.8% water-agar plates at 22°C for 48h in darkness.

2. *L. albus* and *M. atropurpureum*: seeds were surface sterilised 1 min in EtOH 70% (v/v), 3min NaClO 25% followed by several washes with sterile water. Seeds were transferred to 0.8% water-agar plates and incubated 24h at 4°C in darkness; afterwards seeds were germinated at 22°C in darkness for 48h in the case of *L. albus* and 72h in the case of *M. atropurpureum*.

3.4.2 Trap plant assay

M. sativa, *L. albus* and *M. atropurpureum* were used as trap plants. Leonard jars containing Jensen's solution were filled with sterile vermiculite:soil (1:1 v:v) and two to three seedlings were transferred to each jar. Plants were cultivated in a greenhouse at 22°C under a 16/8-h light regime for three to five weeks.

3.4.3 Most probable number (MPN)

To calculate the most probable number (MPN) (Toomsan *et al.*, 1984), Leonard jars containing Jensen's solution were filled with sterile vermiculite and two to three seedlings were transferred to each jar. Serial dilutions of the soil (10^{-1} , 10^{-2} , 10^{-3}) were made the previous day in a salts buffer and left shaking overnight at 200 rpm (Infors HT Multitron standard) at 28°C. Each Leonard jar containing three seedlings corresponded to a given dilution. Seedlings were inoculated with 1 ml of each dilution. Plants were cultivated in a greenhouse at 22°C under a 16/8-h light regime for three to five weeks.

3.4.4 Isolation of Rhizobia from nodules

For Rhizobia isolation from nodules (symbiotic isolates), nodules were surface sterilised 1 min in EtOH 70% (v/v), 3min NaClO 25%, followed by several washes with sterile water. Nodules were individually crushed in 20 µl sterile water and the nodule content was streaked on YMB agar plates and incubated at 28°C for 48h. Colonies were re-streaked on YMB agar plates until pure cultures were obtained (Vincent, 1970).

3.4.5 Inoculation assay to test for nodulation capacity

In order to test the symbiotic capacity of the symbiotic strains isolated, surface-sterilised, germinated seedlings were inoculated with 1 ml of an exponentially-growing bacterial culture

and grown for three weeks in a growth chamber at 22°C under a 16/8-h light regime with a light intensity of 120 $\mu\text{E}/\text{m}^2\text{s}$. Nodule development was examined after three to five weeks.

3.5 Bacterial isolation and techniques

3.5.1 Total aerobic counts in soil

For total aerobic viable counts in Tomejil, Motril and Burguillos soils, serial soil dilutions (10^{-1} , 10^{-2} , 10^{-3} , 10^{-4} , 10^{-5} , 10^{-6} , 10^{-7}) were made in a salts buffer. 100 μl of each dilution was plated on the rich medium Tryptic soy broth (TSB). Plates were incubated at 28°C and single bacterial colonies were counted every day for three days.

3.5.2 Phenotypic characterisation

3.5.2.1 Carbon and Nitrogen sources

The GEN III MicroPlates (Biolog, Inc.) system was used for phenotypic characterisation of carbon and nitrogen sources utilization and sensitivity to some chemicals, following the manufacturer's instructions except for the following modifications. Strains were grown on TY plates, cultures were resuspended in IFA buffer and adjusted to an O.D 600nm of 0.2. Portions of 0.1 ml were added to each well, and plates were incubated at 28°C at 150 rpm on a rotary shaking (Infors HT Multitron standard) for 48h. For growth at different pH, Yeast Mannitol (YM) medium was adjusted to pH 8-9.6 with 50mM Tris.

3.5.2.2 Fatty acids identification

Cellular fatty acid methyl esters (FAMES) were prepared and separated using the method described by Sasser (1990) and identified with the MIDI Sherlock Microbial Identification System (version 6.1), using the TSBA6 database at the Spanish Type Culture Collection (CECT, Burjassot, Valencia). *Neorhizobium tomejilense* T20_22, T17_20 and T11_12 and the *Neorhizobium* sp. type strains were grown in TY for 48h at 28°C and harvested in late exponential growth phase.

3.5.2.3 MALDI-TOF analysis

To determine the whole-cell protein profile, a matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry (MALDI-TOF MS) analysis was carried out at the Scientific-Technical Services (University of Balearic Islands, Spain). Phenotypic characterisation of

freshly-grown colonies was carried as described by (Sanchez *et al.*, 2014). Strains were cultured on TY plates at 28°C for 48h. One microlitre of the extract (these extracts were obtained based on the ethanol/formic acid extraction procedure of Bruker Daltonics) was placed onto a spot of a ground steel plate. Each sample was overlaid with 1 µL of matrix solution (saturated solution of α -cyano-4-hydroxy-cinnamic acid in 50% acetonitrile–2.5% trifluoroacetic acid) and air dried at room temperature. Measurements were performed on an Autoflex III MALDI-TOF/TOF mass spectrometer (Bruker Daltonics, Leipzig, Germany). Spectra were recorded in the linear, positive mode in the mass range 2–20 kDa (IS1 = 20 kV, IS2 = 18.7 kV, lens = 6.50 kV, and PIE = 200 ns). Spectra were obtained for all strains analysed twice each with three replicates. The profiles obtained for each strain was analysed and compared, and the corresponding dendrogram was constructed with the MALDI BioTyper software (version 1.0; Bruker Daltonics).

3.5.3 Isolation of *Neorhizobium* from soil

For *Neorhizobium* isolation the methodology shown in Figure 3 was used. A 10^{-1} soil dilution was made in an Erlenmeyer flask using a salts buffer. Serial dilutions up to 10^{-6} , depending on the soil, were made, and 100 µl portions of each dilution were plated in the semi-selective media MNBP supplemented with 25 ppm of Congo red. Confirmed LB-negative colonies were grown on YMB agar plates and purified until pure cultures were obtained. Genomic DNA was extracted from the pure cultures using the alkaline lysis method and tested for *fnrN* PCR amplification. Positive *fnrN* isolates were phylogenetically characterised and screened for –if present– the symbiosis-related genes *nodC* and *nifH*.

3.5.4 Isolation of *Ensifer* from soil

For *Ensifer* isolation the methodology shown in Figure 3 was used. A 10^{-1} soil dilution was made in an Erlenmeyer flask using a salts buffer. Serial dilutions up to 10^{-6} depending on the soil were made and 100 µl of each dilution was plated on the semi-selective medium MNBP supplemented with 25 ppm of Congo red. Selected white pink colonies from MNBP were grown in YMB agar plates and purified until pure cultures were obtained. Genomic DNA was extracted from the pure cultures using the alkaline lysis method and tested for the *Ensifer* genus marker *asmA* in a PCR amplification. Positive *asmA* isolates were phylogenetically characterised and screened for –if present–the symbiosis-related genes *nodC* and *nifH*.

3.5.5 Isolation of *Bradyrhizobium* from soil

For *Bradyrhizobium* isolation, a detailed explanation of the methodology shown in Figure 3 is described here. A 10^{-1} soil dilution was made using the *Bradyrhizobium* recovery medium. Serial dilutions up to 10^{-6} depending on the soil were made in the same recovery medium and 100 μ l portions of each dilution were plated in the semi-selective medium BSJM supplemented with 25 ppm Congo red. Slow-growth, small white-pink colonies were selected, inoculated in YMB agar plates, incubated at 28°C for up to two weeks, and purified until pure cultures were obtained. Genomic DNA was extracted from the pure cultures using the alkaline lysis method and tested for *Bradyrhizobium* genus marker *thrC* PCR amplification. Positive *thrC* isolates were phylogenetically characterised and screened for –if present– the symbiosis-related genes *nodC* and *nifH*.

3.5.6 Isolation of *Serratia* from soil

For *Serratia* isolation a detailed explanation of the methodology shown in Figure 3 is described here. A 10^{-1} soil dilution was made using the *Bradyrhizobium* recovery medium. Serial dilutions up to 10^{-6} , depending on the soil, were made in the same recovery medium, and 100 μ l portions of each dilution were plated in the semi-selective medium BSJM supplemented with 25 ppm Congo red. White and red, fast-growing colonies were selected and grown in YMB agar plates at 28°C for 24h, and purified until pure cultures were obtained. Genomic DNA was extracted from the pure cultures using the alkaline lysis method and isolates were phylogenetically characterised using the 16S rDNA sequence and screened for the nitrogen-fixation gene *nifH*.

Figure 3. *Neorhizobium*, *Ensifer*, *Bradyrhizobium* and *Serratia* direct isolation methods.

3.5.7 Phenotypic *Serratia* assays

3.5.7.1 Swarming and swimming

To test for swarming or swimming capacity, swarming and swimming plates were prepared by pouring 25 ml of the corresponding medium the day before use. Cultures were grown overnight at 30°C in 5 ml of Luria-broth (LB), cultures were adjusted to $OD_{600} \approx 1$ with LB, and 2.5 μ l portions from each sample were spotted in triplicate on the centre of swarming or swimming plates. Plates were carefully closed and incubated at 30°C for up to two days (swarming) or overnight (swimming).

3.5.7.2 Protease activity and plant cell wall-degrading enzymes: cellulase and pectate lyase

To test for cellulase, pectate lyase, or protease activities, agar plates were prepared by pouring 25 ml of CMC-Sucrose, Pel or Prt media, respectively, the day before use. Cultures were grown overnight at 30°C and 37°C in 5 ml LB, OD_{600} was adjusted to $OD_{600} \approx 1$ with LB, and 2.5 μ l of each sample was spotted on CMC-Sucrose plates. Up to 4-5 samples were spotted in the same plate, in addition to the positive and negative controls, that were: *D. solani* MK10 and *E. coli* K-12 MG1655 respectively, for cellulase and pectate lyase, and *Serratia plymuthica* S13 and *E. coli* K-12 MG1655, respectively, for protease. Once the spots were totally dry, plates were inverted and incubated at 30°C for 48h (for cellulase and pectate lyase) or overnight (for protease). Plates were developed as follows: i) for cellulase, by the addition of a 0.2% Congo Red solution for 20min, followed by a rinse with 1 M NaCl and a further incubation with 1 M NaCl for 15min. Finally, plates were incubated with 1 M HCl for 5min at room temperature and excess HCl was removed (Monson *et al.*, 2013). Cellulase production is indicated by the presence of a purple halo surrounding spots on a blue background; ii) for pectate lyase, plates were treated with 7.5% copper (II) acetate $Cu(CO_2CH_3)_2$ for 1h at room temperature. After the incubation period excess copper (II) acetate was removed (Cubitt *et al.*, 2013). Pectate lyase activity production is indicated by the presence of a semi-transparent halo surrounding spots on a light blue background; iii) for protease, plates were treated with 4 M ammonium sulphate $(NH_4)_2SO_4$ for 5min at room temperature. After incubation, excess ammonium sulphate was removed (Cubitt *et al.*, 2013). Protease production is indicated by a transparent halo surrounding spots.

3.5.7.3 Quorum sensing

To test whether the strains could produce N-acyl homoserine lactones signal molecules (AHLs), two sensor systems were used. The first sensor was *Chromobacterium violaceum* CV026. This strain is a mini-Tn5 mutant, violacein-negative, white mutant (CV026), defective in the production of AHLs but capable of sensing them by turning purple (McClellan *et al.*, 1997). This strain can be induced by AHLs and N-acylhomocysteine thiolactones (AHTs) with N-acyl side chains from C₄ to C₈ in length. However, this biosensor is limited by its weak response to short C₄ AHL chains. For that reason, an additional biosensor strain was used, *Serratia* sp. ATCC 39006 SP19 is a *pigX* and *pigZ* mutant with prodigiosin hyperproduction. This sensor is highly sensitive to short C₄ acyl-AHL chains and responds by turning orange (Poulter *et al.*, 2010).

In order to test for AHL production, cultures were grown overnight at 30°C and 37°C in 5 ml LB, and OD₆₀₀ was adjusted to OD₆₀₀≈1 with LB. In a 5 ml tube, 4 ml of 0.7% LB agar (top agar) were mixed with 100 µl of either a *C. violaceum* CV026 or a *Serratia* sp. ATCC 39006 SP19 culture. The mixture was inverted a couple of times and was quickly poured onto LB plates as an overlay. Once the top agar was solidified, 2.5 µl portions of each culture sample were spotted on the surface of the agar lawns containing sensor strains. Up to 4-5 samples were spotted per plate. Once the spots were totally dry, plates were incubated overnight at 30°C. AHL production is indicated by a red/orange or purple halo surrounding spots for sensor strains *Serratia* sp. ATCC 39006 SP19 or *C. violaceum* CV026, respectively.

3.5.7.4 Siderophore activity

To test for siderophore activity, blue agar CAS assay plates were used (Schwyn and Neilands, 1987). The following media and solutions had to be made separately: 10 x MM9, CAS-HDTMA Deferrated Casamino Acids and CAS Agar. After autoclaving, they were added (per L) in the following order: i) 100 ml 10 x MM9, ii) 30 ml 10% Deferrated Casamino Acids, iii) 10 ml 20% Glycerol, iv) 1 ml 1 M MgCl₂ and v) 1 ml 1 M CaCl₂. After mixing, 100 ml CAS-HDTMA solution was added with constant mixing. The final medium had a uniform blue/green colour and plates were made with 25 ml of this mixture. Cultures were grown overnight at 30°C and 37°C in 5 ml LB, cultures were adjusted to OD₆₀₀≈1 with LB and 2.5 µl of each sample was spotted in the plate. Up to 4-5 samples were spotted in the same plate, in addition to the positive and negative controls *S. plymuthica* S13 and *E. coli* K-12 MG1655, respectively. Once the spots were totally dry, plates were incubated 48h at 30°C.

Siderophore production is indicated by an orange halo surrounding spots on a blue/green background.

3.5.7.5 Antibiotic production activity

To test for antibiotic production, an *E. coli* β -lactam super sensitive strain (ESS or *E. coli* ESS2231) (Nishida *et al.*, 1977) was used as indicator. Cultures were grown overnight at 30 °C and 37°C in 5 ml LB. In a 5 ml tube, 4 ml of 0.7% LB agar (top agar) were mixed with 100 μ l of ESS culture. The mixture was inverted a couple of times and was quickly poured onto LB agar plates as an overlay. Once the top agar was solidified, 2.5 μ l portions of each sample were spotted on the surface of the ESS agar lawn. Up to 7 samples were spotted in per plate, in addition to the positive control *Serratia* sp. 39006. Once the spots were totally dry, plates were incubated overnight at 30°C. Antibiotic production is indicated by a transparent halo surrounding spots.

3.6 Bacteriophage techniques and assays with *Serratia*

3.6.1 Bacteriophage stock and titrate preparation

In order to obtain phage lysates, phages were first titrated by making serial dilutions (neat, 10^{-1} , 10^{-2} and 10^{-3}) of phage lysates in phage buffer. In a 5 ml tube, 10 μ l of each phage dilution was mixed with 100 μ l of an overnight culture of the host bacteria and 3 ml top 0.35% LB agar, and the mixture was inverted a couple of times and quickly poured onto LB agar plates as an overlay. Plates were incubated overnight at the appropriate temperature. Those plates showing near-confluent lysis (a mosaic of phage plaques with very little remaining bacterial lawn), were chosen for lysate preparation. To the selected plate, 1 ml phage buffer was added and, with the help of a glass rod, the top agar was scraped off and poured into a glass tube containing 3 ml of phage buffer. Chloroform (500 μ l) was added and the mixture was vortexed vigorously for 2min. After 30min at room temperature, mixtures were centrifuged at 2,200 x g for 20min, 4°C. The supernatant was decanted into a glass bijoux tube for storage with 100 μ l of chloroform to help minimize contamination. Phage lysates were stored at 4°C. In order to roughly estimate high phage titres, serial dilutions of the phage stock were made in LB (neat, 10^{-1} , 10^{-2} , 10^{-3} , 10^{-4} , 10^{-5} , 10^{-6} , 10^{-7} and 10^{-8}) and mixed in a 5 ml tube with 100 μ l of an overnight culture of the host bacteria and 3 ml 0.35% top LB agar. The mixture was inverted a couple of times and quickly poured onto LB plates as an overlay. With a multichannel pipette, 2.5 μ l portions of each of the dilutions were

spotted on the surface of the phage agar lawn. Once the spots were totally dry, plates were incubated overnight at the corresponding temperature. For each phage, those dilutions where single plaques could be seen were used to calculate plaque forming units per ml (pfu/ml) and efficiency of plating (EOP) using the following formulas:

Plaque forming units per ml (pfu ml⁻¹):

Number of plaques / dilution factor x amount plates in ml

Efficiency of plating (EOP):

Titre on test strain / titre on control strain

3.6.2 Phage sensitivity assay

To test for phage sensitivity, decimal serial dilutions (neat through 10⁻⁸) of the phage stocks were made using LB, and mixed with 100 µl of an overnight culture of the strain to be tested and 3 ml top 0.35% LB agar in a 5 ml tube. The tube was inverted a couple of times and quickly poured onto LB plates as an overlay. Once the top agar was solidified, 2.5 µl portions of each of the phage stock dilutions were spotted with a multichannel pipette on the surface of the agar lawn and left to dry. A total of three phages were spotted per plate. Plates were incubated overnight at the appropriate temperature before scoring lysis, and therefore phage sensitivity.

3.7 Fungal techniques and assays with *Serratia*

3.7.1 Yeast growth inhibition

To test for anti-yeast activity, YE plates were made with 25 ml of medium. *Serratia* and *Saccharomyces cerevisiae* cultures were grown overnight at 30°C in 5 ml LB or YE, respectively, and adjusted to OD₆₀₀ ≈ 1 with LB. In a 5 ml tube, 4 ml of 0.8% YE top agar were mixed with 100 µl *S. cerevisiae* culture. The mixture was inverted a couple of times and quickly poured onto LB plates as an overlay. Once the top agar was solidified, 2.5 µl portions of each *Serratia* culture were spotted on the surface of the fungal agar lawn. Up to 7 samples were spotted per plate, in addition to the positive control *D. solani* MK10. Once spots were totally dry, plates were incubated overnight at 30°C. Anti-yeast activity is indicated by a transparent halo surrounding spots.

3.7.2 Fungal activity against *Verticillium dahliae*

For fungicide assays, the protocol developed by (Matilla *et al.*, 2012b) was used. Indicator top agar of *V. dahliae* was prepared by vortexing a 5-mm fungal plug in 10 ml sterile distilled water. Then, 15 ml PDA + 10% LB was added and mixed, and 5 ml of top lawn were poured onto PDA + 10% LB plates. 5 µl portions of overnight cultures of the strains to be tested were spotted on the surface of the fungal agar lawn and incubated for 7–10 days at 25°C.

3.8 Recombinant DNA techniques

3.8.1 DNA purification and visualisation

3.8.1.1 Genomic DNA extractions

For PCR amplifications, DNA was extracted using a fast alkaline lysis protocol (Birnboim and Doly, 1979). Cells were picked from a plate using a toothpick, 20 µl of lysis solution added, and samples were incubated 15min at 90°C. After the incubation period, 100 µl sterile water was added to the sample followed by a centrifugation of 5min at 12.000 x *g* and the DNA was recovered from the supernatant.

For total bacterial genome sequencing, rhizobia samples were grown in TY medium. CTAB (Cetyl Trimethyl Ammonium Bromide) bacterial genomic DNA isolation method was generally employed following the protocol described by Wilson (2001). In cases where extraction was not successful, the MoBio PowerSoil® DNA isolation kit was used followed by a DNA clean-up and concentration step as indicated by the manufacturer's instructions.

3.8.1.2 DNA quantification

Genomic DNA quantification and quality was measured by Nanodrop spectrophotometry (NanoDrop Technologies, Wilmington, USA) and by fluorescence (Qubit, Invitrogen by Life Technologies Holdings Pte. Ltd, Singapore). Integrity and purity were checked by electrophoresis in a 0.8% agarose gel.

3.8.1.3 Agarose gel electrophoresis

DNA fragment size was checked by electrophoresis, using horizontal agarose gels submerged in TBE buffer. Agarose gels were prepared as described (Table 7). 6x loading buffer was added to samples prior to loading in the gel, and commercial linear size

standards (Nippon Genetics, Germany) were included in every gel. Gels were visualised using the Gel Doc™ EZ Imager (Bio-Rad Laboratories, California, USA).

3.8.1.4 DNA clean-up after enzymatic reactions

Following PCR amplification or restriction digests, unincorporated small DNA fragments were removed either directly from enzymatic reactions mixes or by extraction from electrophoresis gels using the NucleSpin® gel and PCR clean-up (Macherey-Nagel) or QIAquick (Qiagen) kits, according to the manufacturers' instructions. DNAs were eluted and stored in sterile water at -20 °C.

3.8.2 DNA manipulations

3.8.2.1 Polymerase chain reaction (PCR)

Standard PCR reactions were carried out with standard Taq DNA polymerase (Biotools, Madrid). A high-fidelity Phusion Taq DNA polymerase (New England Biolabs or Thermo Scientific) was used in cases when fragments could not be amplified with Taq DNA polymerase (Biotools), even changing PCR conditions, or when high fidelity was required, for instance during cloning. Annealing temperatures and extension times were adjusted according to the primers used, shown in Table 8. Table 9 and Table 10 show standard PCR components and conditions respectively.

Table 8. PCR components.

Components	Final volume (25µl)	Final volume (25µl)
	Taq Biotools®	Taq Phusion®
5X Buffer	2.5µl	5µl
dNTPs	0.5µl	0.5µl
Primer 5'	1µl	0.5µl
Primer 3'	1µl	0.5µl
DMSO (if present)	1µl	0.75
DNA Polymerase	1µl	0.25
DNA	1-2 µl	1-2 µl
dH2O	up to 25µl	up to 25µl

Table 9. PCR standard conditions.

Cycle	Temperature	Time	No. cycles
Initial denaturation	95°C	3 min	1
Denaturation	95°C	30 s	30
Annealing	50-64°C	30 s	
Extension	72°C	30 s / Kb	
Final extension	72°C	2 min	1
Hold	12°C	∞	1

3.8.2.2 Oligonucleotide primers

Oligonucleotide primers used in this work are listed in Table 10, and were obtained from Sigma-Aldrich.

Table 10. Primers used in this study.

Use	Name	Sequence (5'- 3')	Annealing T°C	Reference
<i>Bradyrhizobium</i> <i>m</i> sp. marker	F_threonine synthase_387	TCGATGGGCAAGGCG	56	This study
	R_threonine synthase_682	CCCATSGTCTTCTTGCC		
<i>Bradyrhizobium</i> <i>m</i> sp. full 16S rDNA amplification	Bradj16S70F	GCGGGCGTAGCAATACGTCA GC	68	(Jones <i>et al.</i> , 2016)
	Bradj16S1430R	GCCGGCTGCCTCCCTTGCGG GTTA		
<i>Ensifer</i> sp. marker	F_Asma_1113	ACGCTCGCCTTCGACCG	64	This study
	R_Asma_1635	CCTTGCGCACGAGATCGG		
Full 16S rDNA amplification	fD1	ccgaattcgtcgacaacAGAGTTTGA TCCTGGCTCAG	55	(Weisburg <i>et al.</i> , 1991)
	rDI	cccgggatccaagcttAAGGAGGTG ATCCAGCC		
Partial <i>glnII</i> amplification	GSII-1	AACGCAGATCAAGGAATTCG	56	(Turner and Young, 2000)
	GSII-4	GCGAACGATCTGGTAGGGGT		
Partial <i>nifH</i> universal amplification	DVV_nifH_413_ R	ATIGCRAAICCCICRCAIACIACR TC	58	(Ando <i>et al.</i> , 2005)
	IGK3_nifH_19_F	GCIWHTHTAYGGIAARGGIGGIA THGGIAA		

Use	Name	Sequence (5'- 3')	Annealing T°C	Reference
Partial <i>nodC</i> universal amplification	nodC_F_540	TGATYGAYATGGARTAYTGGC T	50	(Sarita <i>et al.</i> , 2005)
	nodC_R_1160	CGYGACARCCARTCGCTRTTG		
Partial <i>recA</i> amplification	recA_41F	TTCGGCAAGGGMTGRTSAT G	55	(Vinuesa <i>et al.</i> , 2005b)
	recA_640R	ACATSACRCCGATCTTCATGC		
Partial <i>rpoB</i> amplification	F_rpoB_583	GARTTCGACGCCAAGGAYAT	49	(Jorrín, 2016)
	R_rpoB_1200	GAAGAACAGCGAGTTGAACAT		
<i>Rhizobium</i> sp. screening	F_Rleg_FnrN	GGTGATGCCGGACGGGCG	54	(Jorrín, 2016)
	R_Rleg_FnrN	TGGAGCAGGCTTGCGACCTT		

3.8.3 DNA sequencing methods

3.8.3.1 Sanger sequencing

Amplicon and plasmid Sanger DNA sequencing was outsourced to StabVida (Lisbon, Portugal). Sequence chromatograms were analysed with 4Peaks v 1.7.2 software (Nucleobytes B.V., Aalsmeer, The Netherlands), and nucleotide and protein database searches were done using BLAST (Basic Local Alignment Search Tool) (Altschul *et al.*, 1990).

3.8.3.2 Whole genome sequencing

Individual genomes were sequenced externally either at the ASU Biodesign Institute (Illumina MiSeq v.3, 600 bp PE libraries, 2 x 300 bp reads, from 0.6 to 1.3 Mreads; ASU Genomics Core, Tempe, AZ, USA) or at Microbes NG (Illumina MiSeq v3, PE 2x300 bp reads; University of Birmingham, Birmingham, UK).

3.9 Bioinformatics

3.9.1 Phylogenetic analysis

For phylogenetic and sequence analyses, nucleotide sequences obtained from PCR amplifications of 16S rDNA, *recA* and *glnII* genes were corrected and assembled if necessary with SerialCloner2-6 (http://serialbasics.free.fr/Serial_Cloner.html) and 4peaks

(<http://nucleobytes.com/4peaks/>) and aligned with the ClustalW algorithm (Chenna *et al.*, 2003) in MEGA 6.0 (Tamura *et al.*, 2013). Randomized Axelerated Maximum Likelihood (RAxML) (Stamatakis, 2014) was used for the construction of phylogenetic trees. Phylogenetic trees were visualised in FigTree c1.4.3 (<http://tree.bio.ed.ac.uk/software/figtree/>) and edited with Adobe Illustrator CS5 (<http://www.adobe.com/products/illustrator.html>).

3.9.2 Genome assembly and annotation

For genome analyses, original fasta files were quality filtered with Trimmomatic (Bolger, Lohse and Usadel, 2014), assembled with SPAdes (Bankevich *et al.*, 2012) and annotated either with the NCBI Prokaryotic Genome Annotation Pipeline (Tatusova *et al.*, 2016) or with the Prokka package (Seemann, 2014).

3.9.3 Genomic comparisons

Genomes were studied at different levels. Synteny among different genomes was studied with Mauve (Darling, Mau and Perna, 2010). First, genomic contigs were reordered according to the reference genome with Mauve contig reordering. Afterwards, reordered genomes were aligned against each other. In order to study genomic identity among strains, average nucleotide identity (ANIm, calculation based on MUMmer; or ANIb, calculation based on BLAST) was calculated with JSpeciesWS online server (Richter *et al.*, 2016). Distances were calculated as 1- ANI and trees were computed by the UPGMA (Sneath and Sokal, 1973) method using MEGA 7 package (Kumar, Stecher and Tamura, 2016). Alternatively, distance dendrograms were generated by hierarchical cluster analysis of 100 - % ANI matrices (Chan *et al.*, 2012) with StataSE v.14.0 (StataCorp, College Station, TX, USA) after computation of Euclidean distances with the Average Linked method.

3.10 Data analyses

In order to try and identify similarities within large datasets, Principal component analysis (PCA) (Abdi and Williams, 2010) was carried out (script developed by R. E. Monson).

RESULTS

4.1 *Neorhizobium*

4.1.1 Direct isolation of *Neorhizobium* from soil

A total of twenty-eight independent isolations were carried out from Tomejil soil. After four days, pink-white colonies appearing on MNBP plates were chosen as putative fast-growing Rhizobia. From those, we chose to discard all that were able to grow on LB agar (Figure 3; Section 3.5.3). In our hands, this includes most *Ensifer* spp. and many non-rhizobial isolates. However, most *Rhizobium* spp. were unable to grow on the 5 g·l⁻¹ NaCl present in LB agar, and thus this step resulted in a further enrichment of *Rhizobium* spp. (Table 11).

Table 11. Summary of results obtained for *Neorhizobium* sp. isolation.

Soil	No. colonies tested for <i>fnrN</i> marker	Total no. <i>fnrN</i> positive soil isolates	Total no. (%) <i>Neorhizobium</i> sp. ¹
Tomejil	342	99	53 (55.5%)

¹Assigned from the 16S rRNA and *rpoB* sequences of *fnrN* positive strains.

DNA from three hundred and forty-two of these putative *Rhizobium* spp. isolates was amplified for the *fnrN* gene by PCR (Figure 6A). This gene is important for microaerobic metabolism (Gutiérrez *et al.*, 1997) and has been found in all sequenced *Rhizobium* spp. and *Agrobacterium* spp., but not in other Rhizobia. Ninety-nine of the DNAs resulted in good amplification of a major band of a size similar to what was expected. However, two different band sizes were observed. While 46 strains resulted in amplification of a PCR band of the expected size (279 bp), the remaining 53 strains amplified a larger fragment (353 bp, Figure 6A). In order to reduce the number of isolates for further characterisation, a fragment of the *rpoB* gene was amplified and sequenced from all of them. The *rpoB* marker has been shown to be very effective in discriminating phylogenetically close bacteria (Mollet, Drancourt and Raoult, 1997; Khamis *et al.*, 2003; Case *et al.*, 2007), including Rhizobia (Jorin and Imperial, 2015). Forty-six of the ninety-nine partial *rpoB* sequences corresponded, as expected, to *Rhizobium* sp. (data not shown), while the remaining fifty-three, surprisingly, clustered with sequences from genus *Neorhizobium* (Figure 5). Total *Neorhizobium* abundance estimation using the direct isolation method was $2.98 \times 10^4 \pm 2.47 \times 10^4$ CFU/g of soil. These data were compared with the total viable aerobic bacteria counts using TSB as a rich medium in soil, which resulted in $8.43 \times 10^7 \pm 9.81 \times 10^6$ CFU/g of soil. The *rpoB* sequences obtained were compared and classified into groups (twenty in total) if they

differed in at least one nucleotide (Supplemental Table S2). In order to facilitate further studies, a representative strain of each of these *rpoB* sequence types was chosen. Near complete 16S rDNA sequences were obtained from these representative strains and compared against those in databanks, confirming that they formed a diverse clade that was closely related to known members of the genus *Neorhizobium*, but distant from the *Agrobacterium-Rhizobium* group (Figure 4).

Figure 4. Phylogenetic tree of representative Tomejil soil isolates based on PCR amplified near-complete 16S rDNA (1,234 bp) sequences. Maximum likelihood trees (RAxML) were derived from ClustalW alignments. T: Tomejil soil isolates. The number of strains within each genotype group is indicated within parentheses. Trees include sequences from type strains of *Neorhizobium* species, as the closest taxonomic relatives, and of *A. tumefaciens*, as outgroup. Bootstrap support (1,000 replications) for the different nodes is indicated. Bars represent the number of substitutions per base. Genbank accession numbers are listed on Supplemental Table S1.

Figure 5. Phylogenetic tree of representative Tomejil soil isolates based on PCR amplified partial *rpoB* (356 bp) sequences. Maximum likelihood trees (RAxML) were derived from ClustalW alignments. T: Tomejil soil isolates. The number of strains within each genotype group is indicated within parentheses. Trees include sequences from type strains of *Neorhizobium* species, as the closest taxonomic relatives, and of *A. tumefaciens*, as outgroup. Bootstrap support (1,000 replications) for the different nodes is indicated. Bars represent the number of substitutions per base. Genbank accession numbers are listed on Supplemental Table S1.

4.1.2 Genotypic and genomic characterisation of *Neorhizobium* isolates

Since all previously characterised *Neorhizobium* isolates had been obtained from nitrogen-fixing legume root nodules, we tried to amplify *nodC* and *nifH*, markers related to symbiosis and nitrogen fixation, respectively, from our Tomejil isolates. Negative results were obtained in all cases, suggesting that these isolates were non-symbiotic. Since direct tests of

symbiotic ability with legume plants were hampered by the fact that the known hosts of *Neorhizobium* sp. (*Galega officinalis* and *G. orientalis*, (Mousavi *et al.*, 2014); *Sesbania herbacea*, (Wang *et al.*, 1998); *Caragana intermedia*, (Li Lu *et al.*, 2009)) are very diverse and are not found in the Tomejil area, we reasoned that obtaining and characterizing the genome sequence of some of the Tomejil isolates, even at draft level, would help clarify their symbiotic / non-symbiotic status.

We chose five representatives of the most abundant/diverse *rpoB* groups, so as to try to obtain a picture of the genomic diversity within this *Neorhizobium* sp. clade. Genomic DNA was sequenced and assembled to draft level and this assembly was used for subsequent analyses. All five isolates had similar genome size and G+C (%) composition (Table 12).

Table 12. Genomic features of *Neorhizobium* genomes sequenced in this work.

Strain	Number of contigs	Largest contig (bp)	Total genome length (bp)	G+C (%)	N50
T20_22	37	1,052,711	6,608,977	61.47	508,270
T7_12	52	1,197,185	6,627,103	61.44	347,929
T25_27	27	1,446,028	6,462,352	61.49	734,252
T25_13	42	721,675	6,322,993	61.56	419,328
T6_25	69	488,027	6,750,064	61.35	186,129

We first searched for *fnrN* genes. This was important because no *fnrN* gene has been described in *Neorhizobium* sp. isolates, and it was absent from the *Neorhizobium* sp. genomic sequences available in databanks (data not shown). Using the well-characterised *R. leguminosarum* *fnrN* gene for BLAST comparisons, we were unable to find any relevant hit against genomic sequences from Tomejil isolates, despite the fact that the isolates were chosen because they showed clear amplification of single (although of different size) band with *fnrN* primers (Figure 6A). Amplified PCR bands from some of the isolates were sequenced and compared with databanks. Part of the amplified region showed similarity with

genes encoding a poly (3-hydroxybutyrate) depolymerase from *Rhizobia*, including *Neorhizobium galegae* (but not *N. alkanisoli* or *N. huautlense*, data not shown). This gene, together with an upstream ORF, was present in genome assemblies of the Tomejil strains and contained sequences partially complementary to those of *fnrN* primers that may explain the successful amplifications observed (Figure 6B).

Figure 6. PCR amplification of a genomic DNA band from Tomejil *Neorhizobium* sp. strains with *fnrN* primers. (A) Agarose gel electrophoresis separation of PCR products in DNAs from *Neorhizobium* species type strains and *Neorhizobium* Tomejil strains (353 bp). *R. leguminosarum* bv. *viciae* 3841 (279 bp) was used as positive control. (B) Schematic representation of the genomic region amplified by *fnrN* primers in Tomejil *Neorhizobium* sp. strains. Genomic sequences partially complementary to *fnrN* primers were identical in all Tomejil strains: F_ *fnrN* (5'-GGTGATGCCGGACGGGCG-3') hybridized to 5'-ccgctcGCCGGtCGGGCG -3', and R_ *fnrN* (5'- TGGAGCAGGCTTGCGACCTT-3') hybridized to 5'-cGGAAtCAGGtTTGCGACCTT-3'.

We then tried to find *nod* and *nif* genes by running BLAST searches of known *Neorhizobium* *nod* and *nif* genes against draft genome sequences, with negative results in all cases (data not shown). Since *nod* and *nif* genes are harboured on large megaplasmids in *N. galegae*

(Österman *et al.*, 2014), we searched for similar plasmids in the genomes of Tomejil strains. All five genomes contained a set of *repABC* sequences characteristic of rhizobial plasmids (Cevallos, Cervantes-Rivera and Gutierrez-Rios, 2008; Pinto, Pappas and Winans, 2012), highly similar (85-86%) to the *repABC* cluster from the 1.8 Mb megaplasmid from *N. galegae* bv. *orientalis* (Table 13) and that are probably responsible for the replication a large plasmid in each of these strains. This is supported by the fact that *repABC* homologues are present in large contigs (83-523 kb) in the Tomejil strains, despite the draft level quality of their genome sequences (27-69 contigs, Table 12). Multiple sequence alignment of these *repABC* sequences suggests that plasmids from the Tomejil strains are highly-related and separated from the symbiotic plasmids from *N. galegae* bv. *orientalis* and *N. galegae* bv. *officinalis* (Figure 7). The completely sequenced strain *N. galegae* bv. *officinalis* HAMBI 1140 harbours an additional megaplasmid (175 kb) that appeared to be highly unrelated to both *N. galegae* symbiotic plasmids and plasmids from Tomejil strains (Table 13, Figure 7).

Table 13. Presence of *repABC* regions in the genomes of Tomejil strains. DNA regions similar to the 3,628 bp region containing *repABC* genes from the symbiotic megaplasmid from *N. galegae* bv. *orientalis* HAMBI 540 were located in genome sequences by BLAST, extracted and compared by multiple alignment (ClustalW).

Genome	Number of <i>repABC</i> regions	(%) identity to HAMBI 540	size of contig (bp)
<i>N. galegae</i> bv. <i>orientalis</i>			
HAMBI 540	1	100	1,807,065
<i>N. galegae</i> bv. <i>officinalis</i>			
HAMBI 1141	2	95	1,638,739
T20_22	1	53	175,279
T7_12	1	85	376,046
T25_27	1	85	523,062
T25_13	1	85	263,545
T6_25	1	86	721,675
	1	85	82,656

Figure 7. Phylogenetic tree based on *repABC* sequences from Tomejil sequences and *Neorhizobium* megaplasms. Sequences similar to the *N. galegae* bv. orientalis HAMBI 540^T 1.8 Mb megaplasid *repABC* region (3,628 bp) were extracted from Tomejil draft genome sequences and from *N. galegae* bv. officinalis HAMBI 1140 genome sequence. Sequences were aligned with ClustalW, and a Neighbor-Joining consensus tree derived. Bootstrap support (1,000 replications) for the different nodes is indicated. Bar represents the number of substitutions per base. Genbank accession numbers are shown in Supplemental Table S1.

In order to ascertain the phylogenetic relationships among the sequenced Tomejil strains, and between this clade and the type strains of the three *Neorhizobium* species: *N. galegae*, *N. alikisoli*, *N. huautlense*, we pulled out the complete sequences of the genes *atpD*, *glnII*, *recA*, *rpoB* and *thrC* from the genome assemblies. These genes have been previously used successfully to define the *Neorhizobium* genus and its species (Mousavi *et al.*, 2014). Figure 8 shows a maximum-likelihood phylogenetic tree derived from a multiple alignment of the concatenated genes, rooted by using the *Agrobacterium tumefaciens* Ach5 sequences. Very similar results were obtained when the phylogenetic tree was derived by the neighbour-joining method (data not shown). There was very strong support for two clades within the *Neorhizobium* genus; one formed by *N. alikisoli* and *N. huautlense*, and the other by *N. galegae* and the Tomejil strains. Four of the Tomejil strains (T7_12, T20_22, T25_13, T25_27) formed a well-supported group clearly separated from the fifth strain, T6_25, and all five, in turn, clearly separated from *N. galegae*.

Figure 8. Phylogenetic tree of sequenced Tomejil strains and of *Neorhizobium* type strains based on a concatenation of complete *atpD*, *glnII*, *recA*, *rpoB*, and *thrC* genes (8,949 bp). Maximum likelihood trees (RAxML) were derived from ClustalW alignments. T: Tomejil soil strains. The tree includes the *A. tumefaciens* type strain as outgroup. Bootstrap support (1,000 replications) for the different nodes is indicated. Bar represents the number of substitutions per base. Genbank accession numbers are shown in Supplemental Table S1.

Genome-wide comparisons were carried out by calculating pairwise average nucleotide identities (Richter and Rossello-Mora, 2009; Richter *et al.*, 2016) with the above genomes. Both ANIm and ANIb scores were calculated, and the ANIb matrix is shown in Table 14. A distance matrix was generated from the ANIb matrix using 100-ANIb (Chan *et al.*, 2012) and Euclidean distances. These were represented in the dendrogram shown in Figure 10. The dendrogram faithfully reproduced the topology of the multilocus phylogenetic tree (Figure 9). Using a 95% ANI value as a widely accepted delimiter for genomic species (Richter and Rossello-Mora, 2009), our results are consistent with Tomejil strains representing at least a clearly differentiated genospecies within *Neorhizobium*, and probably two, represented by strains T6_25 on one hand, and T7_12, T20_22, T25_13, and T25_27, on the other.

Table 14. Pairwise average nucleotide identities (ANiB) between representative *Neorhizobium* sp. Tomejil isolates and *Neorhizobium* sp. type strains.

Strain name	<i>N. galegae</i> HAMBI 540 ^T	<i>N. alkalisoli</i> DSM 21826 ^T	<i>N. huautlense</i> DSM 21817 ^T	T7_12	T25_27	T25_13	T6_25	T20_22
<i>N. galegae</i>								
HAMBI 540 ^T	*	84.75	84.51	91.70	91.70	91.76	91.59	91.71
<i>N. alkalisoli</i>								
DSM 21826 ^T	84.74	*	89.40	84.77	84.83	84.79	84.75	84.76
<i>N. huautlense</i> DSM 21817 ^T								
DSM 21817 ^T	84.52	89.41	*	84.61	84.66	84.61	84.50	84.59
T7_12	91.70	84.77	84.61	*	97.43	96.47	95.51	97.43
T25_27	91.69	84.83	84.66	97.43	*	96.45	95.53	97.73
T25_13	91.76	84.79	84.61	96.47	96.45	*	95.50	96.46
T6_25	91.59	84.76	84.51	95.51	95.53	95.50	*	95.51
T20_22	91.71	84.76	84.59	97.43	97.73	96.46	95.51	*

Figure 9. Dendrogram representation of a Euclidean distance matrix derived from pairwise average nucleotide identities (ANIb) between Tomejil and *Neorhizobium* type strain genomes. The vertical red line indicated the 95 % ANI threshold.

4.1.3 Description of a new species, *N. tomejilense* sp. nov.

We set out to investigate whether any of the Tomejil *Neorhizobium* strains are in fact representatives of a new *Neorhizobium* species. T20_22 represented the genotype with the highest number of strains, twelve in total. Although we genome sequenced three isolates from this group (T20_22, T17_20 and T11_12), only T20_22 was included in the *Neorhizobium* diversity analyses as the genotype representative. However, among the subset of isolates from the most abundant genotypes whose genome was sequenced to draft level, strains T20_22, T17_20 and T11_12 appeared to be quite homogeneous. All three isolates had similar genome size and G+C (%) composition (Table 15). These were further characterised, and the genomic, phylogenetic and phenotypic data obtained support that they should be classified as a new *Neorhizobium* species.

Table 15. Genomic features of *Neorhizobium* sp. T20_22, T17_20 and T11_12 genomes sequenced in this work.

Strain	Number of contigs	Largest contig (bp)	Total genome length (bp)	G+C (%)	N50
T20_22	37	1,052,711	6,608,977	61.47	508,270
T17_20	27	1,714,448	6,563,491	61.46	1,151,780
T11_12	26	2,227,499	6,514,075	61.47	1,529,274

4.1.4 Phylogenetic analysis

The analysis of 16S rDNA confirmed the placement of T20_22, T17_20 and T11_12 within the *Neorhizobium* genus. A ML tree derived from full 16S rDNA alignments is shown in Figure 10; similar results were obtained using MP and NJ methods (data not shown). Since the genus *Neorhizobium* is a combination of earlier described *Rhizobium* species, sequences from type strains of close *Rhizobium* species were included to support their phylogenetic differences. Tomejil isolates clustered together (in fact, they have identical 16S rDNA gene sequences), close to *N. galegae* HAMBI 540^T and clearly separated from the other *Neorhizobium* species, *N. huautlense* DSM 21817^T and *N. alkalisoli* DSM 21826^T, and from the genus *Rhizobium*, with a very high bootstrap support.

Figure 10. Maximum likelihood phylogenetic tree based on 16S rDNA sequences (1,476 bp) of *Neorhizobium tomejilense* sp. nov., *Neorhizobium* type strains and representative *Rhizobium* type strains. Bootstrap support (1,000 replications) for the different nodes is indicated. Bar represents the number of substitutions per base. Genbank accession numbers are shown in Supplemental Table S1.

An in-depth evaluation at species level was performed with a multilocus sequence analyses of the full-length, concatenated genes *atpD*, *glnII*, *recA*, *rpoB* and *thrC*, used to describe the *Neorhizobium* genus. NJ and MP methods (data not shown) gave results similar to those shown in the ML tree in Figure 11. In this case, and due to the higher resolution obtained, T20_22, T17_20 and T11_12 also kept closely related to *N. galegae* HAMBI 540^T, but were clearly grouped in a different branch with very high bootstrap support. These results therefore suggest that this group of Tomejil soil strains belong to a new *Neorhizobium* species differentiated from all other described *Neorhizobium*.

Figure 11. Maximum likelihood phylogenetic tree based on full *atpD*, *glnII*, *recA*, *rpoB* and *thrC* concatenated sequences (8,386 nt) of *Neorhizobium tomejilense* sp. nov, and sequences from *Neorhizobium* type strains and type strains from representative *Rhizobium* species. Bootstrap support (1,000 replications) for the different nodes is indicated. Bar represents the number of substitutions per base. Genbank accession numbers are shown in Supplemental Table S1.

4.1.5 Genome properties and ANI values

In order to provide support for the classification of these Tomejil isolates as a novel species, average nucleotide identity (ANI) was used to estimate homologies at the genomic level including all the *Neorhizobium* species described. Draft genomes were obtained from T20_22, T17_20, T11_12 and two type species *N. huautlense* DSM 21817^T and *N. alkalisoli* DSM 21826^T whose genomic sequence were not available in public databases. Calculated ANIb values are indicated in Table 16. Pairwise ANI genomic comparisons between T20_22, T17_20 and T11_12 was above 97%, over the ≈95-96% boundary set for species definition (Richter and Rossello-Mora, 2009). Within previously described *Neorhizobium* species, *N. galegae* HAMBI 540^T showed the highest (91.7%) ANI value with T20_22, T17_20 and T11_12, while it was quite distant from *N. huautlense* DSM 21817^T and *N. alkalisoli* DSM 21826^T. These results confirm what had been observed with the 16S rDNA and concatenated *atpD*, *glnII*, *recA*, *rpoB* and *thrC* phylogenetical analyses. In order to avoid clonality of the isolates, we performed a Randomly Amplified Polymorphic DNA- Polymerase chain reaction (RAPD-PCR) using M13 primers. *N. tomejilense* T20_22, T17_20 and T11_12 showed distinctive band profiles in the agarose gel among them. These profiles were also

different to the *Neorhizobium* species, in particular to *N. galegae* HAMBI 540^T, the closest species at phylogenetic and genomic level to *N. tomejilense* (Figure 12).

Table 16. Pairwise average nucleotide identities (ANIb) among T20_22, T7_12, T25_27 and type strains from the different *Neorhizobium* species.

Strain name	<i>N. galegae</i> HAMBI 540 ^T	<i>N. alkalisoli</i> DSM 21826 ^T	<i>N. huautlense</i> DSM 21817 ^T	<i>N. tomejilense</i> T20_22	<i>N. tomejilense</i> T17_20 ^T	<i>N. tomejilense</i> T20_22
<i>N. galegae</i> HAMBI 540 ^T	*	79.34	78.89	90.11	90.10	90.12
<i>N. alkalisoli</i> DSM 21826 ^T	79.49	*	87.36	79.56	79.49	79.52
<i>N. huautlense</i> DSM 21817 ^T	79.26	87.52	*	79.50	79.46	79.49
<i>N. tomejilense</i> T11_12	90.22	79.41	79.03	*	99.29	99.54
<i>N. tomejilense</i> T17_20 ^T	90.16	79.38	79.04	99.31	*	99.19
<i>N. tomejilense</i> T20_22	90.06	79.39	79.12	99.43	99.03	*

Figure 12. M13-RAPD-PCR band pattern fingerprints of *N. tomejilense* and type strains of *Neorhizobium* species: 1, *N. galegae* HAMBI 540^T; 2, *N. huautlense* DSM 21817^T; 3, *N. alkanisoli* DSM 21826^T; 4, *N. tomejilense* T20_22; 5, *N. tomejilense* T11_12; 6. *N. tomejilense* T17_20^T.

Symbiotic genes are usually found as part of mobile genetic elements in Rhizobia (Sullivan *et al.*, 1995) and therefore they should not be taxonomically relevant. However, they provide valuable information to understand symbiotic specificity and legume host range in Rhizobia. T20_22, T17_20 and T11_12 do not harbour symbiotic genes in their genomes, as concluded from the PCR screening with *nifH* and *nodC* universal primers, and also from BLAST genomic searches (data not shown).

4.1.6 Phenotypic analysis

Differential phenotypic characteristics of the strains in the proposed new species and *Neorhizobium* species are indicated in Table 17. In the case of carbon and nitrogen sources, *N. tomejilense* was not able to use L-aspartic acid, in comparison to the other *Neorhizobium* species. In contrast, it was able to use pectin and sorbitol in comparison to its closest *Neorhizobium* species (*N. galegae* HAMBI 540^T). *N. tomejilense* strains were positive for many antibacterial and antifungal compounds, which was not the case for *N. galegae* HAMBI 540^T for example. They were also able to use many proton and electron donors, this was differential to *N. alkanisoli* DSM 21826^T. *N. tomejilense* and *N. huautlense* DSM 21817^T exhibit many differential characteristics, as *N. huautlense* DSM 21817^T showed to have a wide metabolic capacity, this was differential not only of *N. tomejilense* but of the other *Neorhizobium* species.

Table 17. Differential characteristics among *N. tomejilense* strains T11_12, T17_20, T20_22 and type strains from *Neorhizobium* species.

Growth	1	2	3	4	5	6
37 °C	-	+	-	-	-	-
pH 8	+	+	+	+	+	-
pH 10	-	-	-	-	-	-
Carbon and Nitrogen sources:						
D-Sorbitol	-	+	+	+	+	+
Pectin	-	+	-	+	+	+
Dextrin	w	+	-	-	-	-
D-Lactic Acid Methyl Ester	-	+	-	-	-	-
Mucic Acid	-	w	+	-	-	-
D-Malic Acid	+	+	-	+	+	+
D-Serine	-	+	-	-	-	-
L-Serine	-	+	+	-	-	-
L-Aspartic Acid	+	+	+	-	-	-
L-Histidine	-	+	-	-	-	-
D-Aspartic Acid	-	+	-	-	-	-
L-Pyroglutamic Acid	-	+	-	-	-	-
N-Acetyl Neuraminic Acid	+	w	-	-	-	-
Propionic Acid	+	+	-	+	+	+
Formic Acid	-	+	-	+	+	+
Proton and electron donors:						
Sodium Butyrate	-	+	-	+	+	+
Tetrazolium Blue	+	+	-	+	+	+
Potassium Tellurite	+	+	-	+	+	+
Antibiotic and antifungal resistance:						
Troleandomycin	+	+	-	+	+	+
Lincomycin	-	+	+	+	+	+
Vancomycin	-	+	+	+	+	+
Nalidixic Acid	-	+	+	+	+	+
Fusidic Acid	-	+	-	+	+	+

Species / Strains: 1, *N. galegae* HAMBI 540^T; 2, *N. huautlense* DSM 21817^T; 3, *N. alkalisoli* DSM 21826^T; 4, *N. tomejilense* T11_12; 5, *N. tomejilense* T17_20^T; 6. *N. tomejilense* T20_22. All data were obtained in this study. +, positive; -, negative; w, weak.

In addition to the classical phenotypic analysis we also performed a whole-cell matrix-assisted laser-desorption time-of-flight mass spectrometry (WC MALDI-TOF-MS) fingerprint analysis. The dendrogram (Figure 13) representing the distances calculated from the fingerprint profiles among type strains from the *Neorhizobium* species and the Tomejil strains shows a clearly distinct group formed by the three Tomejil strains (T20_22, T17_20 and T11_12) separated from all the *Neorhizobium* type species described.

Figure 13. Dendrogram constructed by applying average linkage clustering and Pearson's distance correlation from a similarity matrix of identical mass peaks, computed from whole cell mass spectra of all *Neorhizobium* strains analysed. The dendrogram was generated by considering the average value of the duplicates for each strain. The bar indicates distance.

Fatty acid composition and abundance varied among all *Neorhizobium* species (Table 18). Fatty acid abundance did not change significantly among *N. tomejilense* strains, that had very similar fatty acid composition, except for 17:0 3OH fatty acid in strain T11_12. Fatty acid pattern and abundance in *N. tomejilense* strains was clearly different to those in the most similar strain within the genus, *N. galegae* HAMBI 540^T. *N. galegae* HAMBI 540^T has distinct fatty acids such as 17:0 cyclo, 20:2 w6,9c, and summed feature 1, which differed from *N. tomejilense*, as well as from *N. huautlense* DSM 21817^T and *N. alkalisoli* DSM 21826^T.

Table 18. Fatty acid patterns of *N. tomejilense* strains T11_12, T17_20, T20_22 and type strains from *Neorhizobium* species.

	1	2	3	4	5	6
15:1 w8c	0.29	0.32	-	-	-	-
16:00	13.22	7.07	11.69	9.41	8.77	8.64
16:0 3OH	2.83	2.55	2.53	2.99	3.13	3.19
17:00	1.58	0.64	-	1.71	0.88	1.11
17:0 cyclo	0.65	-	-	-	-	-
17:0 3OH	0.77	-	-	1.45	-	-
17:1 w8c	0.79	-	-	1.32	0.72	0.65
18:00	0.68	2.57	0.75	0.82	1.41	1.81
18:0 3OH	0.80	1.41	0.64	1.29	0.81	1.19
18:1 w7c 11-methyl	0.66	-	0.33	-	-	-
18:1 2OH	-	0.98	-	-	-	-
19:0 cyclo w8c	27.49	3.50	11.51	7.51	9.52	7.06
19:0 10-methyl	1.84	1.80	2.27	2.17	2.50	2.51
20:2 w6,9c	0.61	-	-	-	-	-
Summed Feature 2	4.71	5.12	4.75	5.02	5.71	5.35
Summed Feature 3	1.72	0.66	1.61	0.89	0.89	1.31
Summed Feature 8	41.12	73.39	63.92	65.43	65.66	67.18
Summed Feature 1	0.25	-	-	-	-	-

Strains: 1, *N. galegae* HAMBI 540^I; 2, *N. huautlense* DSM 21817^I; 3, *N. alkalisoli* DSM 21826^I; 4, *N. tomejilense* T11_12; 5, *N. tomejilense* T17_20^T; 6, *N. tomejilense* T20_22. All data were obtained in this study. Summed features are groups of two or three fatty acids that cannot be separated by GLC with the MIDI system. Summed Feature 1, 13:0 3OH/15:1 I; Summed Feature 2, 14:0 3OH/16:1 iso I; Summed Feature 3, 16:1 w7c/16:1 w6c; Summed Feature 8, 18:1 w7c/18:1 w6c.

Description of *Neorhizobium tomejilense* sp. nov.

Neorhizobium tomejilense (to.me.jil.en'se. N.L. neut. adj. *tomejilense*, pertaining to Tomejil, the name of the experimental farm from which the type strain was isolated).

Cells are Gram-negative, aerobic, motile, non-spore-forming rods. Colonies on YMB are circular, cream-coloured, semi-translucent and mucilaginous. Older colonies have a white centre. Growth occurs at 12-30°C and pH 6-8 on YMB with optimum growth at 28°C at pH 7. Generation time is 2.7h. Growth occurs in YMB supplemented with 1% NaCl at pH 8. Uses L-glutamic acid and glucuronamide as nitrogen sources. Uses methyl pyruvate, ψ -amino-butyric acid, D-maltose, D-fructose, L-fucose, D-arabitol, D-galactose, D-trehalose, D-gluconic acid, L-lactic acid, glycerol, D-salicin, D-cellobiose, L-malic acid, acetic acid as carbon sources. Resistant to troleandomycin, lincomycin, vancomycin, nalidixic acid and fusidic acid. The predominant fatty acids are summed feature 8 (18:1 w7c/18:1 w6c), 16:00 and 19:0 cyclo w8c. The G+C content of the type strain is 61.46 mol% (derived from genome sequence).

The type strain, T17_20 (LMG 30623) and Genbank genome accession number PRJNA434362), was isolated from the Tomejil experimental farm, IFAPA (Instituto de Investigación y Formación Agraria y Pesquera (Centro Las Torres-Tomejil, Junta de Andalucía), Spain and presents no symbiotic genes.

4.2 *Ensifer*

4.2.1 Development of an *Ensifer* genetic marker

Highly-conserved, single copy marker genes were identified using the pipeline described by Wu, Jospin and Eisen (2013). Nine complete or permanent draft *Ensifer* sp. genomes: *E. meliloti* 1021, *E. meliloti* 2011, *E. meliloti* RMO17, *E. meliloti* BL225C, *E. meliloti* SM11, *E. meliloti* Rm41, *E. meliloti* GR4, *E. meliloti* AK83, *E. medicae* WSM 419, and *E. fredii* NGR 234, were selected for the analyses. Genbank accession numbers of the genomes are included in Supplemental Table S1. Gene *asmA* (outer membrane assembly factor) represented the best candidate, since it had a high grade of conservation in all *Ensifer* sp. genomes and it was present in a single copy in the chromosome. *asmA* is 1,761 bp long in the genus *Ensifer*. Primers were designed flanking the most conserved region of the gene, between positions 1,113 and 1,635, that amplify a 522 bp fragment. In order to test the specificity of the marker, PCR amplification was done with several *Ensifer* sp. type strains, and other related and laboratory strains (Figure 14). The *asmA* marker proved to be very specific and generally applicable to *Ensifer* sp. strains. The complete *asmA* alignment is shown in Supplemental Figure S1.

Figure 14. PCR amplification of the *Ensifer* marker gene *asmA* (522 pb) from *Ensifer* laboratory strain collection and from other laboratory strains.

4.2.2 *Ensifer* direct isolation

The direct isolation strategy consisted of a combination of the semi-selective culture medium MNBP already described (Louvrier, Laguerre and Amarger, 1995) and used for direct

isolation of *Rhizobium leguminosarum* from soils, used together with the *Ensifer* marker *asmA*, as shown in Figure 3 and detailed in Section 3.5.4. A total of three and six independent isolations were carried out from Tomejil and Motril soils, respectively. The results obtained with Tomejil and Motril soils are shown in Table 19. Eighteen and twenty-eight strains were isolated from Tomejil and Motril soils, respectively, out of two hundred and twenty-one (8,15%) and three hundred and ten (9%) colonies tested, respectively. Interestingly, 90% and 85% of the isolates that tested positive in the *asmA* PCR screening in Tomejil and Motril soils, respectively, were true *Ensifer* sp. isolates. We seek to analyse the *asmA* marker sequence conservation throughout type and isolated strains. To that end, we sequenced the amplified *asmA* marker fragment (456 bp), performed a multiple alignment, and estimated its degree of conservation (Table 20). Genbank accession numbers are included in Supplemental Table S1. *E. meliloti* 1021 was chosen as the reference sequence. Percentage identity among strains varied from 100% to 55.9%, suggesting that the *asmA* sequence marker is not only highly conserved and specific, but it also allows ample variability within the sequence.

Table 19. Summary of results obtained with the *asmA Ensifer* genus marker.

Soil	No. colonies tested	% true positives ¹	Total no. <i>Ensifer</i> sp. strains
Tomejil	221	90	18
Motril	310	85	28

¹Assigned from the 16S rDNA, *recA* and *glnII* sequences of *asmA* positive strains.

Table 20. *asmA* marker gene (456 bp) conservation in *Ensifer* sp. strains.

Strain	% identity	Source of <i>asmA</i> marker gene
<i>Ensifer meliloti</i> 1021 (reference)	100.0	Genome from Genbank
<i>Ensifer meliloti</i> 2011	100.0	Genome from Genbank
<i>Ensifer meliloti</i> RMO17	100.0	Genome from Genbank
<i>Ensifer meliloti</i> BL225C	100.0	Genome from Genbank
<i>Ensifer meliloti</i> SM11	98.9	Genome from Genbank
<i>Ensifer meliloti</i> Rm41	98.9	Genome from Genbank

Strain	% identity	Source of <i>asmA</i> marker gene
<i>Ensifer meliloti</i> GR4	98.9	Genome from Genbank
<i>Ensifer meliloti</i> AK83	98.9	Genome from Genbank
<i>Ensifer medicae</i> WSM 419	85.5	Genome from Genbank
<i>Ensifer fredii</i> NGR 234	72.2	Genome from Genbank
<i>Ensifer meliloti</i> 1021	100.0	Laboratory strain collection, this work
<i>Ensifer americanum</i> CFNBI 156 ^T	72.7	Laboratory strain collection, this work
<i>Ensifer fredii</i> USDA 191	100.0	Laboratory strain collection, this work
<i>Ensifer xinjiangense</i> LMG 17930 ^T	74.0	Laboratory strain collection, this work
<i>Ensifer arboris</i> LMG 14919 ^T	55.9	Laboratory strain collection, this work
T1_71	100.0	Tomejil isolate, this work
T1_81	99.3	Tomejil isolate, this work
T1_27	85.5	Tomejil isolate, this work
T1_45	100.0	Tomejil isolate, this work
T1_100	98.9	Tomejil isolate, this work
T2_8	99.3	Tomejil isolate, this work
.M4_29	95.4	Motril isolate, this work
.M4_46	95.8	Motril isolate, this work

Estimations of total *Ensifer* soil populations by the direct isolation method were compared with those for symbiotic *Ensifer* soil populations calculated by a Most Probable Number (MPN) method (Toomsan *et al.*, 1984) using *Medicago sativa* as a trap plant (Table 21). MPN counts of symbiotic *Ensifer* with *M. sativa* in Tomejil and Motril soils were low ($10^1 \cdot \text{g}^{-1}$). In contrast, total *Ensifer* soil counts were much higher, 10^4 CFU per gram of soil in both Tomejil and Motril soils. This higher abundance (up to three order of magnitude) of non-symbiotic vs. symbiotic *Ensifer* provides a rationale for our observation that all soil isolates failed to amplify the symbiotic markers *nodC/nifH*, while all alfalfa nodule isolates did amplify the symbiotic markers (Table 22).

Table 21. Total viable aerobic bacteria count (CFU/g), *Ensifer* CFU/g and nodule MPN/g counts in Tomejil and Motril soils.

Soil	TSB CFU/g	Soil <i>Ensifer</i> CFU/g	Nodule (<i>M. sativa</i>) MPN/g
Tomejil	$8.43 \times 10^7 \pm$	$4.44 \times 10^4 \pm$	4.30×10^1
	9.81×10^6	1.92×10^4	95% confidence level 9-180
Motril	$5.28 \times 10^7 \pm$	$2.46 \times 10^4 \pm$	2.00×10^1
	1.30×10^7	7.54×10^3	95% confidence level 4.5-42

Table 22. *Ensifer* isolates obtained from Tomejil and Motril soils.

Soil	Soil isolates (<i>nodC</i> ⁻ / <i>nifH</i> ⁻)	Nodule isolates (<i>M. sativa</i>) (<i>nodC</i> ⁺ / <i>nifH</i> ⁺)
Tomejil	18	31
Motril	28	10

4.2.3 Characterisation of *Ensifer* populations in Tomejil and Motril soils

16S rDNA, *recA* and *glnII* PCR amplifications and sequencing were carried out for all forty-six *Ensifer* isolates from soil and forty-one *Ensifer* isolates from nodules of *M. sativa*. Isolates were classified into genotypes according to their 16S rDNA full sequence: if sequences differed in one or more nucleotides they were classified as different genotypes. Nodule isolates were distributed into six and five different genotypes in the case of the nodule isolates from Tomejil and Motril soils, respectively. Tomejil and Motril soil isolates, on the other hand, were distributed into six different genotypes in both cases (Supplemental Table S2). Multilocus sequence analysis with the phylogenetically-relevant, housekeeping genes 16S rDNA, *recA* and *glnII* of representative strains from Tomejil and Motril soils and type strains for different *Ensifer* species are shown in Figure 15 and Figure 16. Tomejil and Motril isolates distributed mainly within *E. meliloti*, *E. medicae*, *E. numidicus* and *E. adherens*. The more resolutive phylogenetic analyses of the concatenated housekeeping genes *recA* and *glnII* (Figure 17) grouped the strains in three main clusters. Cluster 1 grouped with *E. medicae* and included all the strains that nodulated *M. sativa* in Tomejil soil (TMN_25, TMN_1, TMN_26, TMN_14, TMN_13, TMN_12), one non-symbiotic

Tomejil soil isolate (T1_27) and one *M. sativa* nodule isolate from Motril (MMN_6). Cluster 2 was related, but not identical, to *E. meliloti*, and included only Motril isolates, several non-symbiotic (M_4_45, M_5_5, M_7_2, M_4_46) and a *M. sativa* nodule strain (MMN_5). Cluster 3 grouped within *E. meliloti* and included Motril *M. sativa* nodule isolates (MMN_7, MMN_3, MMN_1) and some Tomejil non-symbiotic isolates (T_1_100, T_1_45). Finally, T2_8 soil isolate did not associate with any *Ensifer* sp. type species, while M5_14 soil isolate appeared to be related to *E. adherens*. Only one of the three clusters (Cluster 2) was specific for a particular soil. In the other two clusters, representative strains from both soils were obtained. Regarding the distribution of nodule and soil isolates, in the case of Tomejil, all nodule isolates grouped with *E. medicae*, whereas soil isolates spread across *E. meliloti*/*E. medicae* and presented a higher diversity. In the case of Motril, soil isolates clustered together except for M5_14; in contrast, nodule isolates were more heterogeneously spread.

Figure 15. Phylogenetic tree of representative Tomejil and Motril soil and nodule isolates based on PCR amplified near complete 16S rDNA sequences (1,277 bp) and type strains from *Ensifer* species. Maximum likelihood trees (RAxML) were derived from ClustalW alignments. TMN: Tomejil *Medicago sativa* Nodule isolates; MMN: Motril *Medicago sativa* Nodule isolates; T: Tomejil soil isolates; M: Motril soil isolates. Symbiotic isolates are indicated by a green square; non-symbiotic isolates are indicated by a red square. The number of strains with a given genotype is shown in parentheses. Bootstrap support (1,000 replications) for the different nodes is indicated. Bar represents the number of substitutions per base. GenBank accession numbers are shown in Supplemental Table S1.

Figure 16. Phylogenetic tree of representative Tomejil and Motril soil and nodule isolates based on PCR amplified partial concatenated *recA* (372 bp) and *glnII* (471 bp) sequences and type strains from *Ensifer* species. Maximum likelihood trees (RAxML) were derived from ClustalW alignments. TMN: Tomejil *Medicago sativa* Nodule isolates; MMN: Motril *Medicago sativa* Nodule isolates; T: Tomejil soil isolates; M: Motril soil isolates. Symbiotic isolates are indicated by a green square; non-symbiotic isolates are indicated by a red square. The number of strains within each genotype group is indicated within parentheses. Bootstrap support (1,000 replications) for the different nodes is indicated. Bar represents the number of substitutions per base. GenBank accession numbers are shown in Supplemental Table S1.

4.2.4 Genome comparison between a soil non-symbiotic isolate and a nodule symbiotic isolate from the same soil

Since, for the first time, we were able to isolate symbiotic and non-symbiotic *Ensifer* strains from the same soil, we decided to choose two closely related strains, one symbiotic and one non-symbiotic, for further comparison. *Ensifer* sp. Motril nodule isolate MMN_5 and nonsymbiotic *Ensifer* sp. Motril soil isolate M4_45, that appeared to be very closely related (Figure 15 and Figure 16) were genome sequenced to high-quality draft, and their genome characteristics are shown on Table 23. MMN_5 and M4_45 showed very similar G+C (%) content, 62.18 and 62.36, respectively. The genome size of the non-symbiotic isolate M4_45 was slightly smaller (6.34 Mb) than that of the nodule isolate (6.63 Mb), consistent with the absence of symbiotic genes. Pairwise average nucleotide identity (ANI) was calculated for *Ensifer* sp. MMN_5, *Ensifer* sp. M4_45 and other *E. meliloti* strains, since this was the closest species according to phylogenetic analyses (Table 24). ANIb values indicated that MMN_5 and M4_45 were highly similar in the aligned portions of their genome sequences. Regarding their similarity with *E. meliloti* strains, Motril isolates showed low similarity < 95%, below species level with the all the *E. meliloti* strains included in the analyses. Therefore this could indicate that MMN_5 and M4_45 and the rest of Cluster 2 (Figure 15 and Figure 16) could correspond to a new genospecies.

Table 23. Genomic features of *Ensifer* sp. MMN_5 and *Ensifer* sp. M4_45 genomes sequenced in this work.

Strain	Number of contigs	Largest contig (bp)	Total genome length (bp)	G+C (%)	N50
<i>Ensifer</i> sp. MMN_5	111	558,778	6,628,752	62.18	285,319
<i>Ensifer</i> sp. M4_45	40	1,098,544	6,339,782	62.36	421,852

Table 24. Pairwise average nucleotide identities (ANIb) between *Ensifer* sp. M4_45, *Ensifer* sp. MMN_5 and *Ensifer meliloti* genomes.

Strains	<i>Ensifer</i> sp. M4_45	<i>Ensifer</i> sp. MMN_5	<i>E. meliloti</i> RMO17	<i>E. meliloti</i> 1021	<i>E. meliloti</i> 2011	<i>E. meliloti</i> Rm41	<i>E. meliloti</i> BL225C	<i>E. meliloti</i> AK83	<i>E. meliloti</i> SM11	<i>E. meliloti</i> RU11/001	<i>E. meliloti</i> L5-30
<i>Ensifer</i> sp. M4_45	*	98.94	93.28	93.44	93.44	93.17	93.30	93.14	93.21	93.25	93.34
<i>Ensifer</i> sp. MMN_5	98.83	*	93.40	93.52	93.51	93.32	93.49	93.13	93.34	93.28	93.32
<i>E. meliloti</i> RMO17	93.14	93.33	*	99.04	99.04	98.97	99.26	98.73	99.09	99.02	98.94
<i>E. meliloti</i> 1021	93.39	93.50	99.10	*	100.00	98.72	99.26	98.69	98.94	98.81	98.68
<i>E. meliloti</i> 2011	93.40	93.55	99.12	100.00	*	98.74	99.27	98.70	99.00	98.86	98.70
<i>E. meliloti</i> Rm41	92.68	92.86	98.55	98.24	98.25	*	98.43	98.31	98.58	98.55	99.21
<i>E. meliloti</i> BL225C	93.17	93.33	99.19	99.10	99.11	98.75	*	98.60	98.89	98.76	98.77
<i>E. meliloti</i> AK83	92.98	93.10	98.71	98.58	98.55	98.81	98.78	*	98.76	98.64	98.92
<i>E. meliloti</i> SM11	92.87	93.05	98.86	98.75	98.74	98.78	98.73	98.57	*	99.43	98.85
<i>E. meliloti</i> RU11/001	92.90	92.90	98.88	98.66	98.65	98.76	98.72	98.44	99.42	*	98.77
<i>E. meliloti</i> L5-30	92.98	92.94	98.65	98.35	98.34	99.36	98.55	98.67	98.68	98.70	*

For comparative genomics, MMN_5 and M4_45 genomes were aligned against each other using Mauve (Darling *et al.*, 2010). Genomes were first reordered individually using the *E. meliloti* 1021 genome as reference (Figure 17). The corresponding reordered genomes were then aligned against each other (Figure 18). The alignment between *Ensifer* sp. MMN_5 and *Ensifer* sp. M4_45 revealed highly syntenic chromosomes in contrast to the pSym plasmids. This synteny was analysed in detail by separately comparing the different replicons. As shown on Table 25, both estimated size and percentage G+C (%) of the chromosome and plasmids of MMN_5 and M4_45 showed values similar to those of the reference genome. Furthermore, M4_45 pSymA plasmid reported a smaller size, as it would be expected in

view of the absence of symbiotic genes. We also performed separate pairwise ANIb analyses for chromosomes and plasmids of MMN_5, M4_45 and *E. meliloti* 1021 (Table 26). Lower ANIb values were obtained for the aligned portions of the genome corresponding to the pSyms, in particular to pSymA, in contrast to the chromosome. This was a clear trend shown in all comparisons, between MMN_5 and M4_45 and with the reference genome *E. meliloti* 1021. Taken together, these results suggest that native *Ensifer* sp. populations undergo plasmid adaptations to symbiosis.

Figure 17. Progressive Mauve genome alignments after contig reordering with the reference genome *Ensifer meliloti* 1021. A) Contig reordering between *E. meliloti* 1021 and *Ensifer* sp. M4_45 (non-symbiotic isolate from motril soil). B) Contig reordering between *E. meliloti* 1021 and *Ensifer* sp. MMN_5 (symbiotic isolate from motril soil). Red lines mark contig boundaries.

Figure 18. Progressive Mauve genome alignment between *Ensifer* sp. M4_45 (Motril non-symbiotic isolate) and *Ensifer* sp. MMN_5 (Motril symbiotic isolate). Contigs were first reordered individually with the reference genome *Ensifer meliloti* 1021 and then aligned against each other using progressive Mauve. The region containing symbiotic genes in MMN_5 genome is marked with a red star.

Table 25. Size and G+C (%) comparison of the chromosome and pSyms of the reference genome *E. meliloti* 1021, *Ensifer* sp. MMN_5 and *Ensifer* sp. M4_45.

Genome		Chromosome	pSymB	pSymA
<i>E. meliloti</i> 1021	Size	3,654,135	1,683,333	1,354,226
	G+C (%)	62.7	62.4	60.4
<i>Ensifer</i> sp. MMN_5	Size	3,627,149	1,794,996	1,206,607
	G+C (%)	62.8	62.4	60.1
<i>Ensifer</i> sp. M4_45	Size	3,571,509	1,744,340	1,024,061
	G+C (%)	62.8	62.4	60.8

Table 26. Pairwise average nucleotide identities (ANIb) calculated for individual replicons from *Ensifer* sp. MMN_5, *Ensifer* sp. M4_45, and the reference genome *E. meliloti* 1021.

Strain name	<i>E. meliloti</i> 1021	<i>Ensifer</i> sp. MMN_5	<i>Ensifer</i> sp. M4_45
Chromosome			
<i>E. meliloti</i> 1021	*	94.92	94.96
<i>Ensifer</i> sp. MMN_5	94.80	*	99.89
<i>Ensifer</i> sp. M4_45	94.78	99.93	*
pSymB			
<i>E. meliloti</i> 1021	*	93.60	93.43
<i>Ensifer</i> sp. MMN_5	93.42	*	98.53
<i>Ensifer</i> sp. M4_45	93.29	98.43	*
pSymA			
<i>E. meliloti</i> 1021	*	90.51	90.43
<i>Ensifer</i> sp. MMN_5	90.82	*	96.56
<i>Ensifer</i> sp. M4_45	90.64	96.96	*

4.3 *Bradyrhizobium*

4.3.1 *Bradyrhizobium* semi-selective medium BSJM optimisation and marker development

The BSJM semi-selective medium was developed for the specific isolation of soybean symbionts of the *B. japonicum* and *B. elkanii* species (Tong and Sadowsky, 1994). In order to ensure that a high diversity of *Bradyrhizobium* species –including those that do not establish symbiosis with soybeans– were able to grow on this medium, several *Bradyrhizobium* type and other strains available in the laboratory collection were grown on the BSJM medium (Table 27). All twenty-three *Bradyrhizobium* strains assessed grew on it, except for *B. retamae* Ro19, *Bradyrhizobium* sp. RSA 104 and *B. icense* LmTR13. BSJM contains an AG medium base (Tong and Sadowsky, 1994) supplemented with the heavy metals cobalt and zinc, Brilliant Green (BG) and PCNB. We modified the original BSJM medium formulation by adding cycloheximide (300 mg/L) and Congo Red (25 ppm). Cycloheximide helped control eukaryotic growth –especially fungal– an important consideration due to the long incubation periods of the plates (2 weeks), and Congo Red helped identify putative rhizobia growing on the plates, since these acquired to a white/pink colour. The final BSJM medium formulation is listed in Table 5.

Table 27. *Bradyrhizobium* strains tested for growth on semi-selective medium BSJM.

Strain	BSJM medium growth
<i>B. betae</i> LMG21987 ^T	+
<i>B. canariense</i> BTA-1 ^T	+
<i>B. cytisi</i> CTAW11 ^T	+
<i>B. diazoefficiens</i> USDA 110 ^T	+
<i>B. elkanii</i> USDA 76 ^T	+
<i>B. icense</i> LMTR 13 ^T	-
<i>B. iriomotense</i> LMG24129 ^T	+
<i>B. japonicum</i> USDA 6 ^T	+
<i>B. jicamae</i> PAC68 ^T	+
<i>B. lablabi</i> CEBAU 223086 ^T	+
<i>B. liaonigense</i> LMG18230 ^T	+
<i>B. lupini</i> ISLU 101	+

Strain	BSJM medium growth
<i>B. pachyrhizi</i> PAC 48 ^T	+
<i>B. paxllaeri</i> LmTR 21 ^T	+
<i>B. retamae</i> Ro19 ^T	-
<i>B. rigense</i> CTAW71 ^T	+
<i>Bradyrhizobium</i> sp. Btai-1	+
<i>Bradyrhizobium</i> sp. Lmj C	+
<i>Bradyrhizobium</i> sp. P.22A	+
<i>Bradyrhizobium</i> sp. RSA 104	-
<i>Bradyrhizobium</i> sp. ORS 278	+
<i>Bradyrhizobium valentinum</i> LmjM3 ^T	+
<i>B. yuanmingense</i> LMG21827 ^T	+

Despite the exceptions, it was clear that BSJM medium would allow growth of a wide range of *Bradyrhizobium* strains and thus it would be appropriate for our purpose. The next step was the development of a specific genus marker for *Bradyrhizobium* that would allow the identification using a PCR screening. Thirty-two complete or permanent draft *Bradyrhizobium* genomes from databases were selected for the analyses (Table 28). Genome accession numbers are included in Supplemental Table S1. Highly conserved single copy maker genes were identified using the pipeline described by (Wu, Jospin and Eisen, 2013). Among all the candidate genes, we selected *Threonine synthase* (*thrC*) as the best target in view of its high grade of conservation in all of the *Bradyrhizobium* genomes and of its putative, relevant function. The *Threonine synthase* gene is 1,761 bp long in *Bradyrhizobium* and is involved in step 5 of the subpathway that synthesizes L-threonine from L-aspartate:

O-Phospho-L-homoserine + H₂O $\rightleftharpoons$ L-Threonine + Orthophosphate (Shames *et al.*, 1984)

Primers were designed in the most conserved regions of the gene, between nucleotide positions 387 and 683, making up a 296 bp band in the PCR amplification. In order to test for specificity of the marker, we first performed a PCR amplification with the *Bradyrhizobium* strains available in the laboratory collection and other, related bacteria from the laboratory collection (Figure 19). The *thrC* marker was successfully amplified from all the tested

Bradyrhizobium strains, except for *Bradyrhizobium* sp. Btai-1, whose *thrC* gene sequence differed from the rest of *Bradyrhizobium* strains (data not shown). On the other hand, the marker could not be amplified from other bacteria tested, except for *E. meliloti* 2011. However, given that this strain was unable to grow on BSJM medium (data not shown) this observation should be irrelevant for the direct isolation of *Bradyrhizobium*.

Table 28. *Bradyrhizobium* genomes used in the pipeline for identification of possible genus-specific markers.

Strain	Strain
<i>B. elkanii</i> USDA 76 ^T	<i>Bradyrhizobium</i> sp. EC3.3
<i>B. elkanii</i> USDA 94	<i>Bradyrhizobium</i> sp. ORS285
<i>B. elkanii</i> USDA 3254	<i>Bradyrhizobium</i> sp. ORS 375
<i>B. elkanii</i> USDA 3259	<i>Bradyrhizobium</i> sp. STM 3809
<i>B. elkanii</i> WSM1741	<i>Bradyrhizobium</i> sp. S23321
<i>B. elkanii</i> WSM2783	<i>Bradyrhizobium</i> sp. TV2a.2
<i>B. japonicum</i> 22	<i>Bradyrhizobium</i> sp. USDA 3384
<i>B. japonicum</i> USDA 4	<i>Bradyrhizobium</i> sp. WSM471
<i>B. japonicum</i> USDA 6 ^T	<i>Bradyrhizobium</i> sp. WSM1253
<i>B. japonicum</i> USDA 38	<i>Bradyrhizobium</i> sp. WSM1417
<i>B. japonicum</i> USDA 122	<i>Bradyrhizobium</i> sp. WSM1743
<i>B. japonicum</i> USDA 124	<i>Bradyrhizobium</i> sp. WSM2254
<i>B. japonicum</i> USDA 135	<i>Bradyrhizobium</i> sp. WSM2793
<i>Bradyrhizobium</i> genosp. SA-4 CB756	<i>Bradyrhizobium</i> sp. WSM3983
<i>Bradyrhizobium</i> sp. ARR65	<i>Bradyrhizobium</i> sp. WSM4349
<i>Bradyrhizobium</i> sp. CCGE-LA001	<i>Bradyrhizobium</i> sp. YR681

Figure 19. PCR amplification of the *Bradyrhizobium* marker *thrC* (296 bp). A) *Bradyrhizobium* strains available in the laboratory strain collection; and B) *Bradyrhizobium* type strain (*B. diazoefficiens* USDA 110) and other related bacteria. Rlv3841, *Rhizobium leguminosarum* bv. viciae 3841; Pst DC3000, *Pseudomonas syringae* pv. tomato DC3000.

4.3.2 *Bradyrhizobium* direct isolation method strategy

The direct isolation strategy consisted of a combination of growth on the semi-selective, optimised BSJM medium followed by PCR screening with the *Bradyrhizobium* genus marker *Threonine synthase* (*thrC*) that was developed in this work. As commented before, members of the genus *Bradyrhizobium* are characterised for having slow or extra-slow growth rates. Competition against other, faster-growing bacterial soil populations represents an important drawback for *Bradyrhizobium* isolation. This strategy was, therefore, far more challenging than those for *Neorhizobium* (see Section 4.1) or *Ensifer* (see Section 4.2). For this reason, and in contrast to them, we had to resort to, first resuspending soil samples in a *Bradyrhizobium* recovery buffer, and then to incubating them overnight in order to enhance *Bradyrhizobium* growth (Figure 3; Section 3.5.5). Other procedures were performed in a manner similar to other isolation schemes (Figure 3). The *Bradyrhizobium* isolation protocol was tested with the three soils: Tomejil, Motril and Burguillos. None of these soils had a legume cultivation history, but they differed in important soil characteristics, such as pH (Table 3). A total of five, five and fifteen independent isolations were carried out from Tomejil, Motril and Burguillos soil samples, respectively; however, successful

Bradyrhizobium isolation was only achieved from Burguillos soil samples. In both Tomejil and Motril samples, although a certain number of colonies grew on BSJM, most of them were fast-growers (see Section 4.4 below), and the few that were slow growers failed the genus-specific PCR marker amplification (data not shown). The results obtained with Burguillos soil are shown in Table 29. Forty-nine colonies were selected for PCR screening. All of these colonies were small, white-coloured and presented a slow growth on YMB agar plates. Successful PCR amplification of the *Bradyrhizobium* genus marker was achieved with only six colonies (12%). These six isolates were further characterised by PCR amplification and sequencing of the housekeeping genes 16S rDNA, *recA* and *glnII*. BLAST searches confirmed that they belonged to *Bradyrhizobium*, thus both showing that the genus marker screening was very efficient and, importantly, suggesting that *Bradyrhizobium* populations are scarce in these soils.

Table 29. Summary of direct *Bradyrhizobium* isolation results.

Soil	No. colonies tested	Total no. <i>Bradyrhizobium</i> sp. isolates	% true positives ¹
Burguillos	49	6	100

¹Assigned from the 16S rDNA, *recA* and *glnII* sequencing

Besides the ecological implications that this finding may have, it bears acutely on the importance of a robust genus marker for uncovering these scarce populations. Therefore, we were interested in the *thrC* marker conservation pattern across *Bradyrhizobium* strains, including those from the laboratory collection as well as the soil six isolates. We therefore sequenced the *thrC* amplified fragments with the same primers used for PCR screening (215 bp useable sequence) and performed a multiple alignment to ascertain their degree of conservation (Table 30). Genbank accession numbers are included in Supplemental Table S1 and the complete *thrC* alignment is shown in Supplemental Figure S2. The percentage identity among sequences varied from 87% to 100% although most values were between 90% and 100%. As expected from the method used to select it, this suggests that the amplified sequence is highly conserved and has a low variability. Altogether, we can conclude that the direct isolation strategy was able to access *Bradyrhizobium* populations in soils with no recent history of growth of *Bradyrhizobium* legume macrosymbionts, something that to our knowledge, had been rarely tested.

Table 30. Sequence conservation of the *Bradyrhizobium* genus-specific *thrC* marker (215 bp) in *Bradyrhizobium* sp. isolates.

Strain	% identity	Source of <i>thrC</i> marker sequence
<i>B. japonicum</i> USDA 6 ^T (reference)	100.0	Genome from JGI
<i>B. elkanii</i> USDA 76 ^T	91.5	Genome from JGI
<i>B. elkanii</i> USDA 94	90.6	Genome from JGI
<i>B. elkanii</i> USDA 3254	92.0	Genome from JGI
<i>B. elkanii</i> USDA 3259	92.0	Genome from JGI
<i>B. elkanii</i> WSM1741	91.5	Genome from JGI
<i>B. elkanii</i> WSM2783	91.0	Genome from JGI
<i>B. japonicum</i> 22	92.5	Genome from JGI
<i>B. japonicum</i> USDA 4	92.5	Genome from JGI
<i>B. japonicum</i> USDA 38	98.9	Genome from JGI
<i>B. japonicum</i> USDA 122	94.3	Genome from JGI
<i>B. japonicum</i> USDA 124	93.4	Genome from JGI
<i>B. japonicum</i> USDA 135	91.5	Genome from JGI
<i>Bradyrhizobium</i> genosp. SA-4 CB756	93.4	Genome from JGI
<i>Bradyrhizobium</i> sp. ARR65	86.8	Genome from JGI
<i>Bradyrhizobium</i> sp. CCGE-LA001	92.9	Genome from JGI
<i>Bradyrhizobium</i> sp. EC3.3	92.5	Genome from JGI
<i>Bradyrhizobium</i> sp. ORS285	91.5	Genome from JGI
<i>Bradyrhizobium</i> sp. ORS 375	90.6	Genome from JGI
<i>Bradyrhizobium</i> sp. STM 3809	90.6	Genome from JGI
<i>Bradyrhizobium</i> sp. S23321	93.9	Genome from JGI
<i>Bradyrhizobium</i> sp. TV2a.2	89.2	Genome from JGI
<i>Bradyrhizobium</i> sp. USDA 3384	93.4	Genome from JGI
<i>Bradyrhizobium</i> sp. WSM471	91.5	Genome from JGI
<i>Bradyrhizobium</i> sp. WSM1253	91.0	Genome from JGI
<i>Bradyrhizobium</i> sp. WSM1417	92.9	Genome from JGI
<i>Bradyrhizobium</i> sp. WSM1743	92.0	Genome from JGI
<i>Bradyrhizobium</i> sp. WSM2254	94.3	Genome from JGI

Strain	% identity	Source of <i>thrC</i> marker sequence
<i>Bradyrhizobium</i> sp. WSM2793	93.9	Genome from JGI
<i>Bradyrhizobium</i> sp. WSM3983	92.0	Genome from JGI
<i>Bradyrhizobium</i> sp. WSM4349	92.9	Genome from JGI
<i>Bradyrhizobium</i> sp. YR681	91.5	Genome from JGI
<i>B. betae</i> LMG 21987 ^T	90.6	Laboratory strain collection, this work
<i>B. diazoefficiens</i> USDA 110 ^T	92.9	Laboratory strain collection, this work
<i>B. elkanii</i> USDA 76 ^T	91.5	Laboratory strain collection, this work
<i>B. japonicum</i> USDA 6 ^T	100.0	Laboratory strain collection, this work
<i>B. liaonigense</i> LMG 18230 ^T	91.5	Laboratory strain collection, this work
<i>B. valentinum</i> LmjM3 ^T	90.1	Laboratory strain collection, this work
<i>B. yuanmingense</i> LMG 21827 ^T	93.4	Laboratory strain collection, this work
<i>Bradyrhizobium</i> sp. P.22A	93.4	Laboratory strain collection, this work
B2_2	92.0	Burguillos isolate, this work
B2_4	92.0	Burguillos isolate, this work
B2_9	92.5	Burguillos isolate, this work
B3_1	92.9	Burguillos isolate, this work
B3_2	92.9	Burguillos isolate, this work
B10_118	90.1	Burguillos isolate, this work

4.3.3 Characterisation of *Bradyrhizobium* populations

Symbiotic *Bradyrhizobium* population estimations in Burguillos soil were done by the MPN method (Toomsan *et al.*, 1984), while total *Bradyrhizobium* population was estimated as Colony Forming Units (CFU) per gram of soil using the direct isolation results. Two plant species were used for MPN estimations, *Macroptilium atropurpureum*, as it has repeatedly been shown to have a broad microsymbiont range (Thies, Bohlool and Singleton, 1991; Hafeez *et al.*, 1995) and *Lupinus albus*, as it is normally found in the Burguillos geographic zone. Table 31 shows total viable aerobic bacteria counts (CFU/g) in Burguillos soil using TSB as a rich medium and *Bradyrhizobium* estimations using the direct isolation and the MPN methods. Burguillos soil aerobic, viable bacterial counts were approximately $10^7 \cdot \text{g}^{-1}$. MPN counts of symbiotic *Bradyrhizobium* with *M. atropurpureum* and *L. albus* were very low, $10^1 \cdot \text{g}^{-1}$ in both cases. Counts using the direct isolation method were higher, increasing up to 10^3 CFU per gram of soil.

Table 31. Total viable aerobic bacteria count (CFU/g), *Bradyrhizobium* CFU/g and nodule MPN/g counts in Burguillos soil.

Soil	TSB CFU/g	Soil CFU/g	Nodule MPN/g	
			<i>M. atropurpureum</i>	<i>L. albus</i>
Burguillos	9.3×10^7 $\pm 5.7 \times 10^7$	4.0×10^3 $\pm 2.5 \times 10^3$	3×10^1 95% confidence level 0.15-11	7.4×10^1 95% confidence level 1.3-20

Table 32 summarises results of symbiotic phenotype analyses with Burguillos *Bradyrhizobium* isolates. A total of twenty-three isolates were obtained from the trap plant experiments and six directly from soil (see above). Within the soil isolates, 2 (33%) amplified the symbiotic marker *nifH* but none *nodC*. Despite this negative result, and given that *Bradyrhizobium* nodulation has been described in the absence of *nod* genes (Giraud *et al.*, 2007), nodulation experiments were carried out with *M. atropurpureum* and *L. albus*, however neither of the isolates was able to nodule either of the plants used.

Table 32. Symbiotic phenotypes Burguillos *Bradyrhizobium* isolates.

Soil	Soil isolates ¹		Nodule isolates % <i>nodC</i> ⁺ / <i>nifH</i> ⁺	
	% <i>nodC</i> ⁻ / <i>nifH</i> ⁺	% <i>nodC</i> ⁻ / <i>nifH</i> ⁻	<i>M. atropurpureum</i>	<i>L. albus</i>
Burguillos	33.3 (2 ²)	66.6 (4 ²)	100 (12 ²)	100 (11 ²)

¹No *nodC*⁺ isolates were found. ²The number of isolates is indicated in parentheses.

To further characterise these *Bradyrhizobium* isolates, they were first classified into genotypes according to their 16S rDNA full sequence, and to the partial *nodC* and *nifH* sequences of symbiotic isolates. If full or partial sequences differed in one or more nucleotides they were classified as different genotypes. According to the 16S rDNA full sequence, isolates were distributed into ten and five genotypes for nodule and soil isolates, respectively. Twenty-one genotypes were obtained for the partial *nodC* sequence of the

nodule isolates, and nine and two genotypes for the partial *nifH* sequence in nodule and soil isolates, respectively (Supplemental Table S2).

With representative strains from Burguillos soil for each genotype, and with type strains for *Bradyrhizobium* species, we performed a phylogenetic analysis based on the relevant housekeeping genes 16S rDNA, *recA* and *glnII* and also on the symbiotic genes *nodC* and *nifH*. Generally, Burguillos isolates were distributed across the genus and were present in both of the two main clusters that define the *Bradyrhizobium* genus (Menna, Barcellos and Hungria, 2009; Vinuesa *et al.*, 2005b) (Figure 20 and Figure 21). Symbiotic strains isolated from the trap plants *M. atropurpureum* and *L. albus* showed low diversity and clustered together within *B. canariense* BTA-1^T in the 16S rDNA phylogenetic tree (Figure 20), and with *B. canariense* BTA-1^T and *B. lupini* USDA 3051^T in the more resolutive phylogenetic tree derived from concatenated *recA* and *glnII* sequences (Figure 21). Soil isolates, on the other hand, were much more diverse, and were distributed across the genus. Interestingly, both soil isolates containing *nifH* (B2_4 and B3_1) each paired with a *nodC*⁻/*nifH*⁻ isolate in both trees. Generally, the *recA-glnII* concatenated phylogenetic tree showed a better resolution of the isolates, as it was expected, although only minor changes appeared. Among them, isolate B2_2, for example, switched from one cluster to the other (Figure 20 and Figure 21).

Figure 20. Phylogenetic tree of representative Burguillos soil and nodule isolates based on PCR amplified near complete 16S rDNA sequences (1,216 bp) and type strains from *Bradyrhizobium* species. Maximum likelihood trees (RAxML) were derived from ClustalW alignments. B: Burguillos soil isolates; BSN: Burguillos *Macroptilium atropurpureum* nodule isolates; BLN: Burguillos *Lupinus albus* nodule isolates. Symbiotic, non-symbiotic, and nodC⁻/nifH⁺ isolates are indicated with green, red, and purple squares, respectively. The number of strains with a given genotype is shown in parentheses. Bootstrap support (1,000 replications) for the different nodes is indicated. Bar represents the number of substitutions per base. GenBank accession numbers are shown in Supplemental Table S1.

Figure 21. Phylogenetic tree of representative Burguillos soil and nodule isolates based on PCR amplified partial concatenated *recA* (375 bp) and *glnII* (470 bp) sequences and type strains from *Bradyrhizobium* species. Maximum likelihood trees (RAxML) were derived from ClustalW alignments. B: Burguillos soil isolates; BSN: Burguillos *Macroptilium atropurpureum* nodule isolates; BLN: Burguillos *Lupinus albus* nodule isolates. Symbiotic, non-symbiotic, and nodC⁻/nifH⁺ isolates are indicated with green, red, and purple squares, respectively. The number of strains with a given genotype is shown in parentheses. Bootstrap support (1,000 replications) for the different nodes is indicated. Bar represents the number of substitutions per base. GenBank accession numbers are shown in Supplemental Table S1.

The phylogenetic tree of *nodC* sequences showed a large diversity within Burguillos nodule isolates (Figure 22). All isolates clustered together with *B. lupini* USDA 3051^T and *B. canariense* BTA-1^T –as was the case with the 16S rDNA and *recA/glnII* phylogenetic trees–, and also with *B. cytisi* CTAW11^T, *B. rifense* CTAW71^T and *B. liaoningense* 2281^T all them not too distant in the 16S rDNA and *recA/glnII* phylogenetic trees. In general terms, *M. atroporpureum* and *L. albus* isolates were distributed across all the type species mentioned above, but some differences were observed: many *M. atroporpureum* isolates clustered within *B. rifense* CTAW71^T and many *L. albus* isolates –with a larger diversity– clustered with *B. canariense* BTA-1^T. The *nifH* phylogenetic tree, that also included two soil isolates, showed a lower diversity of isolates (Figure 23). This was not surprising because: i) the amplified *nifH* fragment is much shorter; and ii) the phylogeny of *nodC* genes correlates with that of the host plant (Ueda *et al.*, 1995), whereas the variation of *nifH* genes is strongly constrained by structural requirements of the nitrogenase reductase enzyme they encode (Raymond *et al.*, 2004). Most *M. atropopureum* isolates clustered within *B. rifense* CTAW71^T and *B. cytisi* CTAW11^T. The rest of the isolates, included all *L. albus* isolates, the rest of *M. atropopureum* isolates and the two soil isolates did not cluster with any *Bradyrhizobium* sp. type strain. Furthermore, soil isolates clustered together and far from any nodule isolates, suggesting that they have an altogether different origin. In order to further characterise these isolates, we obtained draft genome sequences for the *nifH*⁺ soil isolate, B2_9 and for the corresponding *nifH* isolate B3_1 that clustered together in the 16S rDNA phylogenetic tree and another *nifH* isolate, B3_2.

Figure 22. Phylogenetic tree of representative Burguillos nodule isolates based on PCR amplified partial *nodC* (441 bp) sequences and type strains from *Bradyrhizobium* species. Maximum likelihood trees (RAxML) were derived from ClustalW alignments. BSN: Burguillos *Macroptilium atropurpureum* nodule isolates; BLN: Burguillos *Lupinus albus* nodule isolates. Nodule isolates are indicated with an orange square. The number of strains with a given genotype is shown in parentheses. Bootstrap support (1,000 replications) for the different nodes is indicated. Bar represents the number of substitutions per base. GenBank accession numbers are shown in Supplemental Table S1.

Figure 23. Phylogenetic tree of representative Burguillos soil and nodule isolates based on PCR amplified partial *nifH* (147 bp) sequences and type strains from *Bradyrhizobium* species. Maximum likelihood trees (RAxML) were derived from ClustalW alignments. B: Burguillos soil isolates; BSN: Burguillos *Macroptilium atropurpureum* nodule isolates; BLN: Burguillos *Lupinus albus* nodule isolates. Nodule isolates are indicated with an orange square; soil isolates are indicated with a purple square. The number of strains with a given genotype is shown in parentheses. Bootstrap support (1,000 replications) for the different nodes is indicated. Bar represents the number of substitutions per base. GenBank accession numbers are shown in Supplemental Table S1.

Genome features of *Bradyrhizobium* isolated genomes from Burguillos soil are included in Table 33. *Bradyrhizobium* sp. B2_9, B3_1 and B3_2 genomes vary in size and G+C(%), however, they were relatively small compared to studied *Bradyrhizobium* genomes whose size can be as large as 9 Mb (Kaneko *et al.*, 2011). Average nucleotide identity (ANI) analyses were performed with the draft genomes and *Bradyrhizobium* type strains Table 34. ANIb analyses displayed low values across the *Bradyrhizobium* genus, which manifests the diversity of the genus. *Bradyrhizobium* sp. B2_9, B3_1 and B3_2 pairwise ANI values were also low (<90%), indicating that *Bradyrhizobium* isolates from Burguillos soil were very diverse. It did not come as a surprise that although B2_9 and B3_1 clustered together in the 16S rDNA phylogenetic tree, they were so different at genomic level, since different *Bradyrhizobium* species have been shown to have identical 16S rDNA sequence (van Berkum and Fuhrmann, 2000; Willems, Coopman and Gillis, 2001; Willems, 2006). On the other hand, we also looked for genes that could provide clues about the metabolic capacities of these isolates. B2_9 presented *nif* genes but not *nod* genes, this confirmed the results obtained in the screening. It also presented *fix* and *hup* genes and genes for Rubisco. This could indicate the presence of a Calvin-Benson cycle in this strain and a possible chemolithotrophic, autotrophic lifestyle based on hydrogen, as it has been described for *B. diazoefficiens* USDA 122 (Hanus, Maier and Evans, 1979). B3_1 and B3_2 did not present either *fix* or *nod* genes in their genomes, also verifying the screening results. B3_1 did present *fix* and *hup* genes and genes for Rubisco, which again could indicate the presence of the Calvin cycle and hydrogen-dependent chemolithotrophic, autotrophic growth in B3_1. In contrast, B3_2, did not present either *hup* or *fix* genes, only Rubisco.

Table 33. Genomic features of *Bradyrhizobium* sp. B2_9, B3_1 and B3_2 genomes sequenced in this work.

Strain	Number of contigs	Largest contig (bp)	Total genome length (bp)	G+C (%)	N50
B2_9	43	642,737	6,640,846	64.97	370,113
B3_1	55	993,218	7,144,237	63.86	545, 806
B3_2	96	327,494	6,343,136	63.74	129,195

Table 34. Pairwise average nucleotide identities (ANIb) between sequenced *Bradyrhizobium* sp. isolates and *Bradyrhizobium* genomes.

Strain name	B2_9	B3_1	B3_2	<i>B. diazoefficiens</i> USDA 110 ^T	<i>B. liaoningense</i> CCNWSX0360	<i>B. japonicum</i> USDA 6 ^T	<i>B. daqingense</i> CGMCC 1.10947 ^T	<i>B. canariense</i> BTA-1 ^T	<i>B. yuanmingense</i> CCBAU 35157
B2_9	*	87.11	87.04	87.20	88.11	86.88	87.22	76.95	87.85
B3_1	86.86	*	86.24	86.05	88.24	85.74	86.45	76.41	87.47
B3_2	87.03	86.43	*	87.36	87.35	87.02	88.01	76.69	86.95
<i>B. diazoefficiens</i> USDA110 ^T	86.15	85.07	86.29	*	86.29	89.11	87.54	76.40	85.79
<i>B. liaoningense</i> CCNWSX0360	87.40	87.83	86.66	86.68	*	86.16	86.76	76.40	88.01
<i>B. japonicum</i> USDA 6 ^T	85.84	84.91	85.89	89.04	85.68	*	87.26	76.29	85.54
<i>B. daqingense</i> CGMCC 1.10947 ^T	86.70	86.10	87.39	88.18	86.92	87.94	*	76.48	86.87
<i>B. canariense</i> BTA-1 ^T	76.43	75.89	76.25	76.42	76.22	76.33	76.24	*	76.20
<i>B. yuanmingense</i> CCBAU 35157	87.52	87.29	86.51	86.66	88.29	86.36	86.96	76.59	*

4.4 *Serratia*

4.4.1 *Serratia* isolation using the BSJM semi-selective medium

In the multiple, unsuccessful attempts to isolate *Bradyrhizobium* from Tomejil soil using the isolation method consisting of a recovery buffer and the optimised BSJM semi-selective medium, we noticed that a certain type of fast-growing, white colonies invariably appeared. These grew after 24h of incubation at 28°C in YMB medium. We decided to further characterise these serendipitously isolated strains that so resisted the highly stringent BSJM medium. Towards this aim we amplified and sequenced the full 16S rDNA from these isolates. Sequences revealed that these strains belong to the genus *Serratia*. We further tested whether members of *Serratia* were also present in Motril and Burguillos soils. We were particularly interested in the presence of *Serratia* in the Burguillos soil since its pH value is much lower (pH 6.8) as compared to Tomejil (pH 8.2). A detailed description of the isolation method used for *Serratia* is shown in Figure 3 and Section 3.5.6. Generally, the isolation consisted on the selection of fast-growing colonies from the optimised semi-selective medium BSJM followed by plating onto YMB medium. Colonies that grew after 24h of incubation at 28°C were chosen and characterised by sequencing their 16S rDNA.

Five, three and one independent isolations were carried out from Tomejil, Motril and Burguillos soils, respectively, and a total of eighty-nine *Serratia* isolates were obtained in this work. Most of the isolates did not produce prodigiosin, but some prodigiosin-producing isolates were also obtained. Table 35 summarises results for *Serratia* isolations. Numbers of *Serratia* isolates differed in the Tomejil, Motril and Burguillos soils. The highest numbers of *Serratia* strains, sixty, were obtained from Tomejil soil. These represented 67% of the total colonies characterised. Lower rates were obtained for Motril and Burguillos soils. The least number of *Serratia* strains were isolated from Motril soil, seven strains, only 11% of the characterised strains; interestingly prodigiosin positive isolates were only obtained from this soil. Finally, twenty-two isolates were obtained from Burguillos soil, 19% of the total colonies characterised.

Table 35. *Serratia* strains isolated from Tomejil, Motril and Burguillos soils.

Soil	Colonies characterised	Total no. <i>Serratia</i> strains ¹
Tomejil	99	60
Motril	66	7
Burguillos	116	22

¹Assigned from the 16S rDNA sequencing

Serratia isolates were also screened for *nifH*, using the same universal primers as for Rhizobia. None of the strains was positive for *nifH* amplification.

4.4.2 Characterisation of *Serratia* populations

The above numbers of *Serratia* strains isolated from the different soils do not reflect the sizes of the respective soil populations, but rather how simple the isolation was, and this in turn was influenced by the composition of the accompanying microbiota. Estimations of *Serratia* soil populations were derived from calculation of Colony Forming Units (CFU) per gram of soil from direct isolation results. Table 36 shows these estimations for Tomejil, Motril and Burguillos soils against estimations of total viable aerobic bacteria counts using TSB as a rich medium for the corresponding soils. Viable bacterial counts were similar in all soils (ca. 10^7 - 10^8), with a high abundance of *Serratia* in Tomejil soil (ca. 10^5 CFU / g), and lower abundances for Burguillos and Motril soils (ca. 10^4 CFU/g).

Table 36. Estimation of *Serratia* soil populations.

Soil	TSB	<i>Serratia</i> CFU /g
Tomejil	$8.4 \times 10^7 \pm 9.8 \times 10^6$	$2.9 \times 10^5 \pm 1.7 \times 10^5$
Motril	$5.3 \times 10^7 \pm 1.3 \times 10^7$	$7.2 \times 10^3 \pm 5.4 \times 10^3$
Burguillos	$9.33 \times 10^7 \pm 5.7 \times 10^7$	$1.1 \times 10^4 \pm 3.1 \times 10^3$

In order to ascertain the phylogenetic and taxonomic position of *Serratia* isolates they were grouped into genotypes according to their 16S rDNA full sequence. By convention,

sequences differing in one or more nucleotides were classified as different genotypes. *Serratia* isolates were grouped in fifteen, three and six different genotypes in Tomejil, Motril and Burguillos soils, respectively (Supplemental Table S2). A representative strain from each genotype was used for subsequent phylogenetical analyses. A low resolution phylogenetic analysis was carried out with all eighty-seven isolates (two of the sequences failed and were not repeated) in order to gauge the distribution and diversity across the genus using the housekeeping 16S rDNA marker (Figure 24). Since we only used 16S rDNA, notorious for its lack of resolution at the species level (Fox, Wisotzkey and Jurtshuk, 1992), we could not assign precise phylogenetic affiliations. Despite these limitations, we were surprised by the extent of diversity in the isolates obtained: it covered most of the diversity within the genus and, presumably, many of the isolates could represent new species. Tomejil isolates clustered together with either *S. odorifera* DSM 4582^T, *S. proteamaculans* DSM 4543^T, or *S. quinivorans* DSM 4597^T, while two additional isolates, T4_8 and T4_36, clustered by themselves in a different clade. Burguillos isolates assembled into two groups close to *S. ureilytica* NiVa 51^T/*S. nematidiphila* DZ0503SBS1^T and *S. plymuthica* DSM 4540^T / *S. symbiotica* CWBI-2.3^T. All Motril soil isolates grouped with *S. marcescens*, consistent with the observation that some of these isolates produced prodigiosin.

Figure 24. Phylogenetic tree of representative Tomejl, Motril and Burguillos soil isolates based on PCR amplified near complete 16S rDNA sequences (1,296 bp) and type strains from *Serratia* species. Maximum likelihood trees (RAxML) were derived from ClustalW alignments. T: Tomejl isolates (yellow); M: Motril isolates (dark red); B: Burguillos isolates (blue). The number of strains with a given genotype is shown in parentheses. Bootstrap support (1,000 replications) for the different nodes is indicated. Bar represents the number of substitutions per base. GenBank accession numbers are shown in Supplemental Table S1.

We undertook genome sequencing to high-quality draft level of eight representative *Serratia* strains from the above groups for further studies: B1_6, B1_23, B1_42, T1_10, T1_12, T3_17, T4_5, and M3_1. Genomic features of the genomes are included in Table 37. Genomic features of the genomes show common *Serratia* genomic characteristics, that include similar genome size of around 5 Mb, except for T4_5, whose genome is slightly smaller, and G+C (%) of 56-59.

Table 37. Genomic features of *Serratia* sp. B1_6, B1_23, B1_42, T1_10, T1_12, T4_5, T3_17, M3_1 genomes sequenced in this work.

Strain	Number of contigs	Largest contig (bp)	Total genome length (bp)	G+C (%)	N50
B1_6	19	2,731,268	5,525,162	56.07	533,702
B1_23	37	1,527,521	5,173,999	59.64	456,904
B1_42	29	1,037,461	5,163,250	59.63	456,904
T1_10	28	1,179,634	5,334,657	54.96	639528
T1_12	49	743,028	5,408,046	54.84	364,364
T4_5	32	1,073,263	4,817,419	56.36	447369
T3_17	45	963,832	5,469,541	54.87	317569
M3_1	35	803,337	5,072,180	59.65	368,656

Genome-wide comparisons were carried out by calculating pairwise average nucleotide identities in our *Serratia* genomes and genomes from representatives of related *Serratia* species obtained from databanks. ANIb results are included in Table 38. The following representative isolates shown in the 16S rDNA tree for the sequenced strains are indicated in the parenthesis: B1_6 (B1_6), B1_23 (B1_115), B1_42 (B1_42), T1_10 (T1_1), T1_12 (T1_12), T3_17 (T2_10), T4_5 (T4_16), M3_1 (M3_2). ANI results corroborated the phylogenetic allocation of the isolates obtained in the 16S rDNA phylogenetic tree shown in Figure 24. Out of the representative strains sequenced, only T3_17 could represent a new species since all ANI values comparisons with T3_17 were far below < 95%.

Table 38. Pairwise average nucleotide identities (ANIb) between sequenced representative *Serratia* sp. isolates and *Serratia* genomes.

Strain name	B1_06	B1_23	B1_42	T1_10	T1_12	T3_17	T4_5	M3_1	<i>S. marcescens</i> MCB	<i>S. proteamaculans</i> 568	<i>S. liquefaciens</i> HUMV-21	<i>S. odorifera</i> DSM 4582 ^T	<i>S. ureilytica</i> Lr5/4	<i>S. plymuthica</i> V4	<i>S. plymuthica</i> 4Rx13
B1_06	*	83.46	83.46	86.29	86.26	86.31	80.68	83.56	83.52	86.41	86.22	80.67	83.49	98.72	99.44
B1_23	83.62	*	99.95	82.56	82.60	82.39	80.94	94.69	94.83	82.69	82.78	80.88	98.88	83.40	83.57
B1_42	83.69	99.98	*	82.62	82.66	82.50	80.92	94.74	94.90	82.77	82.86	80.88	98.88	83.45	83.63
T1_10	86.36	82.65	82.65	*	99.43	91.36	80.24	82.79	82.65	97.98	87.60	80.23	82.63	86.32	86.40
T1_12	86.32	82.73	82.74	99.26	*	91.27	80.22	82.90	82.72	97.79	87.53	80.25	82.73	86.22	86.31
T3_17	86.43	82.45	82.44	91.44	91.41	*	80.15	82.68	82.48	91.56	87.57	80.09	82.56	86.30	86.42
T4_5	81.03	81.32	81.32	80.62	80.63	80.49	*	81.32	81.19	80.64	80.60	99.61	81.33	81.07	81.08
M3_1	83.73	94.76	94.76	82.89	82.95	82.72	80.95	*	95.43	82.90	82.96	80.83	94.81	83.67	83.68
<i>S. marcescens</i> MCB	83.57	94.89	94.89	82.58	82.61	82.44	80.75	95.47	*	82.71	82.77	80.63	94.84	83.57	83.51
<i>S. proteamaculans</i> 568	86.61	82.84	82.84	97.93	97.93	91.50	80.42	82.95	82.95	*	87.66	80.45	82.92	86.42	86.48
<i>S. liquefaciens</i> HUMV-21	86.24	82.85	82.84	87.59	87.59	87.58	80.42	82.91	82.85	87.60	*	80.32	82.89	86.21	86.15
<i>S. odorifera</i> DSM 4582 ^T	80.96	81.09	81.09	80.51	80.52	80.27	99.54	81.04	80.89	80.50	80.45	*	81.10	80.98	80.94
<i>S. ureilytica</i> Lr5/4	83.70	98.80	98.80	82.65	82.67	82.52	81.02	94.70	94.88	82.82	82.79	80.87	*	83.40	83.70
<i>S. plymuthica</i> V4	98.68	83.21	83.21	86.28	86.26	86.25	80.63	83.45	83.40	86.31	86.16	80.60	83.26	*	98.69
<i>S. plymuthica</i> 4Rx13	99.48	83.63	83.63	86.42	86.41	86.42	80.84	83.65	83.58	86.44	86.19	80.83	83.69	98.81	*

4.4.3 Characterisation of secondary metabolism and phage sensitivity phenotypes in the *Serratia* soil isolates

It seemed clear that having such a diverse collection, certainly the largest *Serratia* soil collection, we had to characterise it further. For that, we contacted Prof. George Salmond from The University of Cambridge, who directs one of the major laboratories working on *Serratia*, particularly on the secondary metabolism and regulation, to do a research stay in his laboratory. We proceed to carry out a series of screenings with our *Serratia* strain collection looking at different phenotypic characteristics related to secondary metabolism and phage sensitivity. The following assays were done: swimming and swarming; protease activity, plant cell-wall degrading enzymes (PCWDE), such as cellulase and pectate lyase activity; quorum sensing; siderophore production; antibiotic production; antifungal production; anti-yeast activity; and phage sensitivity assays. A detailed description of these assays is included in section 3.5.7. In order to analyse and organise all the data obtained from phenotypic experiments, we performed a Principal component analysis (PCA) excluding those assays where the *Serratia* collection gave no phenotype (PCWDE and anti-yeast activity). Figure 25 shows a bidimensional PCA plot for the different phenotypes (panel A), and of *Serratia* strains against phenotypes (panel B). In the PCA plot for the different phenotypes, axis one accounted for 34% of the total variance and axis two for 24%. According to these plots, phenotypes formed three main groups. The first group consists of several physiological assays, such as antibiotic and protease production, swimming and swarming, and AHL production for quorum sensing. A second group included phage sensitivity-related phenotypes and antifungal activity. Finally, siderophore production appeared to behave differently to the other phenotypes tested. Concerning the PCA of *Serratia* strains against phenotypes, strains assembled into eleven different groups plus four individual isolates (Table 39). Tomejil isolates formed six different groups, Burguillos, four, and, finally, Motril only one, but Motril strains were also present in one Tomejil and one Burguillos group. Generally, there was variability among the isolates obtained from the different soils. This was reflected on the grouping of isolates, where each soil formed different isolate assemblies according to the phenotypes obtained. Furthermore, this grouping was also specific for soil types, since groups rarely had included representatives from different soils.

Figure 25. Principal component analysis of different phenotypes (A) and of *Serratia* strains against phenotypes (B).

Table 39. Strain composition of *Serratia* groups identified by the PCA analysis.

	Burguillos	Tomejil	Motril
Group 1	2		
Group 2		19	1
Group 3	9		2
Group 4			3
Group 5	7	1	
Group 6		8	
Group 7		7	

	Burguillos	Tomejil	Motril
Group 8		8	
Group 9	2		
Group 10		13	
Group 11		2	
T4_36		1	
T2_12		1	
B1_64	1		
B1_10	1		

Table 40 summarises the phenotypes recorded for secondary metabolism and phage sensitivity of a representative of each of the groups defined from the PCA plot. These phenotypes were scored from plate assays, many of them chromogenic or qualitative, of the type shown in Figure 26.

Looking at motility, higher swimming capacity correlated with swarming: Burguillos isolates – except for group nine (two isolates, B1_10 and B1_64)– and Motril soil isolates had both high swimming and swarming capacity. Whereas Tomejil isolates phenotypes regarding motility varied from low swimming capacity and no swarming (groups six, ten and T2_12) to high swimming and swarming (group eight and T4_36), also including high swimming and no swarming (groups seven and eleven). All isolates presented high protease activity. Siderophore production varied greatly among the strains: Tomejil strains (groups six, seven, eight, ten, T2_12 and T4_36) presented very high siderophore activity, in contrast to Burguillos and Motril isolates, which presented low or almost none (with the exception of B1_64). AHLs were detected in all Burguillos strains except for isolate B1_10. In the case of Tomejil and Motril strains, AHLs were detected in most of the strains except for group four from Motril soil, groups six and ten from Tomejil soil, T2_12 and T4_36. More AHLs were detected with the *C. violaceum* CV026 sensor strain than with the *Serratia* sp. ATCC 39006 SP19 sensor strain; furthermore, no AHLs were detected with *Serratia* sp. ATCC 39006 SP19 whenever isolates were negative for *C. violaceum* CV026. This may indicate that the AHL molecules produced by these *Serratia* isolates are C6-C8 AHLs, but not smaller since *Serratia* sp. ATCC 39006 SP19 is able to detect smaller AHLs, particularly C4 AHLs. Antibiotic production was observed exclusively in *Serratia* strains isolated from Burguillos soil, in particular group nine (formed by B1_6 and B1_28) that showed high antibiotic production. These strains also presented antifungal activity against the plant pathogen *V. dahliae*, again with the highest activity detected for group nine strains. A few Tomejil strains

(groups 6, 7, 8, 10 and T2_12) also produced antifungal activity, but none of the Motril isolates did. Generally, more *Serratia* strains exhibited antifungal vs. antibiotic activity, which may be related to biocontrol activities of these bacteria in, for example, plant environments. Regarding phage sensitivity assays, the ability of a phage to infect a host cell is directly related to the receptor it binds to; therefore, the phage sensitivity pattern also represented a relevant phenotypic characteristic. *Serratia* strains showed a broad sensitivity to phages ϕ RAY26, ϕ 3M, ϕ MAM1, and ϕ OT8, since many of the strains were infected by all the phages used. ϕ OT8 was able to infect the least number of strains, whereas ϕ RAY26, ϕ 3M, ϕ MAM1 showed the same infection pattern. The highest number of phage-resistant strains were from Tomejil soil, whereas most of the Burguillos and Motril isolates were sensitive.

Table 40. Secondary metabolism and phage sensitivity phenotypes of representative *Serratia* strains from the groups of the PCA analyses.

	Strain	Swimming (mean)	Swarming	Protease activity	Siderophore production	QS		Antibiotic production	Antifungal activity	Bacteriophage sensitivity			
						CVO26	SP19			φRAY26	φ3M	φMAM1	φOT8
Group 1	B1_98	8.7 ± 0	+	+++	<+	+	+	+	+	-	-	-	Turbid +
Group 2	T3_20	6.7 ± 0.4	-	++	+	+	-	-	-	-	-	-	=
Group 3	B1_38	8.6 ± 0.1	+	++++	<+	+	+	+	+	Turbid +	+	+	Turbid +
Group 4	M1_1	8.4 ± 0.2	+	+++	+	-	-	-	-	Turbid +	+	+	Turbid +
Group 5	B1_45	8.6 ± 0.1	+	+++	<+	+	+	+	+	Turbid +	Turbid +	+	Turbid +
Group 6	T1_2	2.8 ± 0.1	-	++	+++	-	-	-	+	Turbid +	Turbid +	-	=
Group 7	T4_8	8.4 ± 0.1	-	+++	+++	+	-	-	+	Turbid +	Turbid +	Turbid +	=
Group 8	T4_22	7.1 ± 0.6	+	+++	++	+	-	-	+	Turbid +	Turbid +	Turbid +	Turbid +
Group 9	B1_6	5.8 ± 0.3	-	+++	+	+	+	++	+++	Turbid +	Turbid +	Turbid +	=
Group 10	T1_16	2.5 ± 0	-	++	+++	-	-	-	+	Turbid +	Turbid +	Turbid +	=
Group 11	T3_12	6.8 ± 0.1	-	++	+	+	-	-	-	-	-	-	-
Individuals	T2_12	2.9 ± 0.1	-	++	++++	-	-	-	+	-	-	-	=
Individuals	T4_36	7.3 ± 0.3	+	+++	+++	-	-	-	-	-	-	-	-
Individuals	B1_10	5.0 ± 0.1	-	++	+	-	-	+	+	Turbid +	Turbid +	Turbid +	-
Individuals	B1_64	4.1 ± 0.2	-	+	++	+	+	+	-	Turbid +	+	+	Turbid +

- Negative, <+ some activity, +activity, ++ high activity, +++ very high activity and ++++ great activity. In the case of phage sensitivity assays: -, no infection, Turbid + means turbid plaques infection and +, clear infection plaques, =. CVO26: *Chromobacterium violaceum* CV026; SP19: *Serratia* sp. ATCC 39006 SP19.

Figure 26. Representative secondary metabolism plate assays and phage sensitivity assays in *Serratia*. Swimming (A1), swarming (B), protease activity (C), siderophore production (D), quorum sensing using *C. violaceum* CV026 (E1) and *Serratia* sp. ATCC 39006 SP19 (E2), antibiotic production using the *E. coli* β -lactam sensitive strain (F), antifungal production using *V. dahliae* (G), and ϕ MAM1 phage sensitivity (H).

Considering we had the genome sequences for eight *Serratia* isolates, we scanned genomes for gene clusters responsible for the biosynthesis of interesting bioactive products known in *Serratia* and that were also tested in the screening. Hence, we looked at the carbapenem biosynthetic cluster (nine genes) for antibiotic production (Thomson *et al.*, 2000), oocycin A cluster (nine genes) for antifungal activity (Matilla *et al.*, 2012b) and the cluster (four genes) for production of the mVOC sodorifen (Domik, Magnus and Piechulla, 2016b). This information coupled with the phenotypic results obtained could give insights into the bioactive products produced by these strains. Table 41 shows the presence of known antibiotic, antifungal and mVOCs biosynthetic clusters described in *Serratia* in our genomes and a comparison with the results obtained in the phenotypic screening and the antiSMASH computational analyses of the genomes (Weber *et al.*, 2015), an antibiotics and secondary metabolite analysis online tool. We found a general absence of the above biosynthetic

clusters in our *Serratia* strains. Only strain B1_6 contained the biosynthetic clusters for oocydin A production and sodorifen. Sodorifen has been shown to originate from terpene metabolism, and terpene biosynthetic capabilities were also predicted in the antiSMASH analyses for this strain. The presence of oocydin A in B1_6 could explain the results obtained against the plant pathogen *V. dahliae* in the phenotypic screening. However, the very high antifungal activity obtained could indicate that B1_6 might be producing more than one antifungal compound. This is more likely since other *Serratia* isolates positive in the *V. dahliae* assay (T1_12, B1_42, T1_10 and T4_5) do not contain the oocydin A cluster. B1_6 is phylogenetically very close to *S. plymuthica* 4Rx14 (data not shown), a strain known to produce many bioactive compounds. No presence of the carbapenem biosynthetic cluster was detected in any *Serratia* strain, even in B1_6, a strain that presented high levels of antibiotic production. This could indicate that B1_6 and other positive strains for antibiotic production (B1_23 and B1_42) could be making other β -lactam antibiotics, yet to be characterised.

Table 41. Presence of antibiotic, antifungal and mVOCs gene clusters in *Serratia* genomes and comparison with phenotypic screening results.

Strain	Antibiotic production		Antifungal		mVOCs	
	Carbapenem biosynthetic cluster	<i>E. coli</i> ESS screening	Oocydin A biosynthetic cluster	Screening against <i>V. dahliae</i>	Sodorifen biosynthetic cluster	Terpene antiSMASH prediction
B1_6	no	++	Yes	+++	yes	+
B1_23	no	+	No	-	no	-
T1_12	no	-	No	+	no	-
M3_1	no	-	No	-	no	-
B1_42	no	+	No	+	no	-
T1_10	no	-	No	+	no	-
T3_17	no	-	No	-	no	-
T4_5	no	-	No	+	no	+

- Negative, <+ some activity, +activity, ++ high activity, +++ very high activity and ++++ great activity.

4.4.4 Correlation between *Serratia* phylogeny and the phenotypes obtained

We were interested in possible correlations between the phenotypes obtained from the screening above and the phylogenetic distribution of the strains. For that purpose, we compared the strain distribution within the 16S rDNA phylogenetic tree, with the one of the

PCA plot of all *Serratia* strains against the phenotypes for each soil individually. The results of this analysis are represented in Figure 27, Figure 28 and Figure 29. In the case of Tomejil soil, generally there is a correlation between the phenotypes obtained and their phylogenetic distribution. Groups seven and eight, next to each other in the PCA plot, clustered together next to *S. odorifera* DSM 4582^T in the phylogenetic tree. Groups two and eleven, close in the PCA plot as well, also clustered together with *S. proteamaculans* DSM 4543^T. Group six correlated with the cluster of strains next to *S. quinivorans* DSM 4597^T and, finally, group ten did not correlate so well with the phylogeny of the isolates since it was subdivided in two main clusters in the phylogenetical tree, one with isolates from group six and another one next to *S. myotis* 12^T. For Motril soil, isolates were distributed in groups four and three in the PCA plot, although group three was mostly formed by Burguillos isolates. In the phylogenetic tree, all these isolates clustered together next to *S. marcescens* DSM 30121^T. In the case of Burguillos soil, isolates were distributed throughout the PCA plot; however, all isolates clustered in the phylogenetic tree next to *S. ureilytuca* NiVa 51^T, showing that, in this case, there was no correlation between the phenotypes obtained and their phylogenetic distribution.

Figure 27. Correlation between *Serratia* 16S rDNA phylogenetic distribution (left) and PCA analysis of *Serratia* strains against phenotypes (right) for Tomejil soil strains, boxed in yellow. Strain groups are numbered in yellow in both plots.

Figure 28. Correlation between *Serratia* 16S rDNA phylogenetic distribution (left) and PCA analysis of *Serratia* strains against phenotypes (right) of Motril soil strains, boxed in red. Strain groups are in red (or blue for Burguillos groups) in both plots.

DISCUSSION

Soil is one of the most complex environments in the biosphere. Besides intricate biotic interactions, soil organisms face temporally- and physically-changing physico-chemical conditions that include a highly diverse matrix structure. All of the above is especially true for microorganisms, given that their small size makes them more sensitive to soil structural effects, both at the micro and macro levels. Microbial soil populations are large and diverse (Fierer and Jackson, 2006; Vogel *et al.*, 2009; Gilbert, Jansson and Knight, 2014; Alivisatos *et al.*, 2015). Furthermore, a large fraction of the microorganisms present in a given soil cannot be readily cultivated (Pham and Kim, 2012), and all this has traditionally compounded difficulties associated to the study of soil microbial ecology. It is only in recent years, with the advent of metagenomics (Gilbert, Jansson and Knight, 2014; Alivisatos *et al.*, 2015), that significant inroads into this field have been made.

Culture independent methods such as soil and plant shotgun metagenomics and 16S rDNA amplicon sequencing projects have revolutionised the study of environmental microbiology and allowed, for the first time, access to a whole new soil microbial diversity including the major unculturable fraction, what has come to be known as the “unseen majority.” However, even metagenomic studies present limitations, many of them related to either depth of coverage or, on the contrary, to analysis of massive data. In fact, soil metagenomics has limited usefulness for low abundance bacteria (10^1 - 10^4 UCF/g), since sequencing depths of up to 1 Tb would be needed in order to obtain 100 Mb of sequence of such a subpopulation, and even that level of coverage would result in just a 15x coverage of a single genome, insufficient to study these populations. In fact, it has been estimated that a sequencing depth of 50 Tb is required to sample 1 g of soil adequately (Gans, Wolinsky and Dunbar, 2005). To put all the above in perspective, an insufficient 1 Tb sequencing depth would be ten-fold higher than the largest metagenomics datasets available today, (metagenome IDs: 3300002568, 3300000917, or 3300001305, in the JGI IMG/M database; <https://img.jgi.doe.gov/cgi-bin/m/main.cgi>). In view of these limitations, other culture-independent methods, such as direct gene amplification from soil DNA, have been used (Zézé, Mutch and Young, 2001; Sarita *et al.*, 2005). At any rate, attempts to approach the genomic structure of minor components of soil populations with available metagenomics and 16S rDNA amplicon sequencing methodologies are clearly unsatisfactory. Among such minor components of the soil microbiota, Rhizobia are probably the most relevant group in terms of agricultural importance. For the purpose of characterisation of symbiotic subpopulations, researchers have traditionally taken advantage of the strong selectivity and capacity for enrichment of legume plant hosts, and there is a plethora of available publications where symbiotic Rhizobia are selected from soil and trapped by their host.

Conversely, there is a paucity of studies on non-symbiotic Rhizobia populations in soil. It is believed that they exist, and that they contribute to the genetic pool of the different clades, although this has been effectively shown in just a few examples (Segovia *et al.*, 1991; Laguerre, Bardin and Amarger, 1993; Sullivan *et al.*, 1996; Pongsilp *et al.*, 2002).

In view of this situation, the main objective of this work has centred on the establishment of methodologies that would allow the isolation and quantification of soil bacteria that compose such a minor, neglected component of soil. The isolation of these novel bacteria has allowed access to their phenotypic and genomic characteristics, some of which could have interest for the characterisation of functions relevant for sustainable agriculture

5.1 Non-symbiotic rhizobial populations in soil

Direct isolation of Rhizobia from soil without resorting to trapping them by means of their legume host is complicated by the fact that rhizobial populations in soil can be small (Singleton and Tavares, 1986) and because of the paucity of selective characters that can be used. This is especially true for non-symbiotic variants lacking any symbiotic marker. As a result, very few studies have aimed at isolating non-symbiotic Rhizobia from soil. Groundbreaking work by Sullivan and Ronson with *Mesorhizobium loti* (Sullivan *et al.*, 1995), did not only reveal the existence of an abundant population of non-symbiotic variants in the soil, but also showed that the genetic determinants for symbiosis could be readily transferred to these variants within the soil. Some of these non-symbiotic variants were further characterised and suggested to be representatives of species hitherto undescribed. Using a semi-selective culture medium, Tong and Sadowsky (1994) were able to enrich in soybean-specific Rhizobia of the species *Bradyrhizobium japonicum* and *B. elkanii*. Using this medium, the same group later described that about half of the Bradyrhizobia isolated directly from Thai soils were non-symbiotic, although they did not characterise them further (Pongsilp *et al.*, 2002). Louvrier, Laguerre and Amarger (1995) were also able to enrich members of the genus *Rhizobium* using a specific medium. Although they focused on symbiotic isolates (Louvrier, Laguerre and Amarger, 1996) they also isolated a fraction of non-symbiotic Rhizobia (Laguerre, pers. comm, 2011). On the other hand, a few attempts have been made to study free-living Rhizobia. Schmidt, Bankole, and Bohlool (1968) used fluorescent-antibodies (FA) specific of *Rhizobium japonicum*, however non-specific adsorption in soil was a problem, and another group developed a real-time PCR assay for

the detection of free-living *Rhizobium leguminosarum* using specific 16S rDNA and *nodD* primers (Macdonald *et al.*, 2011).

In order to describe the rhizobial soil populations in this work, soil selection was determinant, as we were interested in native rhizobial populations, particularly of non-legumes. The principal characteristic we were looking for was the cultivation history of the soils. No legumes had been grown in either Tomejil (where winter wheat is usually grown, with occasional cultivation of cardoon), Motril (where sugarcane is routinely grown), or – especially– in Burguillos (a pristine “dehesa”) soils in recorded times. These soils also differed in their pH (Table 3) and, given that pH has been shown to be the most important factor shaping bacterial communities in soil (Fierer and Jackson, 2006; Rousk *et al.*, 2010; Liu *et al.*, 2014), it was also reasonable to study soils of different pH. Within Rhizobia, *Ensifer* sp., specially *E. meliloti*, have been reported to be sensitive to acidic soils (Rice, Penney and Nyborg, 1977; Brockwell, Pilka and Holliday, 1991), and *Bradyrhizobium* spp. are generally more acid-tolerant than *Rhizobium* spp. for example (Brockwell, Pilka and Holliday, 1991; Sadowsky, 1998). Differences in pH might explain the results we obtained for *Bradyrhizobium* and *Ensifer* population distribution: *Bradyrhizobium* was successfully isolated only from an acidic soil (Burguillos, pH 6.2) whereas, in contrast, *Ensifer* was only isolated from alkaline soils (Tomejil, pH 8.1, and Motril, pH 8.45).

Our group had previously used a *Rhizobium* semi-selective medium (Louvrier, Laguerre and Amarger, 1995), together with the *fnrN* gene marker, to isolate Rhizobia from soil (Jorrín, 2016; Jorrín and Imperial, unpublished data) in order to study, at the genomic level, how specific legume hosts select specific *R. leguminosarum* genotypes among those present in the soil. This phenomenon had been described by the Amarger-Laguerre (Laguerre *et al.*, 2003) and the Young (Young and Wexler, 1988) research groups. During that study, we had been able to isolate a number of non-symbiotic members of the genus *Rhizobium* (Jorrín, 2016; Jorrín and Imperial, unpublished data). Therefore, it was surprising that the same enrichment methodology resulted in the isolation of a large (over 50%) proportion of non-symbiotic members of the genus *Neorhizobium* from soil samples from the IFAPA Tomejil Experimental Station in Carmona (Seville, Spain), first because non-symbiotic *Neorhizobium* had not been previously isolated, and second because our genetic screening with *fnrN* –a *Rhizobium/Agrobacterium* specific marker– should have left them behind (although the different size of the PCR amplified bands was already an indication that these were atypical members of the genus *Rhizobium* at best). The serendipitous reasons why these particular *Neorhizobium* were chosen as putative *Rhizobium* have been presented in section 4.1.

Clearly, they grow well on MNBP semi-selective medium, with cultural characteristics similar to those of *Rhizobium*. However, it is possible that, among the two hundred and forty-three colonies that tested negative for *fnrN* amplification, other non-symbiotic *Neorhizobium* strains that do not have a high enough conservation of the poly (3-hydroxybutyrate) depolymerase region exist in the Tomejil soil. At any rate, *Neorhizobium* appears to be at least as abundant in the particular sampled soil as *Rhizobium*, and given the high rates of horizontal gene transfer that can take place in soil (Sullivan *et al.*, 2002), they may become relevant in the establishment of symbioses with native legumes after receiving the appropriate symbiotic genes. Acquisition of these genes might be facilitated by the fact that *N. galegae* has been shown to harbour symbiotic genes on a megaplasmid (Radeva *et al.*, 2001; Österman *et al.*, 2014). As shown in section 4.1.2, all the sequenced Tomejil strains contain a set of *repABC* genes characteristic of plasmids from the Rhizobiales (Cevallos, Cervantes-Rivera and Gutierrez-Rios, 2008; Pinto, Pappas and Winans, 2012) in large contigs, suggesting that these strains also harbour a megaplasmid and are, thus, probably able to receive similar, large plasmids containing symbiotic genes. Further phylogenetical analyses of the non-symbiotic *Neorhizobium* isolates from Tomejil soil led to the description of the first non-symbiotic *Neorhizobium* species, *N. tomejilense* sp. nov. Although we only described one species, genotypic and genomic characterisation of these isolates suggests that there might be more than one species within *Neorhizobium* (Table 14). A final consideration is whether the existence of populations of non-symbiotic *Neorhizobium* in soils reflects a normal situation in agricultural soils. To answer this question would require a large screening that is beyond the scope of this work. However, our preliminary studies suggest that in at least two other soils from Southern Spain that we have tested (Motril and Burguillos), members of *Neorhizobium* are not present in detectable numbers, and that the predominant isolates resulting from our semi-selective screening are *Rhizobium* spp. (Soenens and Imperial, unpublished results). This would then shift the question to why the Tomejil soil harbours a sizeable *Neorhizobium* population, a question whose answer would require a better understanding of the ecology of this group in soil.

The success in choosing appropriate molecular markers for screening of candidate isolates was determinant in the isolation of soil Rhizobia, even more so when working with soils where Rhizobia are relatively rare. In the case of *Ensifer* and *Bradyrhizobium*, and due to the lack of selective characters that could be used to directly select rhizobial populations without resorting to their symbiotic capacity, we undertook to develop specific genus markers for both *Ensifer* and *Bradyrhizobium*. To that end, we used available genome sequences and adapted the methodology described by Wu, Jospin and Eisen (2013). To our knowledge this

is the first time that this method has been used to develop a single-copy genus maker. Both markers, *Ensifer* sp. *asmA* and *Bradyrhizobium* sp. *thrC* proved to be useful and convenient markers, as they allowed quick identification of isolates by PCR screening with negligible rates of false positives (either null or ca. 12.5% on average; see Table 19 and Table 29). Furthermore, the chosen markers were conserved enough throughout *Ensifer* and *Bradyrhizobium* genera that we were able to use them to successfully identify a large subset of the known species described for each genus (Figure 6 and Figure 19).

Together with the use of semi-selective media, we have provided a successful strategy that enables the isolation and quantification of novel rhizobial populations from soil without recourse to legume plant assays, in particular the non-symbiotic populations, not previously studied beyond the mere acknowledgement of its existence. Furthermore, this simple methodology that combines cultural enrichment characteristics together with differentiating genetic properties, can be extrapolated for the development of similar strategies to study other specific bacterial groups. In general terms, we found non-symbiotic populations to be much larger and diverse than symbiotic populations for both *Ensifer* and *Bradyrhizobium*. In the case of *Neorhizobium*, only non-symbiotic populations were isolated, and these were also highly diverse. In Tomejil soil, the *Neorhizobium* sp. population was estimated at ca. $4.4 \times 10^4 \pm 2.0 \times 10^4$ CFU/g of soil. In Burguillos soil, non-symbiotic *Bradyrhizobium* populations were an order of magnitude smaller (ca. $4 \times 10^3 \pm 2.5$ CFU/g), but still two orders of magnitude above estimations for symbiotic *Bradyrhizobium* (Table 31). This difference was even larger (three orders of magnitude) for non-symbiotic vs. symbiotic *Ensifer* from Tomejil and Motril soils (Table 21). As shown in Table 1, non-symbiotic Rhizobia had been only sporadically identified previously; after this work, despite its limitations in scope and range, it is clear that non-symbiotic rhizobial populations can be abundant and diverse in soils, even when no legumes had been knowingly cultivated in recent times. In fact, our results suggest that these sizeable, non-symbiotic rhizobial communities might be common in soils where no legumes have been planted. At any rate, the non-symbiotic populations accessed in this work represent novel reservoirs of *Neorhizobium*, *Ensifer* and *Bradyrhizobium* genetic diversity, not accessible before, that could represent in many cases new species.

Despite its limitations, our study shows that rhizobial population distribution varied widely among soils. *Neorhizobium* and *Bradyrhizobium* were only isolated from one of the three soils sampled, Tomejil and Burguillos respectively. This can be taken as indication that these populations are very specific of the soil/cultivation history. *Ensifer*, on the other hand, was isolated from two of the soils (Tomejil and Motril) and were not detected in Burguillos soil.

Both Tomejil and Motril soils had similar, basic pH, and the *Ensifer* isolates were also similar. Contrary to the situation found with *Rhizobium* in a soil from the Bourgogne region in France (Jorrín, 2016; Jorin and Imperial, unpublished results), direct isolation failed to identify symbiotic Rhizobia (with the possible exception of *Bradyrhizobium* from Burguillos, where two *nodC*/*nifH*⁺ strains were isolated). This could be due either to very low abundance of symbiotic Rhizobia in the soils studied, or –despite our efforts to choose plant hosts with little specificity– to the inability of the chosen host plant(s) to trap them. For instance, alfalfa plants were used with *Ensifer*, and any symbiotic *Ensifer* specific for plants other than *M. sativa* would be missed. Furthermore, MPN has a lower detection limit of 10 cells per gram of soil (Brockwell, 1982), this is more sensitive than our direct soil isolation method thanks to the highly specific nature of the Rhizobia-plant interaction. In the direct isolation method, not only rhizobial abundance comes into play, but also the size and nature of the whole bacterial community present in a particular soil. Despite these caveats, the direct isolation method was powerful enough to isolate an *Ensifer* non-symbiotic strain (M4_45) closely related at the phylogenetic level to an *Ensifer* nodule isolate (MMN_5) from *M. sativa*, as well as two *Bradyrhizobium* soil isolates, one containing *nif* genes (B2_9), and one non-symbiotic (B3_1) that presented the same 16S rDNA genotype. The genomes of these four isolates were sequenced to draft level and the absence of symbiosis-related genes (*nod* and *nif* genes) in the genome sequence of the non-symbiotic strains confirmed that the PCR screening worked indeed as predicted. Only the work with the *Ensifer* isolates was taken further within the context of this Dissertation. Both whole genome analyses and alignment in comparison to the reference genome *E. meliloti* 1021 (Figure 17, Figure 18, Table 24 and Table 25), showed important reorganisations of pSyms in the symbiotic MMN_5 and non-symbiotic M4_45 isolates, both originating from the same soil subpopulation. In particular, “pSymA” (or more properly, those genome contigs recruited by the reference strain pSymA) exhibited a greater reorganisation, corresponding to complete absence (either loss in the non-symbiotic, or gain in the symbiotic strains) of the symbiotic genes. These results correlate with earlier studies showing that rhizobial symbiotic and non-symbiotic plasmids have a mosaic structure (Zhang, Kosier and Priefer, 2001; González *et al.*, 2003; Giuntini *et al.*, 2005). Although comparisons between rhizobial free-living and nodulation states have been carried out before (Becker *et al.*, 2004; Queiroux *et al.*, 2012; Gomes *et al.*, 2014), they mainly dealt with differences in expression for genes within a given genomic complement. Our work suggests a novel, comparative genomic strategy that could give new insights into the genetics and molecular mechanisms of symbiosis, particularly in the adaptation and evolution of native *Ensifer* sp. soil populations.

The results obtained in this work, make it clear that non-symbiotic Rhizobia populations in soil do not arise as a consequence of an anecdotal or accidental plasmid loss within symbiotic rhizobial strains. In fact, non-symbiotic *Neorhizobium*, *Ensifer* and *Bradyrhizobium* populations are large and diverse bacterial communities widely present in soils. Questions related to the relative proportions of both symbiotic and non-symbiotic populations, to their interrelations and stability over time are widely open, as is the effect on these parameters of the presence of the plant host.

5.2 *Serratia* populations in soil

A large and diverse soil *Serratia* population was uncovered serendipitously during our efforts to directly isolate *Bradyrhizobium* from soil. In this work, we were able to develop a protocol allowing, for the first time, direct *Serratia* isolation from soil. *Serratia* isolation was extremely convenient due to the nature of these bacteria, as their fast growth rate, characteristic of bacteria belonging to the Enterobacteriaceae family, is a mayor advantage for their isolation, particularly in complex environments such as soil (Hibbing *et al.*, 2010). We were able to isolate *Serratia* from all soils tested (Tomejil, Motril and Burguillos), and this is an indication that *Serratia* is generally present in soils. More interestingly, a wide diversity was present, and populations were not dominated by a singles species. *Serratia* populations differed in both abundance and diversity among soils, ranging from 10^4 to 10^5 CFU per gram of soil. Among the isolates there were species that could be clearly expected to be found in soils, such as *S. proteomaculans* or *S. plymuthica*, since these have been described as plant-related species; however the diversity obtained was beyond these species, and although *Serratia* has been previously isolated from many different niches, it was highly surprising that we obtained representatives of most of the *Serratia* species described (

Figure 24). It is possible that many of our isolates might represent novel species. Although further work is needed, the pairwise ANI values obtained for T3_17 (Table 38) and the 16S rDNA distribution of the corresponding genotype (T20_10) in the 16S rDNA phylogenetic tree (

Figure 24), suggest the existence of at least one new species among our isolates. This genotype also represents the most abundance group of Tomejil soil isolates (Supplemental Table S2). In particular, there were drastic differences in the *Serratia* populations isolated from Tomejil soil versus those from Burguillos and Motril. These differences cannot be attributed to soil pH, a major shaping factor of bacterial soil microbiota (Lauber *et al.*, 2009; Tripathi *et al.*, 2018) in view of the similarities between Tomejil and Motril soils (Table 3). A

clearly differential physico-chemical characteristic between Tomejil and Motril is salinity, and this could perhaps explain the isolation of *S. marcescens* exclusively from Motril soil. Distribution differences with Burguillos soil on the other hand, could then be related to soil pH (6.1 for Burguillos and 8.1 for Tomejil). Regardless of the reasons, the existence of such a diverse collection of *Serratia* isolates made us decide to try to further characterise it.

The genus *Serratia* is characterised by a rich secondary metabolism and, therefore, their phenotypic characterisation was based on secondary metabolism-related phenotypes in addition to a set of phage sensitivity assays. We classified all these phenotypes by means of Principal component analyses, where we grouped the isolates individually based on their phenotypic characteristics. PCA is a multi-dimensional classification technique for collapsing the variance of highly-dimensional datasets into a small number of components, thus increasing interpretability while at the same time minimizing information loss (Jolliffe and Cadima, 2016). In our particular case, PCA worked ideally for our dataset, since it allowed a meaningful classification into biologically sound clusters. In the PCA classification against their phenotypes, *Serratia* isolates clustered mainly based on the soil they were isolated from, although isolates from the same soil clustered in different groups. This observation could be explained by two simultaneous properties of populations, first, by a high soil specificity, and second, by a broad diversity within each of the soil populations. The classification afforded by PCA analyses allowed us to compare phenotypic groups with strain phylogeny. Above specific considerations (see section 1214.4.4), the variability observed in the phylogenetic distribution of the isolates was also observed at the phenotypic level.

Soil is a traditional reservoir for new bioactive compound discovery and, as expected, many of our isolates presented interesting traits, such as antibiotic and antifungal activity. Antibiotic production was limited to Burguillos soil strains and further work on the characterisation of these compounds would probably be worthwhile. *Serratia* strains are known to produce carbapenems, these are β -lactam antibiotics (Thomson *et al.*, 2000). Given that we used a β -lactam sensitive strain (*E. coli* ESS) we could expect that the antibiotics produced would be β -lactams; however, the *E. coli* ESS indicator strain is not only restricted to these (Nishida *et al.*, 1977). We searched in our *Serratia* genomes for *car* genes, responsible for β -lactam production and there were no such genes present in the genomes. Although BLAST similarity search was performed with *Serratia car* genes, the fact that no similar genes were found in our genomes, does not absolutely rule out that they might have other, distinct carbapenem genes. Therefore, further characterisation of the antibiotics produced should be

carried out, at least from those isolates with higher antibiotic activity, such as B1_6 and B1_28 (Table 39). Chemical strategies for such a characterisation are classical nowadays (Talarico and Dobrogosz, 1989; Dizman, Elasri and Mathias, 2005; Lambert *et al.*, 2007, Niveshika *et al.*, 2016). Antifungal activity, in turn, was widely distributed among *Serratia* isolates, this is consistent with the numerous examples of *Serratia* strains used or described as biocontrol agents (Berg, 2000; Kurze *et al.*, 2001; Grosch *et al.*, 2005). A search for genes encoding synthesis of oocydin A –an antifungal compound described in *S. plymuthica*– identified the biosynthetic cluster in one of our sequenced *Serratia* genomes, that of strain B1_6. This strain showed high antifungal production against *V. dahliae* in plate assays. Such a high activity could result from production of oocydin A or by a combination of this and other antifungal compounds. A gene cluster for synthesis of sodorifen, an mVOC described in *Serratia* whose biological role has not been described so far but contains a unique structure, was also present in strain B1_6. This strain therefore, represents a promising candidate for further characterisation of bioactive compounds. The fact that many of the genomes did not contain known *Serratia* biosynthetic clusters for bioactive compounds while exhibiting bioactive phenotypes, also presents an opportunity for the description of putatively new bioactive products in this genus. Siderophore production was also analysed in the collection of *Serratia* isolates. Tomejil strains presented a consistently high siderophore production, and we reasoned that this phenotype could be related to specific levels –low– or forms –insoluble– of iron in that soil. For that purpose, we sought help from one of the foremost groups in iron soil chemistry (Torrent, Schwertmann and Schulze, 1980), who analysed our soils for different iron forms (Table 3). Tomejil soil clearly presented a lower iron availability, and this correlated well with the enhanced siderophore production by its *Serratia* isolates. Finally, in the phage sensitivity assay, two kinds of plaques were observed in *Serratia* top lawn infected by the phage, clear and turbid plaques. This is consistent with the lytic and lysogenic phage life cycles, respectively. Generally, phage sensitivity appeared to be broad. This is highly interesting, not only because some of the phages could be used as genetic tools with these *Serratia* isolates, but also because it allowed us to estimate the host range distribution of these phages in such a diverse collection of *Serratia* species.

Nitrogen-fixing *Serratia* isolates have been identified (Ruppel, 1991; Gyaneshwar *et al.*, 2001), these were able to fix nitrogen endophytically in wheat and rice. For this reason, we decided to screen for *nifH* using the same universal primers as for Rhizobia, with negative results in all the strains tested. This was later on confirmed with the genome sequences obtained for some of the strains.

5.3 Rhizobia and *Serratia*, two different strategies for life in the soils and potential PGPRs

Our work on *Serratia* was focused on their secondary metabolism, particularly on their ability to produce interesting natural compounds, predominantly with antibacterial and antifungal properties, and on how this metabolism could be related with quorum sensing. In the case of Rhizobia, work was focused on the development of adequate methodologies that would allow access to soil native populations and their genomic characterisation, without recourse to plant traps. In both cases we were able to uncover the existence of established soil populations of these bacteria. The significance and role of these populations, especially regarding interactions with plants, still needs to be determined. It is possible that some of these bacteria could have an alternative lifestyle as PGPRs in the right environment or with the adequate host, and this hypothesis seems appropriate since Rhizobia and *Serratia* have been sporadically identified as PGPRs. Throughout this work, we have developed resources and a knowledge base that could allow us to answer this question in the future. We foresee a two-pronged approach, where screenings of well-known PGPR traits –such as phosphorus solubilisation strategies, siderophore production, phytohormone production (IAA and ACC deaminase in particular, since these have been the most extensively studied) and volatile organic compounds (VOCs)– would be accompanied by genomic studies of their possible genetic determinants, taking advantage of our collection of symbiotic and non-symbiotic Rhizobia, in addition to the *Serratia* genomes, to be continued by direct plant inoculation tests. Furthermore, the biocontrol potential of the *Serratia* strains should be further studied in view of our preliminary results, that point to an aggressive lifestyle of *Serratia* populations in soil, and that may have biotechnological applications in directed biocontrol.

At any rate, it is clear that *Serratia* and Rhizobia have drastic lifestyle differences in the soil that they follow in order to adapt to oligotrophic and copiotrophic conditions. These differences are also apparent in the PGPR properties they usually display (Table 2), although these may be not as extreme as originally thought, in view of the observed ability of many of the soil *Serratia* to thrive in the demanding semi-selective media we used for their isolation. In PGPR studies so far, Rhizobia are usually associated with the production of phytohormones or resistance to metals, whereas *Serratia*, on the other hand, are usually associated with biocontrol activities of plant pathogens. It has been traditionally assumed that symbiotic Rhizobia just sit in soil waiting for a host, where they will benefit from the secure and protected niche that the plant provides thus increasing their maintenance and survival in soil once they are released. Non-symbiotic Rhizobia, which are showing up

consistently in the endophytic fraction of the microbiota of many plants, not necessarily legumes (Lundberg *et al.*, 2012; Peiffer *et al.*, 2013; Shakya *et al.*, 2013; Chaparro, Badri and Vivanco, 2014; Ofek-Lalzar *et al.*, 2014; de Souza *et al.*, 2016; Castañeda and Barbosa, 2017) could probably follow similar strategies in order to preserve themselves in the secure and protected environment provided by the plant. This does not appear to be a widespread strategy with *Serratia* which, rather than seek a protective environment, aggressively defend their territory. For example, they have been shown to encode type VI secretion system (T6SS) that target bacterial cells, and that can be used as antibacterial T6SSs against competitors (Murdoch *et al.*, 2011; English *et al.*, 2012; Fritsch *et al.*, 2013). T6SSs are widespread among Gram-negative bacteria and exhibit an intriguing versatility. They have also been identified in Rhizobia, where they appear to facilitate the interaction with their symbiotic host, rather than being a pathogenicity determinant (Nelson and Sadowsky, 2015). A further consideration within *Serratia* is whether regulations would permit their use in agriculture, given that the genus contains some opportunistic human pathogenic species (*S. marcescens*). Despite these caveats, *S. plymuthica* for example, has been used as a biocontrol agent for fungal diseases in plants, and nowadays this species is classified within risk group 1 by the DSMZ (German Collection of Micro-organisms and Cell Cultures), to indicate that it does not pose a threat to human health.

A final consideration regarding potential uses of Rhizobia as PGPR is that the anecdotically low numbers of non-symbiotic *Neorhizobium*, *Ensifer* and *Bradyrhizobium* described to date, together with the lack of access to these populations, has determined that most of the rhizobial PGPRs described until now are symbiotic (Table 2). We cannot, therefore, underestimate the potential of the rhizobial populations isolated in this work as PGPRs.

CONCLUSIONS

1. This work has allowed, for the first time, a comprehensive study of *Neorhizobium*, *Ensifer* and *Bradyrhizobium* non-symbiotic populations and *Serratia* populations in soils with no previous legume cultivation history. The combination of semi-selective media followed by a PCR screening using a genus or species marker, has been shown to be an adequate methodology to characterise soil populations for low abundance bacteria, such as Rhizobia, or of unknown abundance, as in the case of *Serratia*.
2. In all cases, non-symbiotic rhizobial populations were larger and more diverse than the symbiotic ones, thus revealing that soils are a repository of large rhizobial diversity in the form of non-symbiotic variants that have traditionally escaped characterisation.
3. Rhizobia were not equally isolated from all soils studied. The slow-growing *Bradyrhizobium* were only isolated from the acidic Burguillos soil. In contrast, *Ensifer* and *Neorhizobium* populations were only isolated from alkaline soils: Tomejil, in the case of *Neorhizobium*, and Tomejil and Motril, in the case of *Ensifer*. This indicates clear differences in rhizobial distribution according to soil pH.
4. *Serratia* were widely present in soil; in fact, levels of diversity equivalent to those previously described within the genus were obtained from the soils studied in this work. Abundance and diversity of isolates obtained varied among the soils studied.
5. Isolation and identification of non-symbiotic members of the genus *Neorhizobium* from soil was achieved for the first time. Genotypic and genomic characterisation of these isolates suggests that, among them, representatives of one, or perhaps two, new genomospecies within *Neorhizobium* were present. Further phylogenetic characterisation of these novel non-symbiotic *Neorhizobium* isolates led to the description of the first non-symbiotic *Neorhizobium* species, *N. tomejilense* sp. nov.
6. The *Ensifer* sp. direct isolation method allowed isolation of a non-symbiotic strain phylogenetically close to a nodule isolate from the same soil. Genomic comparison showed pSyms reorganisation in these two strains, particularly of pSymA, which harbours the symbiotic genes. These studies could provide insights into the adaptation and evolution of native *Ensifer* populations in soils.

7. The *Bradyrhizobium* population obtained from Burguillos soil was composed of symbiotic (*nodC*⁺/*nifH*⁺), non-symbiotic (*nodC*⁻/*nifH*⁻) and nitrogen-fixing (*nodC*⁻/*nifH*⁺) representatives, reflecting the genomic diversity of this genus. Genomic analyses of some of the soil isolates revealed that these were not coincident with any described species of the genus, indicating that they could represent new species.
8. Phenotypic characterisation of the *Serratia* collection showed that they produce a full range of interesting secondary metabolites such as antibiotics and antifungals. Further characterisation of these compounds, coupled with genomic analysis of the strains could reveal novel bioactive products.
9. Particularly, *Serratia* strain B1_6 represents an exceptional candidate for biocontrol studies, since phenotypic and genomic analyses showed its ability to produce antibiotics (whose nature is still to be determined), antifungals, and mVOCs such as oocydin A and Sodorifen.

BIBLIOGRAPHY

- Abdi H, and Williams LJ. (2010). Principal component analysis. *Wiley Interdiscip Rev: Comput Stat*, 2:433-459.
- Abedon ST, Kuhl SJ, Blasdel BG, and Kutter EM. (2011). Phage treatment of human infections. *Bacteriophage*, 1:66-85.
- Adam E, Müller H, Erlacher A, and Berg G. (2016). Complete genome sequences of the *Serratia plymuthica* strains 3Rp8 and 3Re4-18, two rhizosphere bacteria with antagonistic activity towards fungal phytopathogens and plant growth promoting abilities. *Stand Genomic Sci*, 11:61.
- Albrecht-Gary AM, and Crumbliss AL. (1998). Coordination chemistry of siderophores: thermodynamics and kinetics of iron chelation and release. *Met Ions Biol Syst*, 35:239-327.
- Alivisatos AP, Blaser MJ, Brodie EL, Chun M, Dangl JL, Donohue TJ, Dorrestein PC, Gilbert JA, Green JL, Jansson JK, Knight R, Maxon ME, McFall-Ngai MJ, Miller JF, Pollard KS, Ruby EG, Taha SA, and Consortium UMI. (2015). MICROBIOME. A unified initiative to harness Earth's microbiomes. *Science*, 350:507-508.
- Alström S. (2001). Characteristics of Bacteria from Oilseed Rape in Relation to their Biocontrol Activity against *Verticillium dahliae*. *J Phytopathol* 149:57-64.
- Altschul SF, Gish W, Miller W, Myers EW, and Lipman DJ. (1990). Basic local alignment search tool. *J Mol Biol*, 215:403-410.
- Ando S, Goto M, Meunchang S, Thongra-ar P, Fujiwara T, Hayashi H, and Yoneyama T. (2005). Detection of *nifH* Sequences in Sugarcane (*Saccharum officinarum* L.) and Pineapple (*Ananas comosus* [L.] Merr.). *Soil Sci Plant Nutr*, 51:303-308.
- Andrews M, and Andrews ME. (2017). Specificity in Legume-Rhizobia Symbioses. *Int J Mol Sci*, 18 (4).
- Angerer A, Klupp B, and Braun V. (1992). Iron transport systems of *Serratia marcescens*. *J Bacteriol*, 174:1378-1387.
- Armanhi JSL, de Souza RSC, Damasceno NdB, de Araújo LM, Imperial J, and Arruda P. (2017). A Community-Based Culture Collection for Targeting Novel Plant Growth-Promoting Bacteria from the Sugarcane Microbiome. *Front Plant Sci*, 8:2191.
- Atlas RM. (1996). Handbook of Microbiological Media: Second Edition: *CRC-Press*. 1712 pp.
- Babic KH, Schauss K, Hai B, Sikora S, Redzepovic S, Radl V, and Schlöter M. (2008). Influence of different *Sinorhizobium meliloti* inocula on abundance of genes involved in nitrogen transformations in the rhizosphere of alfalfa (*Medicago sativa* L.). *Environ Microbiol*, 10: 2922-2930.
- Bachmann B. (1996). Derivations and genotypes of some mutant derivatives of *Escherichia coli* K12. Neidhardt, F.C. (ed.). Washington, DC: *ASM Press*, pp. 2460-2488.
- Badri Y, Zribi K, Badri M, Huguet T, van Berkum P, and Aouani ME. (2007). Comparison of rhizobia that nodulate *Medicago laciniata* and *Medicago truncatula* present in a single Tunisian arid soil. *Can J Microbiol*, 53:277-283.

- Bai Y, Souleimanov A, and Smith DL. (2002). An inducible activator produced by a *Serratia proteamaculans* strain and its soybean growth-promoting activity under greenhouse conditions. *J Exp Bot*, 53:1495-1502.
- Bailly X, Olivieri I, De Mita S, Cleyet-Marel JC, and Bena G. (2006). Recombination and selection shape the molecular diversity pattern of nitrogen-fixing *Sinorhizobium* sp. associated to Medicago. *Mol Ecol*, 15:2719-2734.
- Bakker PA, Pieterse CM, and van Loon LC. (2007). Induced Systemic Resistance by Fluorescent *Pseudomonas* spp. *Phytopathology*, 97:239-243.
- Banik JJ, and Brady SF. (2010). Recent application of metagenomic approaches toward the discovery of antimicrobials and other bioactive small molecules. *Curr Opin Microbiol*, 13:603-609.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, and Pevzner PA. (2012). SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol*, 19:455-477.
- Battini F, Cristani C, Giovannetti M, and Agnolucci M. (2016). Multifunctionality and diversity of culturable bacterial communities strictly associated with spores of the plant beneficial symbiont *Rhizophagus intraradices*. *Microbiol Res*, 183:68-79.
- Becker A, Berges H, Krol E, Bruand C, Ruberg S, Capela D, Lauber E, Meilhoc E, Ampe F, de Bruijn FJ, Fourment J, Francez-Charlot A, Kahn D, Kuster H, Liebe C, Puhler A, Weidner S, and Batut J. (2004). Global changes in gene expression in *Sinorhizobium meliloti* 1021 under microoxic and symbiotic conditions. *Mol Plant Microbe Interact*, 17:292-303.
- Bedmar EJ, Robles EF, and Delgado MJ. (2005). The complete denitrification pathway of the symbiotic, nitrogen-fixing bacterium *Bradyrhizobium japonicum*. *Biochem Soc Trans*, 33:141-144.
- Beghalem H, Aliliche K, Chriki A, and Landoulsi A. (2017). Molecular and phenotypic characterization of endophytic bacteria isolated from sulla nodules. *Microb Pathog*, 111:225-231.
- Beijerinck MW. (1888). Die bakterien der papilionaceenknollchen. *Botanische Zeitung*, pp. 740-750.
- Beneduzi A, Ambrosini A, and Passaglia LMP. (2012). Plant growth-promoting rhizobacteria (PGPR): Their potential as antagonists and biocontrol agents. *Genet Mol Biol*, 35:1044-1051.
- Bennett JW, and Bentley R. (2000). Seeing red: the story of prodigiosin. *Adv Appl Microbiol*, 47:1-32.
- Berg G. (2000). Diversity of antifungal and plant-associated *Serratia plymuthica* strains. *J Appl Microbiol*, 88:952-960.
- Beringer JE. (1974). R factor transfer in *Rhizobium leguminosarum*. *J Gen Microbiol*, 84 (1), pp. 188-98.

- Bertani G. (1951). Studies on lysogenesis. I. The mode of phage liberation by lysogenic *Escherichia coli*. *J Bacteriol*, 62:293-300.
- Biondi EG, Tatti E, Comparini D, Giuntini E, Mocali S, Giovannetti L, Bazzicalupo M, Mengoni A, and Viti C. (2009). Metabolic capacity of *Sinorhizobium (Ensifer) meliloti* strains as determined by phenotype MicroArray analysis. *Appl Environ Microbiol*, 75:5396-5404.
- Birnboim HC, and Doly J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res*, 7:1513-1523.
- Bizio B. (1823). Lettera di Bartolomeo Bizio al chiarissimo canonico Angelo Bellani sopra il fenomeno della polenta porporina. *Biblioteca Italiana o sia Giornale di Letteratura, Scienze e Arti (Anno VIII)*, pp. 275-295.
- Boiero L, Perrig D, Masciarelli O, Penna C, Cassán F, and Luna V. (2007). Phytohormone production by three strains of *Bradyrhizobium japonicum* and possible physiological and technological implications. *Appl Microbiol Biotechnol*, 74:874-880.
- Bolger AM, Lohse M, and Usadel B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, 30:2114-2120.
- Bollet C, Gainnier M, Sainty JM, Orhesser P, and De Micco P. (1991). *Serratia fonticola* isolated from a leg abscess. *J Clin Microbiol*, 29:834-835.
- Bollet C, Grimont P, Gainnier M, Geissler A, Sainty JM, and De Micco P. (1993). Fatal pneumonia due to *Serratia proteamaculans* subsp. *quinovora*. *J Clin Microbiol*, 31:444-445.
- Brockhurst MA, Morgan AD, Fenton A, and Buckling A. (2007). Experimental coevolution with bacteria and phage. The *Pseudomonas fluorescens*--Phi2 model system. *Infect Genet Evol*, 7:547-552.
- Brockwell J. (1982). Plant infection counts of rhizobia in soils. In Nitrogen Fixation in Legumes. Vincent, J.M. (ed.). New York, USA: *Academic Press*, pp. 41–57.
- Brockwell J, Pilka A, and Holliday RA. (1991). Soil pH is a major determinant of the numbers of naturally occurring *Rhizobium meliloti* in non-cultivated soils in central New South Wales. *Aust J Exp Agr*, 31:211-219.
- Bromfield ESP, Wheatcroft R, and Barran LR. (1994). Medium for direct isolation of *Rhizobium meliloti* from soils. *Soil Biol Biochem*, 26:423-428.
- Canfield DE, Glazer AN, and Falkowski PG. (2010). The evolution and future of Earth's nitrogen cycle. *Science*, 330:192-196.
- Case RJ, Boucher Y, Dahllöf I, Holmström C, Doolittle WF, and Kjelleberg S. (2007). Use of 16S rRNA and *rpoB* genes as molecular markers for microbial ecology studies. *Appl Environ Microbiol*, 73:278-288.
- Casida LE. (1982). *Ensifer adhaerens* gen. nov., sp. nov.: A Bacterial Predator of Bacteria in Soil. *Int J Syst Evol Microbiol*, 32:339-345.

- Castañeda LE, and Barbosa O. (2017). Metagenomic analysis exploring taxonomic and functional diversity of soil microbial communities in Chilean vineyards and surrounding native forests. *PeerJ*, 5: e3098.
- Cevallos MA, Cervantes-Rivera R, and Gutierrez-Rios RM. (2008). The repABC plasmid family. *Plasmid*, 60:19-37.
- Cha C, Gao P, Chen YC, Shaw PD, and Farrand SK. (1998). Production of acyl-homoserine lactone quorum-sensing signals by gram-negative plant-associated bacteria. *Mol Plant Microbe Interact*, 11:1119-1129.
- Chahboune R, Carro L, Peix A, Barrijal S, Velázquez E, and Bedmar EJ. (2011). *Bradyrhizobium cytisi* sp. nov., isolated from effective nodules of *Cytisus villosus*. *Int J Syst Evol Microbiol*, 61:2922-2927.
- Chahboune, R, Carro L, Peix A, Ramírez-Bahena MH, Barrijal S, Velázquez E, and Bedmar EJ. (2012). *Bradyrhizobium rifense* sp. nov. isolated from effective nodules of *Cytisus villosus* grown in the Moroccan Rif. *Syst Appl Microbiol* 35 (5):302-305.
- Chakraborty SP, and Sen Gupta SP. (1959). Fixation of Nitrogen by the Rice Plant. *Nature*, 184:2033.
- Chan JZ, Halachev MR, Loman NJ, Constantinidou C, and Pallen MJ. (2012). Defining bacterial species in the genomic era: insights from the genus *Acinetobacter*. *BMC Microbiol*, 12:302.
- Chang YL, Wang JY, Wang ET, Liu HC, Sui XH, and Chen WX. (2011). *Bradyrhizobium lablabi* sp. nov., isolated from effective nodules of *Lablab purpureus* and *Arachis hypogaea*. *Int J Syst Evol Microbiol*, 61:2496-2502.
- Chaparro JM, Badri DV, and Vivanco JM. (2014). Rhizosphere microbiome assemblage is affected by plant development. *ISME J*, 8:790-803.
- Chen L, He LY, Wang Q, and Sheng XF. (2016). Synergistic effects of plant growth-promoting *Neorhizobium huautlense* T1-17 and immobilizers on the growth and heavy metal accumulation of edible tissues of hot pepper. *J Hazard Mater*, 312:123-131.
- Chen WX, Yan GH, and Li JL. (1988). Numerical Taxonomic Study of Fast-Growing Soybean Rhizobia and a Proposal that *Rhizobium fredii* Be Assigned to *Sinorhizobium* gen. nov. *Int J Syst Bacteriol*, pp. 392-397.
- Chenna R, Sugawara H, Koike T, Lopez R, Gibson TJ, Higgins DG, and Thompson JD. (2003). Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Res*, 31:3497-3500.
- Chi F, Yang P, Han F, Jing Y, and Shen S. (2010). Proteomic analysis of rice seedlings infected by *Sinorhizobium meliloti* 1021. *Proteomics*, 10:1861-1874.
- Chibani-Chennoufi S, Bruttin A, Dillmann ML, and Brussow H. (2004). Phage-host interaction: an ecological perspective. *J Bacteriol*, 186:3677-3686.

- Chu H, Fierer N, Lauber CL, Caporaso JG, Knight R, and Grogan P. (2010). Soil bacterial diversity in the Arctic is not fundamentally different from that found in other biomes. *Environ Microbiol*, 12:2998-3006.
- Citorik RJ, Mimee M, and Lu TK. (2014). Bacteriophage-based synthetic biology for the study of infectious diseases. *Curr Opin Microbiol*, 19:59-69.
- Cole MA, and Elkan GH. (1979). Multiple antibiotic resistance in *Rhizobium japonicum*. *Appl Environ Microbiol*, 37:867-870.
- Cook MA, and Lopez JJ. (1998). *Serratia odorifera* biogroup I: an emerging pathogen. *J Am Osteopath Assoc*, 98:505-507.
- Costacurta A, and Vanderleyden J. (1995). Synthesis of phytohormones by plant-associated bacteria. *Crit Rev Microbiol*, 21:1-18.
- Cubitt MF, Hedley PE, Williamson NR, Morris JA, Campbell E, Toth IK, and Salmond GP. (2013). A metabolic regulator modulates virulence and quorum sensing signal production in *Pectobacterium atrosepticum*. *Mol Plant Microbe Interact*, 26:356-366.
- Daniel R. (2004). The soil metagenome-a rich resource for the discovery of novel natural products. *Curr Opin Biotechnol*, 15:199-204.
- Darling AE, Mau B, and Perna NT. (2010). progressiveMauve: Multiple Genome Alignment with Gene Gain, Loss and Rearrangement. *PLoS ONE*, 5:e11147.
- De Lajudie P, Willems A, Pot B, Dewettinck D, Maestrojuan G, Neyra M, Collins MD, Dreyfus B, Kersters K, and Gillis M. (1994). Polyphasic Taxonomy of Rhizobia: Emendation of the Genus *Sinorhizobium* and Description of *Sinorhizobium meliloti* comb. nov., *Sinorhizobium saheli* sp. nov., and *Sinorhizobium teranga* sp. nov. *Int J Syst Evol Microbiol*, 44:715.
- de Souza RSC, Okura VK, Armanhi JSL, Jorrin B, Lozano N, da Silva MJ, Gonzalez-Guerrero M, de Araujo LM, Verza NC, Bagheri HC, Imperial J, and Arruda P. (2016). Unlocking the bacterial and fungal communities assemblages of sugarcane microbiome. *Sci Rep*, 6 (28774).
- Delamuta JR, Ribeiro RA, Ormeno-Orrillo E, Melo IS, Martinez-Romero E, and Hungria M. (2013). Polyphasic evidence supporting the reclassification of *Bradyrhizobium japonicum* group Ia strains as *Bradyrhizobium diazoefficiens* sp. nov. *Int J Syst Evol Microbiol*, 63:3342-3351.
- Delamuta JR, Ribeiro RA, Ormeño-Orrillo E, Parma MM, Melo IS, Martínez-Romero E, and Hungria M. (2015). *Bradyrhizobium tropiciagri* sp. nov. and *Bradyrhizobium embrapense* sp. nov., nitrogen-fixing symbionts of tropical forage legumes. *Int J Syst Evol Microbiol*, 65:4424-4433.
- Demain AL. (1995). Why do microorganisms produce antimicrobials? In Fifty Years of Antimicrobials: Past Perspectives and Future Trends (Society for General Microbiology Symposium no. 53). Edited by P. A. Hunter, G. K. Darby & N.J. Russell. Cambridge: Cambridge University Press, pp. 205-228.

- Dimkpa C, Weinand T, and Asch F. (2009). Plant-rhizobacteria interactions alleviate abiotic stress conditions. *Plant Cell Environ*, 32:1682-1694.
- Dizman B, Elasri MO, and Mathias LJ. (2005). Synthesis, characterization, and antibacterial activities of novel methacrylate polymers containing norfloxacin. *Biomacromolecules*, 6:514-520.
- Döbereiner J. (1989). Isolation and identification of root associated diazotrophs. Nitrogen Fixation with Non-Legumes: The Fourth International Symposium on 'Nitrogen Fixation with Non-Legumes', Rio de Janeiro, 23–28 August 1987. Dordrecht: *Springer Netherlands*, pp. 103-108.
- Domik D, Thurmer A, Weise T, Brandt W, Daniel R, and Piechulla B. (2016a). A Terpene Synthase Is Involved in the Synthesis of the Volatile Organic Compound Sodorifen of *Serratia plymuthica* 4Rx13. *Front Microbiol*, 7:737.
- Domik D, Magnus N, and Piechulla B. (2016b). Analysis of a new cluster of genes involved in the synthesis of the unique volatile organic compound sodorifen of *Serratia plymuthica* 4Rx13. *FEMS Microbiol Lett*, 363.
- Dubouix A, Roques C, Segonds C, Jeannot MJ, Malavaud S, Daude S, Chabanon G, and Marty N. (2005). Epidemiological investigation of a *Serratia liquefaciens* outbreak in a neurosurgery department. *J Hosp Infect*, 60:8-13.
- Durán D, Rey L, Mayo J, Zúñiga-Dávila D, Imperial J, Ruiz-Argüeso T, Martínez-Romero E, and Ormeño-Orrillo E. (2014a). *Bradyrhizobium paxllaeri* sp. nov. and *Bradyrhizobium icense* sp. nov., nitrogen-fixing rhizobial symbionts of Lima bean (*Phaseolus lunatus* L.) in Peru. *Int J Syst Evol Microbiol*, 64:2072-2078.
- Durán D, Rey L, Navarro A, Busquets A, Imperial J, and Ruiz-Argüeso T. (2014b). *Bradyrhizobium valentinum* sp. nov., isolated from effective nodules of *Lupinus mariae-josephae*, a lupine endemic of basic-lime soils in Eastern Spain. *Syst Appl Microbiol*, 37:336-341.
- Ecker JR. (1995). The ethylene signal transduction pathway in plants. *Science*, 268:667-675.
- Effmert U, Kalderas J, Warnke R, and Piechulla B. (2012). Volatile mediated interactions between bacteria and fungi in the soil. *J Chem Ecol*, 38:665-703.
- Egamberdieva D, Wirth SJ, Alqarawi AA, Abd_Allah EF, and Hashem A. (2017). Phytohormones and Beneficial Microbes: Essential Components for Plants to Balance Stress and Fitness. *Front Microbiol*, 8 (2104).
- Eisenstein IB. (1990). Enterobacteriaceae. Churchill Livingstone New York: In: Mandell GL, Douglas RG, Bennett JE, Eds. *Principles and Practice Of Infectious Disease* 3rd ed, pp. 1658-1673.
- English G, Trunk K, Rao VA, Srikanthasani V, Hunter WN, and Coulthurst SJ. (2012). New secreted toxins and immunity proteins encoded within the Type VI secretion system gene cluster of *Serratia marcescens*. *Mol Microbiol*, 86:921-936.

- Erez Z, Steinberger-Levy I, Shamir M, Doron S, Stokar-Avihail A, Peleg Y, Melamed S, Leavitt A, Savidor A, Albeck S, Amitai G, and Sorek R. (2017). Communication between viruses guides lysis-lysogeny decisions. *Nature*, 541:488-493.
- Espona-Fiedler M, Soto-Cerrato V, Hosseini A, Lizcano JM, Guallar V, Quesada R, Gao T, and Perez-Tomas R. (2012). Identification of dual mTORC1 and mTORC2 inhibitors in melanoma cells: prodigiosin vs. obatoclox. *Biochem Pharmacol*, 83:489-496.
- Evans WR, Fleischman DE, Calvert HE, Pyati PV, Alter GM, and Rao NS. (1990). Bacteriochlorophyll and Photosynthetic Reaction Centers in *Rhizobium* Strain BTAi 1. *Appl Environ Microbiol*, 56:3445-3449.
- Evans TJ, Crow MA, Williamson NR, Orme W, Thomson NR, Komitopoulou E, and Salmond GP. (2010). Characterization of a broad-host-range flagellum-dependent phage that mediates high-efficiency generalized transduction in, and between, *Serratia* and *Pantoea*. *Microbiology*, 156:240-247.
- Fagervold SK, May HD, and Sowers KR. (2007). Microbial reductive dechlorination of aroclor 1260 in Baltimore harbor sediment microcosms is catalyzed by three phylotypes within the phylum *Chloroflexi*. *Appl Environ Microbiol*, 73:3009-3018.
- Fahraeus G. (1957). The infection of clover root hairs by nodule bacteria studied by a simple glass slide technique. *J Gen Microbiol*, 16:374-381.
- Farrand SK, Qin Y, and Oger P. (2002). Quorum-sensing system of *Agrobacterium* plasmids: analysis and utility. *Methods Enzymol*, 358:452-484.
- Fierer N, and Jackson RB. (2006). The diversity and biogeography of soil bacterial communities. *Proc Natl Acad Sci U S A*, 103:626-631.
- Fierer N, Strickland MS, Liptzin D, Bradford MA, and Cleveland CC. (2009). Global patterns in belowground communities. *Ecol Lett*, 12:1238-1249.
- Fierer N, Leff JW, Adams BJ, Nielsen UN, Bates ST, Lauber CL, Owens S, Gilbert JA, Wall DH, and Caporaso JG. (2012). Cross-biome metagenomic analyses of soil microbial communities and their functional attributes. *Proc Natl Acad Sci U S A*, 109:21390-21395.
- Fox GE, Wisotzkey JD, and Jurtshuk P, Jr. (1992). How close is close: 16S rRNA sequence identity may not be sufficient to guarantee species identity. *Int J Syst Bacteriol*, 42:166-170.
- Francis DD, Schepers JS, and Vigil MF. (1993). Post-Anthesis Nitrogen Loss from Corn. *Agron J*, 85:659-663.
- Frank B. (1889). Ueber die Pilzsymbiose der Leguminosen. *Dtsch Bot Ges*, pp. 332-346.
- Frink CR, Waggoner PE, and Ausubel JH. (1999). Nitrogen fertilizer: retrospect and prospect. *Proc Natl Acad Sci U S A*, 96:1175-1180.
- Fritsch MJ, Trunk K, Diniz JA, Guo M, Trost M, and Coulthurst SJ. (2013). Proteomic identification of novel secreted antibacterial toxins of the *Serratia marcescens* type VI secretion system. *Mol Cell Proteomics*, 12:2735-2749.

- Fürnkranz M, Adam E, Müller H, Grube M, Huss H, Winkler J, and Berg G. (2012). Promotion of growth, health and stress tolerance of Styrian oil pumpkins by bacterial endophytes. *Eur J Plant Pathol*, 134:509-519.
- Furukawa K. (2003). 'Super bugs' for bioremediation. *Trends Biotechnol*, 21:187-190.
- Galardini M, Mengoni A, Brilli M, Pini F, Fioravanti A, Lucas S, Lapidus A, Cheng JF, Goodwin L, Pitluck S, Land M, Hauser L, Woyke T, Mikhailova N, Ivanova N, Daligault H, Bruce D, Detter C, Tapia R, Han C, Teshima H, Mocali S, Bazzicalupo M, and Biondi EG. (2011). Exploring the symbiotic pangenome of the nitrogen-fixing bacterium *Sinorhizobium meliloti*. *BMC Genomics*, 12:235.
- Galibert F, Finan TM, Long SR, Puhler A, Abola P, Ampe F, Barloy-Hubler F, Barnett MJ, Becker A, Boistard P, Bothe G, Boutry M, Bowser L, Buhrmester J, Cadieu E, Capela D, Chain P, Cowie A, Davis RW, Dreano S, Federspiel NA, Fisher RF, Gloux S, Godrie T, Goffeau A, Golding B, Gouzy J, Gurjal M, Hernandez-Lucas I, Hong A, Huizar L, Hyman RW, Jones T, Kahn D, Kahn ML, Kalman S, Keating DH, Kiss E, Komp C, Lalaure V, Masuy D, Palm C, Peck MC, Pohl TM, Portetelle D, Purnelle B, Ramsperger U, Surzycki R, Thebault P, Vandenbol M, Vorholter FJ, Weidner S, Wells DH, Wong K, Yeh KC, and Batut J. (2001). The composite genome of the legume symbiont *Sinorhizobium meliloti*. *Science*, 293:668-672.
- Gans J, Wolinsky M, and Dunbar J. (2005). Computational improvements reveal great bacterial diversity and high metal toxicity in soil. *Science*, 309:1387-1390.
- Garbeva P, Postma J, van Veen JA, and van Elsas JD. (2006). Effect of above-ground plant species on soil microbial community structure and its impact on suppression of *Rhizoctonia solani* AG3. *Environ Microbiol*, 8:233-246.
- García-Fraile P, Chudíčková M, Benada O, Pikula J, and Kolařík M. (2015). *Serratia myotis* sp. nov. and *Serratia vespertilionis* sp. nov., isolated from bats hibernating in caves. *Int J Syst Evol Microbiol*, 65:90-94.
- Gault RR, and Schwinghamer EA. (1993). Direct isolation of *Bradyrhizobium japonicum* from soil. *Soil Biol Biochem*, 25:1161-1166.
- Geddes BA, and Oresnik IJ. (2014). Physiology, genetics, and biochemistry of carbon metabolism in the alphaproteobacterium *Sinorhizobium meliloti*. *Can J Microbiol*, 60:491-507.
- Gilbert JA, Jansson JK, and Knight R. (2014). The Earth Microbiome project: successes and aspirations. *BMC Biol*, 12:69.
- Giraud E, Moulin L, Vallenet D, Barbe V, Cytryn E, Avarre J-C, Jaubert M, Simon D, Cartieaux F, Prin Y, Bena G, Hannibal L, Fardoux J, Kojadinovic M, Vuillet L, Lajus A, Cruveiller S, Rouy Z, Mangenot S, Segurens B, Dossat C, Franck WL, Chang W-S, Saunders E, Bruce D, Richardson P, Normand P, Dreyfus B, Pignol D, Stacey G, Emerich D, Verméglio A, Médigue C, and Sadowsky M. (2007). Legumes Symbioses: Absence of Nod Genes in Photosynthetic Bradyrhizobia. *Science*, 316:1307-1312.
- Giuntini E, Mengoni A, De Filippo C, Cavalieri D, Aubin-Horth N, Landry CR, Becker A, and Bazzicalupo M. (2005). Large-scale genetic variation of the symbiosis-required

- megaplasmid pSymA revealed by comparative genomic analysis of *Sinorhizobium meliloti* natural strains. *BMC Genomics*, 6:158-158.
- Glick BR. (2014). Bacteria with ACC deaminase can promote plant growth and help to feed the world. *Microbiol Res*, 169:30-39.
- Gomes DF, da Silva Batista JS, Rolla AAP, da Silva LP, Bloch C, Galli-Terasawa LV, and Hungria M. (2014). Proteomic analysis of free-living *Bradyrhizobium diazoefficiens*: highlighting potential determinants of a successful symbiosis. *BMC Genomics*, 15:643.
- Gomez P, and Buckling A. (2011). Bacteria-phage antagonistic coevolution in soil. *Science*, 332:106-109.
- González V, Bustos P, Ramírez-Romero MA, Medrano-Soto A, Salgado H, Hernández-González I, Hernández-Celis JC, Quintero V, Moreno-Hagelsieb G, Girard L, Rodríguez O, Flores M, Cevallos MA, Collado-Vides J, Romero D, and Dávila G. (2003). The mosaic structure of the symbiotic plasmid of *Rhizobium etli* CFN42 and its relation to other symbiotic genome compartments. *Genome Biol*, 4:R36.
- Goodridge LD, and Bisha B. (2011). Phage-based biocontrol strategies to reduce foodborne pathogens in foods. *Bacteriophage*, 1:130-137.
- Gorlach K, Shingaki R, Morisaki H, and Hattori T. (1994). Construction of eco-collection of paddy field soil bacteria for population analysis. *J Gen Appl Microbiol*, 40:509-517.
- Graham PH, Draeger KJ, Ferrey ML, Conroy MJ, Hammer BE, Martinez E, Aarons SR, and Quinto C. (1994). Acid pH tolerance in strains of *Rhizobium* and *Bradyrhizobium*, and initial studies on the basis for acid tolerance of *Rhizobium tropici* UMR1899. *Can J Microbiol*, 40:198-207.
- Grant CL, and Pramer D. (1962). Minor element composition of yeast extract. *J Bacteriol*, 84:869-870.
- Grimont PAD, Grimont F, and Starr MP. (1978). *Serratia proteamaculans* (Paine and Stansfield) comb. no., a Senior Subjective Synonym of *Serratia liquefaciens* (Grimes and Hennerty) Bascomb et al. *Int J Syst Bacteriol*, pp. 503-510.
- Grimont F, and Grimont PAD. (1984). Genus *Serratia* Bizio 1823, 288 AL. Baltimore: In: Krieg, N. R., and Holt, J.G. (ed.), *Bergey's manual of systematic bacteriology*. Williams and Wilkins, pp. 477-484.
- Grimont PAD, Jackson TA, Ageron E, and Noonan MJ. (1988). *Serratia entomophila* sp. nov. Associated with Amber Disease in the New Zealand Grass Grub *Costelytra zealandica*. *Int J Syst Evol Microbiol*, 38:1.
- Grosch R, Faltin F, Lottmann J, Kofoet A, and Berg G. (2005). Effectiveness of 3 antagonistic bacterial isolates to control *Rhizoctonia solani* Kuhn on lettuce and potato. *Can J Microbiol*, 51:345-353.
- Gruber N, and Galloway JN. (2008). An Earth-system perspective of the global nitrogen cycle. *Nature*, 451:293-296.

- Guerinot ML, Meidl EJ, and Plessner O. (1990). Citrate as a siderophore in *Bradyrhizobium japonicum*. *J Bacteriol*, 172:3298-3303.
- Guerrouj K, Ruíz-Díez B, Chahboune R, Ramírez-Bahena M-H, Abdelmoumen H, Quiñones MA, El Idrissi MM, Velázquez E, Fernández-Pascual M, Bedmar EJ, and Peix A. (2013). Definition of a novel symbiovar (sv. *retamae*) within *Bradyrhizobium retamae* sp. nov., nodulating *Retama sphaerocarpa* and *Retama monosperma*. *Syst Appl Microbiol*, 36:218-223.
- Guo J, and Chi J. (2014). Effect of Cd-tolerant plant growth-promoting *Rhizobium* on plant growth and Cd uptake by *Lolium multiflorum* Lam. and *Glycine max* (L.) Merr. in Cd-contaminated soil. *Plant Soil*, 375 (1-2), pp. 205–214.
- Gutiérrez D, Hernando Y, Palacios JM, Imperial J, and Ruiz-Argüeso T. (1997). FnrN controls symbiotic nitrogen fixation and hydrogenase activities in *Rhizobium leguminosarum* biovar *viciae* UPM791. *J Bacteriol*, 179:5264-5270.
- Gutiérrez RA. (2012). Systems biology for enhanced plant nitrogen nutrition. *Science*, 336:1673-1675.
- Gyaneshwar P, James EK, Mathan N, Reddy PM, Reinhold-Hurek B, and Ladha JK. (2001). Endophytic colonization of rice by a diazotrophic strain of *Serratia marcescens*. *J Bacteriol*, 183:2634-2645.
- Hafeez FY, Asad S, Ahmad T, and Malik KA. (1995). Host specificity and characterization of fast-growing rhizobia from *Macroptilium atropurpureum* cv. Siratro in Pakistan. *Soil Biol Biochem*, 27:729-733.
- Han K, Li ZF, Peng R, Zhu LP, Zhou T, Wang LG, Li SG, Zhang XB, Hu W, Wu ZH, Qin N, and Li YZ. (2013). Extraordinary expansion of a *Sorangium cellulosum* genome from an alkaline milieu. *Sci Rep*, 3:2101.
- Hanus FJ, Maier RJ, and Evans HJ. (1979). Autotrophic growth of H₂-uptake-positive strains of *Rhizobium japonicum* in an atmosphere supplied with hydrogen gas. *Proc Natl Acad Sci U S A*, 76:1788-1792.
- Harnett SJ, Allen KD, and Macmillan RR. (2001). Critical care unit outbreak of *Serratia liquefaciens* from contaminated pressure monitoring equipment. *J Hosp Infect*, 47:301-307.
- Harshey RM, and Matsuyama T. (1994). Dimorphic transition in *Escherichia coli* and *Salmonella typhimurium*: surface-induced differentiation into hyperflagellate swarmer cells. *Proc Natl Acad Sci U S A*, 91:8631-8635.
- Hendrix RW. (2003). Bacteriophage genomics. *Curr Opin Microbiol*, 6:506-511.
- Herskowitz I, and Hagen D. (1980). The lysis-lysogeny decision of phage lambda: explicit programming and responsiveness. *Annu Rev Genet*, 14:399-445.
- Hibbing ME, Fuqua C, Parsek MR, and Peterson SB. (2010). Bacterial competition: surviving and thriving in the microbial jungle. *Nat Rev Microbiol*, 8:15-25.

- Hirel B, Tétu T, Lea PJ, and Dubois F. (2011). Improving Nitrogen Use Efficiency in Crops for Sustainable Agriculture. *Sustainability*, 3.
- Hirsch PR, and Mauchline TH. (2015). The Importance of the Microbial N Cycle in Soil for Crop Plant Nutrition. *Adv Appl Microbiol*, 93:45-71.
- Hökeberg M, Gerhardson B, and Johnsson L. (1997). Biological control of cereal seed-borne diseases by seed bacterization with greenhouse-selected bacteria. *Eur J Plant Pathol*, 103:25-33.
- Honma M, and Shimomura T. (1978). Metabolism of 1-Aminocyclopropane-1-carboxylic Acid. *Agric Biol Chem*, 42:1825-1831.
- Hsieh HY, Shieh JJ, Chen CJ, Pan MY, Yang SY, Lin SC, Chang JS, Lee AY, and Chang CC. (2012). Prodigiosin down-regulates SKP2 to induce p27(KIP1) stabilization and antiproliferation in human lung adenocarcinoma cells. *Br J Pharmacol*, 166:2095-2108.
- Iqbal HA, Feng Z, and Brady SF. (2012). Biocatalysts and small molecule products from metagenomic studies. *Curr Opin Chem Biol*, 16 (1):109-116.
- Islam MS, Kawasaki H, Muramatsu Y, Nakagawa Y, and Seki T. (2008). *Bradyrhizobium iriomotense* sp. nov., isolated from a tumor-like root of the legume *Entada koshunensis* from Iriomote Island in Japan. *Biosci Biotechnol Biochem*, 72:1416-1429.
- Iyer B, Rajput MS, and Rajkumar S. (2017). Effect of succinate on phosphate solubilization in nitrogen fixing bacteria harbouring chick pea and their effect on plant growth. *Microbiol Res*, 202:43-50.
- Jang JH, Lee D, Cha S, and Seo T. (2017). *Ensifer collicola* sp. nov., a bacterium isolated from soil in South Korea. *J Microbiol*, 55:520-524.
- Janssen PH. (2006). Identifying the dominant soil bacterial taxa in libraries of 16S rRNA and 16S rRNA genes. *Appl Environ Microbiol*, 72:1719-1728.
- Jeong H, Yim JH, Lee C, Choi SH, Park YK, Yoon SH, Hur CG, Kang HY, Kim D, Lee HH, Park KH, Park SH, Park HS, Lee HK, Oh TK, and Kim JF. (2005). Genomic blueprint of *Hahella chejuensis*, a marine microbe producing an algicidal agent. *Nucleic Acids Res*, 33:7066-7073.
- Johnston AW, and Beringer JE. (1975). Identification of the rhizobium strains in pea root nodules using genetic markers. *J Gen Microbiol*, 87:343-350.
- Jolliffe IT, and Cadima J. (2016). Principal component analysis: a review and recent developments. *Philos Trans A Math Phys and Eng Sci*, 374 (2065):20150202.
- Jones FP, Clark IM, King R, Shaw LJ, Woodward MJ, and Hirsch PR. (2016). Novel European free-living, non-diazotrophic *Bradyrhizobium* isolates from contrasting soils that lack nodulation and nitrogen fixation genes - a genome comparison. *Sci Rep*, 6:25858.

- Jones KM, Kobayashi H, Davies BW, Taga ME, and Walker GC. (2007). How rhizobial symbionts invade plants: the *Sinorhizobium-Medicago* model. *Nat Rev Microbiol*, 5:619-633.
- Jordan DC. (1982). NOTES: Transfer of *Rhizobium japonicum* Buchanan 1980 to *Bradyrhizobium* gen. nov., a Genus of Slow-Growing, Root Nodule Bacteria from Leguminous Plants. *Int J Syst Evol Microbiol*, 32:136.
- Jorrín B. (2016). Genomics of Specificity in the Symbiotic Interaction between *Rhizobium leguminosarum* and Legumes. Madrid, Spain: Universidad Politécnica de Madrid.
- Jorrin B and Imperial J. (2015). Population Genomics Analysis of Legume Host Preference for Specific Rhizobial Genotypes in the *Rhizobium leguminosarum* bv. *viciae* Symbioses. *Mol Plant Microbe Interact*, 28:310-318.
- Kai M, Effmert U, Berg G, and Piechulla B. (2007). Volatiles of bacterial antagonists inhibit mycelial growth of the plant pathogen *Rhizoctonia solani*. *Arch Microbiol*, 187:351-360.
- Kai M, Crespo E, Cristescu SM, Harren FJ, Francke W, and Piechulla B. (2010). *Serratia odorifera*: analysis of volatile emission and biological impact of volatile compounds on *Arabidopsis thaliana*. *Appl Microbiol Biotechnol*, 88:965-976.
- Kämpfer P, and Glaeser SP. (2016). *Serratia aquatilis* sp. nov., isolated from drinking water systems. *Int J Syst Evol Microbiol*, 66:407-413.
- Kaneko T, Nakamura Y, Sato S, Asamizu E, Kato T, Sasamoto S, Watanabe A, Idesawa K, Ishikawa A, and Kawashima K. (2000). Complete genome structure of the nitrogen-fixing symbiotic bacterium *Mesorhizobium loti*. *DNA Res*, 7.
- Kaneko T, Nakamura Y, Sato S, Minamisawa K, Uchiumi T, Sasamoto S, Watanabe A, Idesawa K, Iriguchi M, and Kawashima K. (2002). Complete genomic sequence of nitrogen-fixing symbiotic bacterium *Bradyrhizobium japonicum* USDA 110. *DNA Res*, 9.
- Kaneko T, Maita H, Hirakawa H, Uchiike N, Minamisawa K, Watanabe A, and Sato S. (2011). Complete Genome Sequence of the Soybean Symbiont *Bradyrhizobium japonicum* Strain USDA6^T. *Genes*, 2:763-787.
- Kant S, Bi Y-M, and Rothstein SJ. (2011). Understanding plant response to nitrogen limitation for the improvement of crop nitrogen use efficiency. *J Exp Bot*, 62:1499-1509.
- Kawauchi K, Shibutani K, Yagisawa H, Kamata H, Nakatsuji S, Anzai H, Yokoyama Y, Ikegami Y, Moriyama Y, and Hirata H. (1997). A possible immunosuppressant, cycloprodigiosin hydrochloride, obtained from *Pseudoalteromonas denitrificans*. *Biochem Biophys Res Commun*, 237:543-547.
- Kearns DB. (2010). A field guide to bacterial swarming motility. *Nat Rev Microbiol*, 8:634-644.

- Khamis A, Colson P, Raoult D, and Scola BL. (2003). Usefulness of *rpoB* gene sequencing for identification of *Afipia* and *Bosea* species, including a strategy for choosing discriminative partial sequences. *Appl Environ Microbiol*, 69:6740-6749.
- Kingsley MT, and Bohlool BB. (1981). Release of *Rhizobium* spp. from Tropical Soils and Recovery for Immunofluorescence Enumeration. *Appl Environ Microbiol*, 42:241-248.
- Kinkle BK, Sadowsky MJ, Johnstone K, and Koskinen WC. (1994). Tellurium and Selenium Resistance in Rhizobia and Its Potential Use for Direct Isolation of *Rhizobium meliloti* from Soil. *Appl Environ Microbiol*, 60:1674-1677.
- Kloepper JW, Ryu CM, and Zhang S. (2004). Induced Systemic Resistance and Promotion of Plant Growth by *Bacillus* spp. *Phytopathology*, 94:1259-1266.
- Konstantinidis KT, and Tiedje JM. (2004). Trends between gene content and genome size in prokaryotic species with larger genomes. *Proc Natl Acad Sci U S A*, 101:3160-3165.
- Koskella B, and Meaden S. (2013). Understanding bacteriophage specificity in natural microbial communities. *Viruses*, 5:806-823.
- Koskella B, and Brockhurst MA. (2014). Bacteria–phage coevolution as a driver of ecological and evolutionary processes in microbial communities. *FEMS Microbiol Rev*, 38:916-931.
- Kumar R, Khurana A, and Sharma AK. (2014). Role of plant hormones and their interplay in development and ripening of fleshy fruits. *J Exp Bot*, 65:4561-4575.
- Kumar S, Stecher G, and Tamura K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol*, 33:1870-1874.
- Kurze S, Bahl H, Dahl R, and Berg G. (2001). Biological Control of Fungal Strawberry Diseases by *Serratia plymuthica* HRO-C48. *Plant Dis*, 85:529-534.
- Kuykendall LD, Saxena B, Devine TE, and Udell SE. (1992). Genetic diversity in *Bradyrhizobium japonicum* Jordan 1982 and a proposal for *Bradyrhizobium elkanii* sp.nov. *Can J Microbiol*, 38:501-505.
- Kwak YS, and Weller DM. (2013). Take-all of Wheat and Natural Disease Suppression: A Review. *Plant Pathol J*, 29:125-135.
- Lagares A, Sanjuán J, and Pistorio M. (2014). The Plasmid Mobilome of the Model Plant-Symbiont *Sinorhizobium meliloti*: Coming up with New Questions and Answers. *Microbiol Spectr*, 2 (5).
- Laguerre G, Bardin M, and Amarger N. (1993). Isolation from soil of symbiotic and nonsymbiotic *Rhizobium leguminosarum* by DNA hybridization. *Can J Microbiol*, 39:1142-1149.
- Laguerre G, Louvrier P, Allard MR, and Amarger N. (2003). Compatibility of rhizobial genotypes within natural populations of *Rhizobium leguminosarum* biovar viciae for nodulation of host legumes. *Appl Environ Microbiol*, 69:2276-2283.

- Lambert M, Wolfender J-L, Stærk D, Christensen SB, Hostettmann K, and Jaroszewski JW. (2007). Identification of Natural Products Using HPLC-SPE Combined with CapNMR. *Anal Chem*, 79:727-735.
- Land M, Hauser L, Jun SR, Nookaew I, Leuze MR, Ahn TH, Karpinets T, Lund O, Kora G, Wassenaar T, Poudel S, and Ussery DW. (2015). Insights from 20 years of bacterial genome sequencing. *Funct Integr Genomics*, 15:141-161.
- Lauber CL, Hamady M, Knight R, and Fierer N. (2009). Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Appl Environ Microbiol*, 75:5111-5120.
- Lavania M, Chauhan PS, Chauhan SV, Singh HB, and Nautiyal CS. (2006). Induction of plant defense enzymes and phenolics by treatment with plant growth-promoting rhizobacteria *Serratia marcescens* NBRI1213. *Curr Microbiol*, 52:363-368.
- Lee B, Farag MA, Park HB, Kloepper JW, Lee SH, and Ryu CM. (2012). Induced resistance by a long-chain bacterial volatile: elicitation of plant systemic defense by a C13 volatile produced by *Paenibacillus polymyxa*. *PLoS ONE* 7:e48744.
- Lee J, Carey J, and Perlman DC. (2006). Pneumonia and bacteremia due to *Serratia odorifera*. *J Infect*, 53:212-214.
- Li Lu Y, Chen WF, Li Han L, Wang ET, and Chen WX. (2009). *Rhizobium alkalisoli* sp. nov., isolated from *Caragana intermedia* growing in saline-alkaline soils in the north of China. *Int J Syst Evol Microbiol*, 59:3006-3011.
- Li Y, Yan J, Yu B, Wang ET, Li X, Yan H, Liu W, and Xie Z. (2016). *Ensifer alkalisoli* sp. nov. isolated from root nodules of *Sesbania cannabina* grown in saline-alkaline soils. *Int J Syst Evol Microbiol*, 66:5294-5300.
- Li Y, Pang HD, He LY, Wang Q, and Sheng XF. (2017). Cd immobilization and reduced tissue Cd accumulation of rice (*Oryza sativa* wuyun-23) in the presence of heavy metal-resistant bacteria. *Ecotoxicol Environ Saf*, 138:56-63.
- Li Z, Ma Z, Hao X, Rensing C, and Wei G. (2014). Genes Conferring Copper Resistance in *Sinorhizobium meliloti* CCNWSX0020 Also Promote the Growth of *Medicago lupulina* in Copper-Contaminated Soil. *Appl Environ Microbiol*, 80:1961-1971.
- Lima-Mendez G, Toussaint A, and Leplae R. (2007). Analysis of the phage sequence space: the benefit of structured information. *Virology*, 365:241-249.
- Lindblad P, Rai AN, and Bergman B. (1987). The *Cycas revoluta*-*Nostoc* Symbiosis: Enzyme Activities of Nitrogen and Carbon Metabolism in the Cyanobiont. *Microbiology*, 133:1695.
- Lindström K. (1989). *Rhizobium galegae*, a New Species of Legume Root Nodule Bacteria. *Int J Syst Bacteriol*, pp. 365-367.
- Liu J, Sui Y, Yu Z, Shi Y, Chu H, Jin J, Liu X, and Wang G. (2014). High throughput sequencing analysis of biogeographical distribution of bacterial communities in the black soils of northeast China. *Soil Biol Biochem*, 70:113-122.

- Liu X, Jia J, Popat R, Ortori CA, Li J, Diggle SP, Gao K, and Cámara M. (2011). Characterisation of two quorum sensing systems in the endophytic *Serratia plymuthica* strain G3: differential control of motility and biofilm formation according to life-style. *BMC Microbiol*, 11(1):26.
- Lloret L, Ormeno-Orrillo E, Rincon R, Martinez-Romero J, Rogel-Hernandez MA, and Martinez-Romero E. (2007). *Ensifer mexicanus* sp. nov. a new species nodulating *Acacia angustissima* (Mill.) Kuntze in Mexico. *Syst Appl Microbiol*, 30:280-290.
- López-Guerrero MG, Ormeño-Orrillo E, Acosta JL, Mendoza-Vargas A, Rogel MA, Ramírez MA, Rosenblueth M, Martínez-Romero J, and Martínez-Romero E. (2012). Rhizobial extrachromosomal replicon variability, stability and expression in natural niches. *Plasmid*, 68:149-158.
- Louvrier P, Laguerre G, and Amarger N. (1995). Semiselective medium for isolation of *Rhizobium leguminosarum* from soils. *Soil Biol Biochem*, 27:919-924.
- Louvrier P, Laguerre G, and Amarger N. (1996). Distribution of Symbiotic Genotypes in *Rhizobium leguminosarum* biovar viciae Populations Isolated Directly from Soils. *Appl Environ Microbiol*, 62:4202-4205.
- Lu CH, Lin SC, Yang SY, Pan MY, Lin YW, Hsu CY, Wei YH, Chang JS, and Chang CC. (2012). Prodigiosin-induced cytotoxicity involves RAD51 down-regulation through the JNK and p38 MAPK pathways in human breast carcinoma cell lines. *Toxicol Lett*, 212:83-89.
- Lu JK, Dou YJ, Zhu YJ, Wang SK, Sui XH, and Kang LH. (2014). *Bradyrhizobium ganzhouense* sp. nov., an effective symbiotic bacterium isolated from *Acacia melanoxylon* R. Br. nodules. *Int J Syst Evol Microbiol*, 64:1900-1905.
- Lucy M, Reed E, and Glick BR. (2004). Applications of free living plant growth-promoting rhizobacteria. *Antonie Van Leeuwenhoek*, 86:1-25.
- Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, Tremblay J, Engelbrektson A, Kunin V, Rio TGd, Edgar RC, Eickhorst T, Ley RE, Hugenholtz P, Tringe SG, and Dangl JL. (2012). Defining the core *Arabidopsis thaliana* root microbiome. *Nature*, 488:86-90.
- Macdonald CA, Clark IM, Hirsch PR, Zhao FJ, and McGrath SP. (2011). Development of a real-time PCR assay for detection and quantification of *Rhizobium leguminosarum* bacteria and discrimination between different biovars in zinc-contaminated soil. *Appl Environ Microbiol*, 77:4626-4633.
- Marschner H. (1995). Mineral Nutrition of Higher Plants: *Academic Press*. 889 pp.
- Matilla MA, and Salmond GPC. (2012a). Complete Genome Sequence of *Serratia plymuthica* Bacteriophage phi MAM1. *J Virol*, 86:13872-13873.
- Matilla MA, Stockmann H, Leeper FJ, and Salmond GPC. (2012b). Bacterial biosynthetic gene clusters encoding the anti-cancer haterumalide class of molecules: biogenesis of the broad spectrum antifungal and anti-oomycete compound, oocydin A. *J Biol Chem*, 287:39125-39138.

- Matilla MA, and Salmond GPC. (2014). Bacteriophage varphiMAM1, a viunalikevirus, is a broad-host-range, high-efficiency generalized transducer that infects environmental and clinical isolates of the enterobacterial genera *Serratia* and *Kluyvera*. *Appl Environ Microbiol*, 80:6446-6457.
- Matilla MA, Leeper FJ, and Salmond GPC. (2015). Biosynthesis of the antifungal haterumalide, oocydin A, in *Serratia*, and its regulation by quorum sensing, RpoS and Hfq. *Environ Microbiol*, 17:2993-3008.
- Mazzola M, and White FF. (1994). A mutation in the indole-3-acetic acid biosynthesis pathway of *Pseudomonas syringae* pv. *syringae* affects growth in *Phaseolus vulgaris* and syringomycin production. *J Bacteriol*, 176:1374-1382.
- McClellan KH, Winson MK, Fish L, Taylor A, Chhabra SR, Camara M, Daykin M, Lamb JH, Swift S, Bycroft BW, Stewart GS, and Williams P. (1997). Quorum sensing and *Chromobacterium violaceum*: exploitation of violacein production and inhibition for the detection of N-acylhomoserine lactones. *Microbiology*, 143 (Pt 12):3703-3711.
- McCullough NB. (1949). Laboratory Tests in the Diagnosis of Brucellosis. *Am J Public Health Nations Health*, 39:866-869.
- McGowan SJ, Sebaihia M, Porter LE, Stewart GSAB, Williams P, Bycroft BW, and Salmond GPC. (1996). Analysis of bacterial carbapenem antibiotic production genes reveals a novel β -lactam biosynthesis pathway. *Mol Microbiol*, 22:415-426.
- Mengel K, and Kirkby E A. (1978). Principles of plant nutrition, 3rd edn. Bern, Switzerland: International Potash Institute. 849 pp.
- Menna P, Barcellos FG, and Hungria M. (2009). Phylogeny and taxonomy of a diverse collection of *Bradyrhizobium* strains based on multilocus sequence analysis of the 16S rRNA gene, ITS region and glnII, recA, atpD and dnaK genes. *Int J Syst Evol Microbiol*, 59:2934-2950.
- Menna P, and Hungria M. (2011). Phylogeny of nodulation and nitrogen-fixation genes in *Bradyrhizobium*: supporting evidence for the theory of monophyletic origin, and spread and maintenance by both horizontal and vertical transfer. *Int J Syst Evol Microbiol*, 61:3052-3067.
- Meyer O. (1994). Functional Groups of Microorganisms. *Biodiversity and Ecosystem Function*. Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 67-96.
- Miao V, and Davies J. (2010). Actinobacteria: the good, the bad, and the ugly. *Antonie Van Leeuwenhoek*, 98:143-150.
- Mitarai N, Brown S, and Sneppen K. (2016). Population Dynamics of Phage and Bacteria in Spatially Structured Habitats Using Phage lambda and *Escherichia coli*. *J Bacteriol*, 198:1783-1793.
- Mitsch MJ, diCenzo GC, Cowie A, and Finan TM. (2017). Succinate Transport Is Not Essential for Symbiotic Nitrogen Fixation by *Sinorhizobium meliloti* or *Rhizobium leguminosarum*. *Appl Environ Microbiol*, 84 (1):16.

- Moak M, and Molineux IJ. (2003). Peptidoglycan hydrolytic activities associated with bacteriophage virions. *Mol Microbiol*, 51:1169-1183.
- Moellering RC, Eliopoulos GM, and Sentochnik DE. (1989). The carbapenems: new broad spectrum beta-lactam antibiotics. *J Antimicrob Chemother*, 24 Suppl A:1-7.
- Mollet C, Drancourt M, and Raoult D. (1997). *rpoB* sequence analysis as a novel basis for bacterial identification. *Mol Microbiol*, 26:1005-1011.
- Molouba F, Lorquin J, Willems A, Hoste B, Giraud E, Dreyfus B, Gillis M, de Lajudie P, and Masson-Boivin C. (1999). Photosynthetic bradyrhizobia from *Aeschynomene* spp. are specific to stem-nodulated species and form a separate 16S ribosomal DNA restriction fragment length polymorphism group. *Appl Environ Microbiol*, 65:3084-3094.
- Monson R, Burr T, Carlton T, Liu H, Hedley P, Toth I, and Salmond GPC. (2013). Identification of genes in the VirR regulon of *Pectobacterium atrosepticum* and characterization of their roles in quorum sensing-dependent virulence. *Environ Microbiol*, 15:687-701.
- Mooshammer M, Wanek W, Hämmerle I, Fuchslueger L, Hofhansl F, Knoltsch A, Schnecker J, Takriti M, Watzka M, Wild B, Keiblinger KM, Zechmeister-Boltenstern S, and Richter A. (2014). Adjustment of microbial nitrogen use efficiency to carbon:nitrogen imbalances regulates soil nitrogen cycling. *Nat Commun*, 5:3694.
- Morris VL, Jackson DP, Grattan M, Ainsworth T, and Cuppels DA. (1995). Isolation and sequence analysis of the *Pseudomonas syringae* pv. tomato gene encoding a 2,3-diphosphoglycerate-independent phosphoglyceromutase. *J Bacteriol*, 177:1727-1733.
- Moulin L, Béna G, Boivin-Masson C, and Stepkowski T. (2004). Phylogenetic analyses of symbiotic nodulation genes support vertical and lateral gene co-transfer within the *Bradyrhizobium* genus. *Mol Phylogenet Evol*, 30:720-732.
- Mousavi SA, Österman J, Wahlberg N, Nesme X, Lavire C, Vial L, Paulin L, de Lajudie P, and Lindström K. (2014). Phylogeny of the *Rhizobium*–*Allorhizobium*–*Agrobacterium* clade supports the delineation of *Neorhizobium* gen. nov. *Syst Appl Microbiol*, 37:208-215.
- Müller H, Westendorf C, Leitner E, Chernin L, Riedel K, Schmidt S, Eberl L, and Berg G. (2009). Quorum-sensing effects in the antagonistic rhizosphere bacterium *Serratia plymuthica* HRO-C48. *FEMS Microbiol Ecol*, 67:468-478.
- Müller H, Fürnkranz M, Grube M, and Berg G. (2013). Genome Sequence of *Serratia plymuthica* Strain S13, an Endophyte with Germination- and Plant-Growth-Promoting Activity from the Flower of Styrian Oil Pumpkin. *Genome Announc*, 1:e00594-00513.
- Mulley G, White JP, Karunakaran R, Prell J, Bourdes A, Bunnewell S, Hill L, and Poole PS. (2011). Mutation of GOGAT prevents pea bacteroid formation and N₂ fixation by globally downregulating transport of organic nitrogen sources. *Mol Microbiol*, 80:149-167.

- Murdoch SL, Trunk K, English G, Fritsch MJ, Pourkarimi E, and Coulthurst SJ. (2011). The opportunistic pathogen *Serratia marcescens* utilizes type VI secretion to target bacterial competitors. *J Bacteriol*, 193:6057-6069.
- Nambiar PTC, and Sivaramakrishnan S. (1987). Detection and assay of siderophores in cowpea rhizobia (*Bradyrhizobium*) using radioactive Fe (⁵⁹Fe). *Lett Appl Microbiol*, 4:37-40.
- Nelson MS, and Sadowsky MJ. (2015). Secretion systems and signal exchange between nitrogen-fixing rhizobia and legumes. *Front Plant Sci*, 6:491.
- Neupane S, Goodwin LA, Högberg N, Kyrpides NC, Alström S, Bruce D, Quintana B, Munk C, Daligault H, Teshima H, Davenport K, Reitenga K, Green L, Chain P, Erkkila T, Gu W, Zhang X, Xu Y, Kunde Y, Chertkov O, Han J, Han C, Detter JC, Ivanova N, Pati A, Chen A, Szeto E, Mavromatis K, Huntemann M, Nolan M, Pitluck S, Deshpande S, Markowitz V, Pagani I, Klenk HP, Woyke T, and Finlay RD. (2013). Non-contiguous finished genome sequence of plant-growth promoting *Serratia proteamaculans* S4. *Stand Genomic Sci*, 8:441-449.
- Nick G, de Lajudie P, Eardly BD, Suomalainen S, Paulin L, Zhang X, Gillis M, and Lindstrom K. (1999). *Sinorhizobium arboris* sp. nov. and *Sinorhizobium kostiense* sp. nov., isolated from leguminous trees in Sudan and Kenya. *Int J Syst Bacteriol* 49, Pt 4:1359-1368.
- Nishida M, Mine Y, Nonoyama S, and Kojo H. (1977). Nocardicin A, a new monocyclic beta-lactam antibiotic III. In vitro evaluation. *J Antibiot (Tokyo)*, 30:917-925.
- Niveshika, Verma E, Mishra AK, Singh AK, and Singh VK. (2016). Structural Elucidation and Molecular Docking of a Novel Antibiotic Compound from Cyanobacterium *Nostoc* sp. MGL001. *Front Microbiol*, 7:1899.
- Nouh F, and Bhandari S. (2000). CAPD peritonitis due to *Serratia plymuthica*. *Perit Dial Int*, 20:349.
- Ofek-Lalzar M, Sela N, Goldman-Voronov M, Green SJ, Hadar Y, and Minz D. (2014). Niche and host-associated functional signatures of the root surface microbiome. *Nat Commun*, 5.
- Okubo T, Tsukui T, Maita H, Okamoto S, Oshima K, Fujisawa T, Saito A, Futamata H, Hattori R, Shimomura Y, Haruta S, Morimoto S, Wang Y, Sakai Y, Hattori M, Aizawa S, Nagashima KV, Masuda S, Hattori T, Yamashita A, Bao Z, Hayatsu M, Kajiya-Kanegae H, Yoshinaga I, Sakamoto K, Toyota K, Nakao M, Kohara M, Anda M, Niwa R, Jung-Hwan P, Sameshima-Saito R, Tokuda S, Yamamoto S, Yokoyama T, Akutsu T, Nakamura Y, Nakahira-Yanaka Y, Takada Hoshino Y, Hirakawa H, Mitsui H, Terasawa K, Itakura M, Sato S, Ikeda-Ohtsubo W, Sakakura N, Kaminuma E, and Minamisawa K. (2012). Complete genome sequence of *Bradyrhizobium* sp. S23321: insights into symbiosis evolution in soil oligotrophs. *Microbes Environ*, 27:306-315.
- Oldroyd GE, Murray JD, Poole PS, and Downie JA. (2011). The rules of engagement in the legume-rhizobial symbiosis. *Annu Rev Genet*, 45:119-144.
- Op den Camp R, Streng A, De Mita S, Cao Q, Polone E, Liu W, Ammiraju JS, Kudrna D, Wing R, Untergasser A, Bisseling T, and Geurts R. (2011). LysM-type mycorrhizal

- receptor recruited for rhizobium symbiosis in nonlegume *Parasponia*. *Science*, 331:909-912.
- Österman J, Marsh J, Laine PK, Zeng Z, Alatalo E, Sullivan JT, Young JPW, Thomas-Oates J, Paulin L, and Lindström K. (2014). Genome sequencing of two *Neorhizobium galegae* strains reveals a noeT gene responsible for the unusual acetylation of the nodulation factors. *BMC Genomics*, 15:500.
- Österman J, Mousavi SA, Koskinen P, Paulin L, and Lindström K. (2015). Genomic features separating ten strains of *Neorhizobium galegae* with different symbiotic phenotypes. *BMC Genomics* 16:348.
- Park YS, Dutta S, Ann M, Raaijmakers JM, and Park K. (2015). Promotion of plant growth by *Pseudomonas fluorescens* strain SS101 via novel volatile organic compounds. *Biochem Biophys Res Commun*, 461:361-365.
- Parker WL, Rathnum ML, Wells JS, Jr., Trejo WH, Principe PA, and Sykes RB. (1982). SQ 27,860, a simple carbapenem produced by species of *Serratia* and *Erwinia*. *J Antibiot (Tokyo)*, 35:653-660.
- Pawlowski K. (2008). Induction of Actinorhizal Nodules by *Frankia*. *Prokar Symbion Plants*, pp. 127-154.
- Peiffer JA, Spor A, Koren O, Jin Z, Tringe SG, Dangl JL, Buckler ES, and Ley RE. (2013). Diversity and heritability of the maize rhizosphere microbiome under field conditions. *Proc Natl Acad Sci U S A*, 110:6548-6553.
- Pemberton CL, Whitehead NA, Sebaihia M, Bell KS, Hyman LJ, Harris SJ, Matlin AJ, Robson ND, Birch PR, Carr JP, Toth IK, and Salmond GPC. (2005). Novel quorum-sensing-controlled genes in *Erwinia carotovora* subsp. *carotovora*: identification of a fungal elicitor homologue in a soft-rotting bacterium. *Mol Plant Microbe Interact*, 18:343-353.
- Pérez-Tomás R, Montaner B, Llagostera E, and Soto-Cerrato V. (2003). The prodigiosins, proapoptotic drugs with anticancer properties. *Biochem Pharmacol*, 66:1447-1452.
- Perret X, Staehelin C, and Broughton WJ. (2000). Molecular basis of symbiotic promiscuity. *Microbiol Mol Biol Rev*, 64:180-201.
- Perry JJ. (1961). Prodigiosin in an actinomycete. *Nature*, pp. 77-78.
- Peters GA. (1991). *Azolla* and other plant-cyanobacteria symbioses: Aspects of form and function. *Plant Soil*, 137:25-36.
- Pham VH, and Kim J. (2012). Cultivation of unculturable soil bacteria. *Trends Biotechnol*, 30:475-484.
- Pinto UM, Pappas KM, and Winans SC. (2012). The ABCs of plasmid replication and segregation. *Nat Rev Microbiol*, 10:755-765.
- Pongsilp N, Teaumroong N, Nuntagij A, Boonkerd N, and Sadowsky MJ. (2002). Genetic structure of indigenous non-nodulating and nodulating populations of *Bradyrhizobium* in soils from Thailand. *Symbiosis*, 33(1):39-58.

- Poulter S, Carlton TM, Su X, Spring DR, and Salmond GPC. (2010). Engineering of new prodigiosin-based biosensors of *Serratia* for facile detection of short-chain N-acyl homoserine lactone quorum-sensing molecules. *Environ Microbiol Rep*, 2:322-328.
- Pritchard L, Humphris S, Baeyen S, Maes M, Van Vaerenbergh J, Elphinstone J, Saddler G, and Toth I. (2013). Draft Genome Sequences of Four *Dickeya dianthicola* and Four *Dickeya solani* Strains. *Genome Announc*, 1.
- Pueppke SG, and Broughton WJ. (1999). *Rhizobium* sp. strain NGR234 and *R. fredii* USDA257 share exceptionally broad, nested host ranges. *Mol Plant Microbe Interact*, 12:293-318.
- Queiroux C, Washburn BK, Davis OM, Stewart J, Brewer TE, Lyons MR, and Jones KM. (2012). A comparative genomics screen identifies a *Sinorhizobium meliloti* 1021 sodM-like gene strongly expressed within host plant nodules. *BMC Microbiol*, 12:74.
- Raaijmakers JM, and Weller DM. (1998). Natural Plant Protection by 2,4-Diacetylphloroglucinol-Producing *Pseudomonas* spp. in Take-All Decline Soils. *Molecular Plant-Microbe Interactions*, 11:144-152.
- Raaijmakers JM, Vlami M, and de Souza JT. (2002). Antibiotic production by bacterial biocontrol agents. *Antonie Van Leeuwenhoek*, 81:537-547.
- Radeva G, Jurgens G, Niemi M, Nick G, Suominen L, and Lindström K. (2001). Description of two biovars in the *Rhizobium galegae* species: biovar orientalis and biovar officinalis. *Syst Appl Microbiol*, 24:192-205.
- Ramírez-Bahena MH, Peix A, Rivas R, Camacho M, Rodríguez-Navarro DN, Mateos PF, Martínez-Molina E, Willems A, and Velázquez E. (2009). *Bradyrhizobium pachyrhizi* sp. nov. and *Bradyrhizobium jicamae* sp. nov., isolated from effective nodules of *Pachyrhizus erosus*. *Int J Syst Evol Microbiol*, 59:1929-1934.
- Ramírez-Bahena MH, Chahboune R, Peix A, and Velázquez E. (2013). Reclassification of *Agromonas oligotrophica* into the genus *Bradyrhizobium* as *Bradyrhizobium oligotrophicum* comb. nov. *Int J Syst Evol Microbiol*, 63:1013-1016.
- Raymond J, Siefert JL, Staples CR, and Blankenship RE. (2004). The natural history of nitrogen fixation. *Mol Biol Evol*, 21:541-554.
- Reeve W, Chain P, O'Hara G, Ardley J, Nandesena K, Brau L, Tiwari R, Malfatti S, Kiss H, Lapidus A, Copeland A, Nolan M, Land M, Hauser L, Chang YJ, Ivanova N, Mavromatis K, Markowitz V, Kyrpides N, Gollagher M, Yates R, Dilworth M, and Howieson J. (2010). Complete genome sequence of the Medicago microsymbiont *Ensifer* (*Sinorhizobium*) *medicae* strain WSM419. *Stand Genomic Sci*, 2:77-86.
- Regue M, Fabregat C, and Vinas M. (1991). A generalized transducing bacteriophage for *Serratia marcescens*. *Res Microbiol*, 142:23-27.
- Reina J, Borrell N, and Llompart I. (1992). Community-acquired bacteremia caused by *Serratia plymuthica*. Case report and review of the literature. *Diagn Microbiol Infect Dis*, 15:449-452.

- Rejili M, Lorite MJ, Mahdhi M, Pinilla JS, Ferchichi A, and Mars M. (2009). Genetic diversity of rhizobial populations recovered from three *Lotus* species cultivated in the infra-arid Tunisian soils. *Prog Nat Sci*, 19:1079-1087.
- Rice WA, Penney DC, and Nyborg M. (1977). Effects of soil acidity on rhizobia numbers, nodulation and nitrogen fixation by alfalfa and red clover. *Can J Soil Sci*, 57:197-203.
- Richter M, and Rossello-Mora R. (2009). Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci U S A*, 106:19126-19131.
- Richter M, Rossello-Mora R, Oliver Glockner F, and Peplies J. (2016). JSpeciesWS: a web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinformatics*, 32:929-931.
- Rivas R, Willems A, Palomo JL, Garcia-Benavides P, Mateos PF, Martinez-Molina E, Gillis M, and Velazquez E. (2004). *Bradyrhizobium betae* sp. nov., isolated from roots of *Beta vulgaris* affected by tumour-like deformations. *Int J Syst Evol Microbiol*, 54:1271-1275.
- Robertson GP, and Vitousek PM. (2009). Nitrogen in Agriculture: Balancing the Cost of an Essential Resource. *Ann Rev Environ Resour*, 34:97-125.
- Roth VR, Arduino MJ, Nobiletti J, Holt SC, Carson LA, Wolf CF, Lenos BA, Allison PM, and Jarvis WR. (2000). Transfusion-related sepsis due to *Serratia liquefaciens* in the United States. *Transfusion*, 40:931-935.
- Rousk J, Baath E, Brookes PC, Lauber CL, Lozupone C, Caporaso JG, Knight R, and Fierer N. (2010). Soil bacterial and fungal communities across a pH gradient in an arable soil. *ISME J*, 4:1340-1351.
- Rouws LFM, Leite J, de Matos GF, Zilli JE, Coelho MRR, Xavier GR, Fischer D, Hartmann A, Reis VM, and Baldani JL. (2014). Endophytic *Bradyrhizobium* spp. isolates from sugarcane obtained through different culture strategies. *Environ Microbiol Rep*, 6:354-363.
- Ruppel S. (1991). *Serratia rubidea* — an associative plant growth promoting nitrogen fixing microorganism. *Zentralbl Mikrobiol*, 146:297-303.
- Ryu CM, Farag MA, Hu CH, Reddy MS, Wei HX, Pare PW, and Kloepper JW. (2003). Bacterial volatiles promote growth in *Arabidopsis*. *Proc Natl Acad Sci U S A*, 100:4927-4932.
- Sabri A, Leroy P, Haubruge E, Hance T, Frère I, Destain J, and Thonart P. (2011). Isolation, pure culture and characterization of *Serratia symbiotica* sp. nov., the R-type of secondary endosymbiont of the black bean aphid *Aphis fabae*. *Int J Syst Evol Microbiol*, 61:2081-2088.
- Sadowsky MJ. (1998). Soil biology of the Rhizobiaceae. In *The Rhizobiaceae: Molecular Biology of Model Plant-Associated Bacteria*. Kluwer, Dordrecht, The Netherlands, pp. 155–172.
- Salmond GPC, and Fineran PC. (2015). A century of the phage: past, present and future. *Nat Rev Microbiol*, 13:777-786.

- Sanchez D, Mulet M, Rodriguez AC, David Z, Lalucat J, and Garcia-Valdes E. (2014). *Pseudomonas aestusnigri* sp. nov., isolated from crude oil-contaminated intertidal sand samples after the Prestige oil spill. *Syst Appl Microbiol* 37:89-94.
- Sanguin H, Sarniguet A, Gazengel K, Moënné-Loccoz Y, and Grundmann GL. (2009). Rhizosphere bacterial communities associated with disease suppressiveness stages of take-all decline in wheat monoculture. *New Phytol*, 184:694-707.
- Santner A, and Estelle M. (2009). Recent advances and emerging trends in plant hormone signalling. *Nature*, 459:1071-1078.
- Sarita S, Sharma PK, Priefer UB, and Prell J. (2005). Direct amplification of rhizobial *nodC* sequences from soil total DNA and comparison to *nodC* diversity of root nodule isolates. *FEMS Microbiol Ecol*, 54:1-11.
- Sasser M. (1990). Identification of bacteria by gas chromatography of cellular fatty acids. *MIDI Technical Note 101*. Microbial ID Inc., Newark, DE, USA. 6 pp.
- Schachtman DP, Reid RJ, and Ayling SM. (1998). Phosphorus Uptake by Plants: From Soil to Cell. *Plant Physiol*, 116:447.
- Schmidt EL, Bakole RO, and Bohlool BB. (1968). Fluorescent-antibody approach to study of rhizobia in soil. *J Bacteriol*, 95:1987-1992.
- Schmidt R, Jager V, Zuhlke D, Wolff C, Bernhardt J, Cankar K, Beekwilder J, Ijcken WV, Sleutels F, Boer W, Riedel K, and Garbeva P. (2017). Fungal volatile compounds induce production of the secondary metabolite Sodorifen in *Serratia plymuthica* PRI-2C. *Sci Rep*, 7:862.
- Schmulling T. (2002). New Insights into the Functions of Cytokinins in Plant Development. *J Plant Growth Regul*, 21:40-49.
- Schultze M, and Kondorosi A. (1998). Regulation of symbiotic root nodule development. *Annu Rev Genet*, 32:33-57.
- Schwyn B, and Neilands JB. (1987). Universal chemical assay for the detection and determination of siderophores. *Anal Biochem*, 160:47-56.
- Seemann T. (2014). Prokka: rapid prokaryotic genome annotation. *Bioinformatics*, 30:2068-2069.
- Segovia L, Piñero D, Palacios R, and Martínez-Romero E. (1991). Genetic structure of a soil population of nonsymbiotic *Rhizobium leguminosarum*. *Appl Environ Microbiol*, 57:426-433.
- Shakya M, Gottel N, Castro H, Yang ZK, Gunter L, Labbé J, Muchero W, Bonito G, Vilgalys R, Tuskan G, Podar M, and Schadt CW. (2013). A Multifactor Analysis of Fungal and Bacterial Community Structure in the Root Microbiome of Mature *Populus deltoides* Trees. *PLoS ONE*, 8:e76382.
- Shames SL, Ash DE, Wedler FC, and Villafranca JJ. (1984). Interaction of aspartate and aspartate-derived antimetabolites with the enzymes of the threonine biosynthetic pathway of *Escherichia coli*. *J Biol Chem*, 259:15331-15339.

- Shaw D P, Ping G, L. Daly S, Cha C, Cronan J, L. Rinehart K, and Farrand S. (1997). Detecting and characterizing *N*-acyl-homoserine lactone signal molecules by thin-layer chromatography. *Proc Natl Acad Sci USA*, 94(12): 6036–6041.
- Silva C, Kan FL, and Martinez-Romero E. (2007). Population genetic structure of *Sinorhizobium meliloti* and *S. medicae* isolated from nodules of *Medicago* spp. in Mexico. *FEMS Microbiol Ecol*, 60:477-489.
- Singh RP, and Jha PN. (2016). The Multifarious PGPR *Serratia marcescens* CDP-13 Augments Induced Systemic Resistance and Enhanced Salinity Tolerance of Wheat (*Triticum aestivum* L.). *PLoS ONE*, 11:e0155026.
- Singleton PW, and Tavares JW. (1986). Inoculation Response of Legumes in Relation to the Number and Effectiveness of Indigenous Rhizobium Populations. *Appl Environ Microbiol*, 51:1013-1018.
- Slater H, Crow M, Everson L, and Salmond GPC. (2003). Phosphate availability regulates biosynthesis of two antibiotics, prodigiosin and carbapenem, in *Serratia* via both quorum-sensing-dependent and -independent pathways. *Mol Microbiol*, 47:303-320.
- Smith SE, Jakobsen I, Grønlund M, and Smith FA. (2011). Roles of Arbuscular Mycorrhizas in Plant Phosphorus Nutrition: Interactions between Pathways of Phosphorus Uptake in Arbuscular Mycorrhizal Roots Have Important Implications for Understanding and Manipulating Plant Phosphorus Acquisition. *Plant Physiol*, 156:1050.
- Sneath PH, and Sokal RR. (1973). Unweighted Pair Group Method with Arithmetic Mean. ed. San Francisco: Freeman: Numerical Taxonomy, pp. 230–234.
- Solé M, Rius N, Francia A, and Lorén JG. (1994). The effect of pH on prodigiosin production by non-proliferating cells of *Serratia marcescens*. *Lett Appl Microbiol*, 19:341-344.
- Somasegaran P, and Hoben JH. (1995). Handbook for Rhizobia: Methods in Legume-Rhizobium Technology. *Q Rev Biol*, 70:224-225.
- Spaepen S, Vanderleyden J, and Remans R. (2007). Indole-3-acetic acid in microbial and microorganism-plant signaling. *FEMS Microbiol Rev*, 31:425-448.
- Stamatakis A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*, 30:1312-1313.
- Stursová M, Zifčáková L, Leigh MB, Burgess R, and Baldrian P. (2012). Cellulose utilization in forest litter and soil: identification of bacterial and fungal decomposers. *FEMS Microbiol Ecol*, 80:735-746.
- Sugawara M, Epstein B, Badgley BD, Unno T, Xu L, Reese J, Gyaneshwar P, Denny R, Mudge J, Bharti AK, Farmer AD, May GD, Woodward JE, Médigue C, Vallenet D, Lajus A, Rouy Z, Martinez-Vaz B, Tiffin P, Young ND, and Sadowsky MJ. (2013). Comparative genomics of the core and accessory genomes of 48 *Sinorhizobium* strains comprising five genospecies. *Genome Biol*, 14:R17.
- Sullivan JT, Patrick HN, Lowther WL, Scott DB, and Ronson CW. (1995). Nodulating strains of *Rhizobium loti* arise through chromosomal symbiotic gene transfer in the environment. *Proc Natl Acad Sci U S A*, 92:8985-8989.

- Sullivan JT, Eardly BD, van Berkum P, and Ronson CW. (1996). Four unnamed species of nonsymbiotic rhizobia isolated from the rhizosphere of *Lotus corniculatus*. *Appl Environ Microbiol*, 62:2818-2825.
- Sullivan JT, Trzebiatowski JR, Cruickshank RW, Gouzy J, Brown SD, Elliot RM, Fleetwood DJ, McCallum NG, Rossbach U, Stuart GS, Weaver JE, Webby RJ, De Bruijn FJ, and Ronson CW. (2002). Comparative sequence analysis of the symbiosis island of *Mesorhizobium loti* strain R7A. *J Bacteriol*, 184:3086-3095.
- Taghavi S, Garafola C, Monchy S, Newman L, Hoffman A, Weyens N, Barac T, Vangronsveld J, and van der Lelie D. (2009). Genome survey and characterization of endophytic bacteria exhibiting a beneficial effect on growth and development of poplar trees. *Appl Environ Microbiol*, 75:748-757.
- Tahir HAS, Gu Q, Wu H, Raza W, Safdar A, Huang Z, Rajer FU, and Gao X. (2017). Effect of volatile compounds produced by *Ralstonia solanacearum* on plant growth promoting and systemic resistance inducing potential of *Bacillus* volatiles. *BMC Plant Biol*, 17:133.
- Talarico TL, and Dobrogosz WJ. (1989). Chemical characterization of an antimicrobial substance produced by *Lactobacillus reuteri*. *Antimicrob Agents Chemother*, 33:674-679.
- Tamura K, Stecher G, Peterson D, Filipowski A, and Kumar S. (2013). MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Mol Biol Evol*, 30:2725-2729.
- Tan Z, Hurek T, Vinuesa P, Müller P, Ladha JK, and Reinhold-Hurek B. (2001). Specific detection of *Bradyrhizobium* and *Rhizobium* strains colonizing rice (*Oryza sativa*) roots by 16S-23S ribosomal DNA intergenic spacer-targeted PCR. *Appl Environ Microbiol*, 67:3655-3664.
- Tate RL III. (1995). Soil microbiology. New York: John Wiley and Sons. 536 pp.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, and Ostell J. (2016). NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res*, 44:6614-6624.
- Tecon R, and Or D. (2017). Cooperation in carbon source degradation shapes spatial self-organization of microbial consortia on hydrated surfaces. *Sci Rep*, 7:43726.
- Thies JE, Bohlool BB, and Singleton PW. (1991). Subgroups of the Cowpea Miscellany: Symbiotic Specificity within *Bradyrhizobium* spp. for *Vigna unguiculata*, *Phaseolus lunatus*, *Arachis hypogaea*, and *Macroptilium atropurpureum*. *Appl Environ Microbiol*, 57:1540-1545.
- Thomson NR, Crow MA, McGowan SJ, Cox A, and Salmond GPC. (2000). Biosynthesis of carbapenem antibiotic and prodigiosin pigment in *Serratia* is under quorum sensing control. *Mol Microbiol*, 36:539-556.
- Thuerig B, Fließbach A, Berger N, Fuchs JG, Kraus N, Mahlberg N, Nietlispach B, and Tamm L. (2009). Re-establishment of suppressiveness to soil- and air-borne diseases by re-inoculation of soil microbial communities. *Soil Biol Biochem*, 41:2153-2161.

- Tian CF, Zhou YJ, Zhang YM, Li QQ, Zhang YZ, Li DF, Wang S, Wang J, Gilbert LB, Li YR, and Chen WX. (2012). Comparative genomics of rhizobia nodulating soybean suggests extensive recruitment of lineage-specific genes in adaptations. *Proc Natl Acad Sci U S A*, 109:8629-8634.
- Tombolini R, van der Gaag DJ, Gerhardson B, and Jansson JK. (1999). Colonization pattern of the biocontrol strain *Pseudomonas chlororaphis* MA 342 on barley seeds visualized by using green fluorescent protein. *Appl Environ Microbiol*, 65:3674-3680.
- Tong Z, and Sadowsky MJ. (1994). A Selective Medium for the Isolation and Quantification of *Bradyrhizobium japonicum* and *Bradyrhizobium elkanii* Strains from Soils and Inoculants. *Appl Environ Microbiol*, 60:581-586.
- Toomsan B, Rupela OP, Mittal S, Dart PJ, and Clark KW. (1984). Counting *Cicer-Rhizobium* using a plant infection technique. *Soil Biol Biochem*, 16:503-507.
- Torrent J, Schwertmann U, and Schulze DG. (1980). Iron oxide mineralogy of some soils of two river terrace sequences in Spain. *Geoderma*, 23:191-208.
- Torres MJ, Bueno E, Mesa S, Bedmar EJ, and Delgado MJ. (2011). Emerging complexity in the denitrification regulatory network of *Bradyrhizobium japonicum*. *Biochem Soc Trans*, 39:284-288.
- Torsvik V, and Øvreås L. (2002). Microbial diversity and function in soil: from genes to ecosystems. *Curr Opin Microbiol*, 5:240-245.
- Trinick MJ, and Hadobas PA. (1989). Competition by *Bradyrhizobium* Strains for Nodulation of the Nonlegume *Parasponia andersonii*. *Appl Environ Microbiol*, 55:1242-1248.
- Tripathi, BM, Stegen JC, Kim M, Dong K, Adams JM, and Lee YK. (2018). Soil pH mediates the balance between stochastic and deterministic assembly of bacteria. *ISME J*, 12 (4):1072-1083.
- Turner SL, and Young JPW. (2000). The glutamine synthetases of rhizobia: phylogenetics and evolutionary implications. *Mol Biol Evol*, 17 (2):309-19.
- Ueda T, Suga Y, Yahiro N, and Matsuguchi T. (1995). Phylogeny of Sym plasmids of rhizobia by PCR-based sequencing of a nodC segment. *J Bacteriol*, 177:468-472.
- Urquiaga S, P. Xavier R, F. de Moraes R, B. Batista R, Schultz N, M. Leite J, Sá J, P. Barbosa K, S. de Resende A, Alves B, and Boddey R. (2011). Evidence from field nitrogen balance and ¹⁵N natural abundance data for the contribution of biological N₂ fixation to Brazilian sugarcane varieties. *Plant Soil*, 356, (1–2), pp. 5–21.
- Ursua P R, Unzaga M, Melero P, Iturburu I, Ezpeleta C, and Cisterna Cancer R. (1996). *Serratia rubidaea* as an invasive pathogen. *J Clin Microbiol*, 34(1): 216–217.
- van Berkum P, and Fuhrmann JJ. (2000). Evolutionary relationships among the soybean bradyrhizobia reconstructed from 16S rRNA gene and internally transcribed spacer region sequence divergence. *Int J Syst Evol Microbiol*, 50 Pt 6:2165-2172.

- van Berkum P, Leibold JM, and Eardly BD. (2006). Proposal for combining *Bradyrhizobium* spp. (*Aeschynomene indica*) with *Blastobacter denitrificans* and to transfer *Blastobacter denitrificans* (Hirsch and Muller, 1985) to the genus *Bradyrhizobium* as *Bradyrhizobium denitrificans* (comb. nov.). *Syst Appl Microbiol* 29 (3):207-15.
- van der Heijden MG, Bardgett RD, and van Straalen NM. (2008). The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecol Lett*, 11:296-310.
- van Elsas JD, Hill P, Chronáková A, Grekova M, Topalova Y, Elhottová D, and Kristůfek V. (2007). Survival of genetically marked *Escherichia coli* O157:H7 in soil as affected by soil microbial community shifts. *ISME J*, 1:204-214.
- van Elsas JD, Costa R, Jansson J, Sjöling S, Bailey M, Nalin R, Vogel TM, and van Overbeek L. (2008). The metagenomics of disease-suppressive soils - experiences from the METACONTROL project. *Trends Biotechnol*, 26:591-601.
- VanInsberghe D, Maas KR, Cardenas E, Strachan CR, Hallam SJ, and Mohn WW. (2015). Non-symbiotic *Bradyrhizobium* ecotypes dominate North American forest soils. *ISME J*, 9:2435-2441.
- Verma V, Ravindran P, and Kumar PP. (2016). Plant hormone-mediated regulation of stress responses. *BMC Plant Biol*, 16:86.
- Vincent JM. (1970). A Manual for the Practical Study of Root-nodule Bacteria: Published for the International Biological Programme by Blackwell Scientific. Oxford. 440 pp.
- Vinuesa P, León-Barrios M, Silva C, Willems A, Jarabo-Lorenzo A, Pérez-Galdona R, Werner D, and Martínez-Romero E. (2005a). *Bradyrhizobium canariense* sp. nov., an acid-tolerant endosymbiont that nodulates endemic genistoid legumes (*Papilionoideae*: Genisteae) from the Canary Islands, along with *Bradyrhizobium japonicum* bv. genistearum, *Bradyrhizobium* genospecies alpha and *Bradyrhizobium* genospecies beta. *Int J Syst Evol Microbiol*, 55:569-575.
- Vinuesa P, Silva C, Werner D, and Martínez-Romero E. (2005b). Population genetics and phylogenetic inference in bacterial molecular systematics: the roles of migration and recombination in *Bradyrhizobium* species cohesion and delineation. *Mol Phylogenet Evol*, 34:29-54.
- Vogel TM, Simonet P, Jansson JK, Hirsch PR, Tiedje JM, van Elsas JD, Bailey MJ, Nalin R, and Philippot L. (2009). TerraGenome: a consortium for the sequencing of a soil metagenome. *Nat Rev Microbiol*, 7:252.
- von Reuss SH, Kai M, Piechulla B, and Francke W. (2010). Octamethylbicyclo[3.2.1]octadienes from the rhizobacterium *Serratia odorifera*. *Angew Chem Int Ed Engl*, 49:2009-2010.
- Wall LG. (2000). The Actinorhizal Symbiosis. *J Plant Growth Regul*, 19:167-182.
- Wang ET, van Berkum P, Beyene D, Sui XH, Dorado O, Chen WX, and Martínez-Romero E. (1998). *Rhizobium huautlense* sp. nov., a symbiont of *Sesbania herbacea* that has a close phylogenetic relationship with *Rhizobium galegae*. *Int J Syst Bacteriol*, 48 Pt 3:687-699.

- Wang JY, Wang R, Zhang YM, Liu HC, Chen WF, Wang ET, Sui XH, and Chen WX. (2013). *Bradyrhizobium daqingense* sp. nov., isolated from soybean nodules. *Int J Syst Evol Microbiol*, 63:616-624.
- Wang YC, Wang F, Hou BC, Wang ET, Chen WF, Sui XH, Chen WX, Li Y, and Zhang YB. (2013). Proposal of *Ensifer psoraleae* sp. nov., *Ensifer sesbaniae* sp. nov., *Ensifer morelense* comb. nov. and *Ensifer americanum* comb. nov. *Syst Appl Microbiol*, 36:467-473.
- Wassenaar TM, Bohlin J, Binnewies TT, and Ussery DW. (2009). Genome comparison of bacterial pathogens. *Genome Dyn*, 6:1-20.
- Weber T, Blin K, Duddela S, Krug D, Kim HU, Bruccoleri R, Lee SY, Fischbach MA, Müller R, Wohlleben W, Breitling R, Takano E, and Medema MH. (2015). antiSMASH 3.0—a comprehensive resource for the genome mining of biosynthetic gene clusters. *Nucleic Acids Res*, 43:W237-W243.
- Weidner S, Becker A, Bonilla I, Jaenicke S, Lloret J, Margaret I, Puhler A, Ruiz-Sainz JE, Schneiker-Bekel S, Szczepanowski R, Vinardell JM, Zehner S, and Gottfert M. (2012). Genome sequence of the soybean symbiont *Sinorhizobium fredii* HH103. *J Bacteriol*, 194:1617-1618.
- Weisburg WG, Barns SM, Pelletier DA, and Lane DJ. (1991). 16S ribosomal DNA amplification for phylogenetic study. *J Bacteriol*, 173:697-703.
- Weise T, Thurmer A, Brady S, Kai M, Daniel R, Gottschalk G, and Piechulla B. (2014). VOC emission of various *Serratia* species and isolates and genome analysis of *Serratia plymuthica* 4Rx13. *FEMS Microbiol Lett*, 352:45-53.
- Weller DM, Raaijmakers JM, Gardener BB, and Thomashow LS. (2002). Microbial populations responsible for specific soil suppressiveness to plant pathogens. *Annu Rev Phytopathol*, 40:309-348.
- Wenke K, Kai M, and Piechulla B. (2010). Belowground volatiles facilitate interactions between plant roots and soil organisms. *Planta*, 231:499-506.
- Wenke K, Weise T, Warnke R, Valverde C, Wanke D, Kai M, and Piechulla B. (2012). Bacterial Volatiles Mediating Information Between Bacteria and Plants. *Biocommuni Plants*, Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 327-347.
- Weyens N, van der Lelie D, Taghavi S, Newman L, and Vangronsveld J. (2009). Exploiting plant-microbe partnerships to improve biomass production and remediation. *Trends Biotechnol*, 27:591-598.
- White PJ, and Brown PH. (2010). Plant nutrition for sustainable development and global health. *Ann Bot*, 105:1073-1080.
- Wilkinson S, Kudoyarova GR, Veselov DS, Arkhipova TN, and Davies WJ. (2012). Plant hormone interactions: innovative targets for crop breeding and management. *J Exp Bot*, 63:3499-3509.

- Willems A, Coopman R, and Gillis M. (2001). Phylogenetic and DNA-DNA hybridization analyses of *Bradyrhizobium* species. *Int J Syst Evol Microbiol*, 51:111-117.
- Willems A, Fernandez-Lopez M, Munoz-Adelantado E, Goris J, De Vos P, Martinez-Romero E, Toro N, and Gillis M. (2003). Description of new *Ensifer* strains from nodules and proposal to transfer *Ensifer adhaerens* Casida 1982 to *Sinorhizobium* as *Sinorhizobium adhaerens* comb. nov. Request for an opinion. *Int J Syst Evol Microbiol*, 53:1207-1217.
- Willems A. (2006). The taxonomy of rhizobia: an overview. *Plant Soil*, (287, 1/2), pp. 3-14.
- Williams RP. (1973). Biosynthesis of prodigiosin, a secondary metabolite of *Serratia marcescens*. *Appl Microbiol*, pp. 396-402.
- Williams RP, Gott CL, and Qadri SM. (1971). Induction of pigmentation in nonproliferating cells of *Serratia marcescens* by addition of single amino acids. *J Bacteriol*, 106:444-448.
- Williams RP, and Qadri SMH. (1980). The pigment of *Serratia*. In the Genus *Serratia*. Edited by A. von Graevenitz & S.J Rubin. Boca Raton, Florida: CRC Press, pp. 31-75.
- Wilson K. (2001). Preparation of genomic DNA from bacteria. *Curr Protoc Mol Biol*, Chapter 2:Unit 2.4.
- Wommack KE, and Colwell RR. (2000). Virioplankton: viruses in aquatic ecosystems. *Microbiol Mol Biol Rev*, 64:69-114.
- Woodmansee RG, Dodd JL, Bowman RA, Clark FE, and Dickinson CE. (1978). Nitrogen budget of a shortgrass prairie ecosystem. *Oecologia*, 34:363-376.
- Wu DY, Jospin G, and Eisen JA. (2013). Systematic Identification of Gene Families for Use as "Markers" for Phylogenetic and Phylogeny-Driven Ecological Studies of Bacteria and Archaea and Their Major Subgroups. *PLoS ONE*, 8(10):e77033.
- Wust PK, Nacke H, Kaiser K, Marhan S, Sikorski J, Kandeler E, Daniel R, and Overmann J. (2016). Estimates of Soil Bacterial Ribosome Content and Diversity Are Significantly Affected by the Nucleic Acid Extraction Method Employed. *Appl Environ Microbiol*, 82:2595-2607.
- Xu LM, Ge C, Cui Z, Li J, and Fan H. (1995). *Bradyrhizobium liaoningense* sp. nov., isolated from the root nodules of soybeans. *Int J Syst Bacteriol*, 45:706-711.
- Yang Y, Wang N, Guo X, Zhang Y, and Ye B. (2017). Comparative analysis of bacterial community structure in the rhizosphere of maize by high-throughput pyrosequencing. *PLoS ONE*, 12:e0178425.
- Yao Y, Sui XH, Zhang XX, Wang ET, and Chen WX. (2015). *Bradyrhizobium erythrophlei* sp. nov. and *Bradyrhizobium ferriligni* sp. nov., isolated from effective nodules of *Erythrophleum fordii*. *Int J Syst Evol Microbiol*, 65:1831-1837.
- Yao ZY, Kan FL, Wang ET, Wei GH, and Chen WX. (2002). Characterization of rhizobia that nodulate legume species of the genus *Lespedeza* and description of *Bradyrhizobium yuanmingense* sp. nov. *Int J Syst Evol Microbiol*, 52:2219-2230.

- Young JPW, and Wexler M. (1988). Sym plasmid and chromosomal genotypes are correlated in field populations of *Rhizobium leguminosarum*. *J Gen Microbiol*, 134.
- Young JM. (2003). The genus name *Ensifer* Casida 1982 takes priority over *Sinorhizobium* Chen *et al.* 1988, and *Sinorhizobium morelense* Wang *et al.* 2002 is a later synonym of *Ensifer adhaerens* Casida 1982. Is the combination "*Sinorhizobium adhaerens*" (Casida 1982) Willems *et al.* 2003 legitimate? Request for an Opinion. *Int J Syst Evol Microbiol*, 53:2107-2110.
- Yu X, Cloutier S, Tambong JT, and Bromfield ES. (2014). *Bradyrhizobium ottawaense* sp. nov., a symbiotic nitrogen fixing bacterium from root nodules of soybeans in Canada. *Int J Syst Evol Microbiol*, 64:3202-3207.
- Zézé A, Mutch Lesley A, and Young JPW. (2001). Direct amplification of *nodD* from community DNA reveals the genetic diversity of *Rhizobium leguminosarum* in soil. *Environ Microbiol*, 3:363-370.
- Zhang X-X, Kosier B, and Priefer UB. (2001). Symbiotic Plasmid Rearrangement in *Rhizobium leguminosarum* bv. *viciae* VF39SM. *J Bacteriol*, 183:2141-2144.
- Zhang YM, Li Y, Chen WF, Wang ET, Sui XH, Li QQ, Zhang YZ, Zhou YG, and Chen WX. (2012). *Bradyrhizobium huanghuaihaiense* sp. nov., an effective symbiotic bacterium isolated from soybean (*Glycine max* L.) nodules. *Int J Syst Evol Microbiol*, 62:1951-1957.
- Zhu J, Chai Y, Zhong Z, Li S, and Winans SC. (2003). Agrobacterium Bioassay Strain for Ultrasensitive Detection of N-Acylhomoserine Lactone-Type Quorum-Sensing Molecules: Detection of Autoinducers in *Mesorhizobium huakuii*. *Appl Environ Microbiol*, 69:6949-6953.

SUPPLEMENTAL TABLES AND FIGURES

Supplemental Table S1. *Neorhizobium*, *Ensifer*, *Bradyrhizobium* and *Serratia* Genbank and JGI (**bold**) Accession numbers.

<i>Neorhizobium</i> accession numbers				
Strain	16S rDNA	<i>rpoB</i>	Genome	
<i>Neorhizobium</i> sp. T9_24	MG957216	MG966473		
<i>Neorhizobium</i> sp. T6_25	MG957217	MG966474	PRJNA434367	
<i>Neorhizobium</i> sp. T13_2	MG957218	MG966475		
<i>Neorhizobium</i> sp. T7_12	MG957219	MG966476	PRJNA434366	
<i>Neorhizobium</i> sp. T6_21	MG957220	MG966477		
<i>Neorhizobium</i> sp. T6_1	MG957221	MG966478		
<i>Neorhizobium</i> sp. T20_22	MG957222	MG966479	PRJNA434360	
<i>Neorhizobium</i> sp. T25_27	MG957223	MG966480	PRJNA434365	
<i>Neorhizobium</i> sp. T4_8	MG957224	MG966481		
<i>Neorhizobium</i> sp. T4_1	MG957225	MG966482		
<i>Neorhizobium</i> sp. T25_7	MG957226	MG966483		
<i>Neorhizobium</i> sp. T25_13	MG957227	MG966484	PRJNA434364	
<i>Neorhizobium</i> sp. T21_15	MG957228	MG966485		
<i>Neorhizobium</i> sp. T24_19	MG957229	MG966486		
<i>Neorhizobium</i> sp. T7_7	MG957230	MG966487		
<i>Neorhizobium</i> sp. T16_9	MG957231	MG966488		
<i>Neorhizobium</i> sp. T5_2	MG957232	MG966489		
<i>Neorhizobium</i> sp. T7_11	MG957233	MG966490		
<i>Neorhizobium</i> sp. T22_47	MG957234	MG966491		
<i>Neorhizobium</i> sp. T23_12	MG957235	MG966492		
<i>Neorhizobium</i> sp. T17_20 ^T			PRJNA434362	
<i>Neorhizobium</i> sp. T11_12			PRJNA434363	
<i>Neorhizobium galegae</i> HAMBI 540 ^T			PRJNA224116	
<i>Neorhizobium alkanisoli</i> DSM 21817 ^T			PRJNA434370	
<i>Neorhizobium huautlense</i> DSM 21826 ^T			PRJNA434371	

Supplemental Tables and Figures

<i>Neorhizobium alkanisoli</i> CCBAU 01393 ^T	EU074168.1	KF278573.1			
<i>Neorhizobium huautlense</i> LMH 18254 ^T	AF025852.1	EF217309.1			
<i>Agrobacterium tumefaciens</i> NCPPB 2437 ^T	NR_115516.1	JN580757.1			
<i>Rhizobium etli</i> CFN 42 ^T			NC_007761.1		
<i>Rhizobium leguminosarum</i> bv. <i>viciae</i> 2370 ^T			PRJNA450070		
<i>Agrobacterium tumefaciens</i> Ach5			PRJNA278497		
Ensifer accession numbers					
Strain	16S rDNA	<i>glnII</i>	<i>recA</i>	<i>asmA</i>	Genome
<i>Ensifer</i> sp. TMN_1	MG230474	MG270002	MG270022		
<i>Ensifer</i> sp. TMN_12	MG230475	MG270003	MG270023		
<i>Ensifer</i> sp. TMN_13	MG230476	MG270004	MG270024		
<i>Ensifer</i> sp. TMN_14	MG230477	MG270005	MG270025		
<i>Ensifer</i> sp. TMN_25	MG230478	MG270006	MG270026		
<i>Ensifer</i> sp. TMN_26	MG230479	MG270007	MG270027		
<i>Ensifer</i> sp. MMN_1	MG230485	MG270008	MG270028		
<i>Ensifer</i> sp. MMN_3	MG230486	MG270009	MG270029		
<i>Ensifer</i> sp. MMN_5	MG230487	MG270010	MG270030		PRJNA407431
<i>Ensifer</i> sp. MMN_6	MG230488	MG270011	MG270031		
<i>Ensifer</i> sp. MMN_7	MG230489	MG270012	MG270032		
<i>Ensifer</i> sp. M_4_50	MG230484	MG270013	MG270041		
<i>Ensifer</i> sp. M_4_45	MG230483	MG270014	MG270033		PRJNA407429
<i>Ensifer</i> sp. M_4_46	MG230482	MG270015	MG270034		
<i>Ensifer</i> sp. M_5_14	MG230481	MG270016	MG270035		
<i>Ensifer</i> sp. M7_2	MG230480	MG270017	MG270036		
<i>Ensifer</i> sp. T1_27	MG230473	MG270018	MG270037	MG270055	
<i>Ensifer</i> sp. T1_45	MG230472	MG270019	MG270038	MG270056	

<i>Ensifer</i> sp. T1_100	MG230471	MG270020	MG270039	MG270059	
<i>Ensifer</i> sp. T2_8	MG230470	MG270021	MG270040	MG270060	
<i>Ensifer</i> sp. T1_71				MG270057	
<i>Ensifer</i> sp. T1_81				MG270058	
<i>Ensifer</i> sp. M4_29				MG270061	
<i>Ensifer meliloti</i> 2011				MG270042	AGG73298
<i>Ensifer meliloti</i> RMO 17				MG270043	AIL98543
<i>Ensifer meliloti</i> BL 225C				MG270044	AEG03270
<i>Ensifer meliloti</i> SM 11				MG270045	AEH77649
<i>Ensifer meliloti</i> Rm 41				MG270046	CCM66388
<i>Ensifer meliloti</i> GR 4				MG270047	AGA05677
<i>Ensifer meliloti</i> AK 83				MG270048	AEG52184
<i>Ensifer medicae</i> WSM 419				MG270049	ABR59173
<i>Ensifer fredii</i> NGR 234				MG270050	ACP24130
<i>Ensifer americanum</i> CFNEI 156 ^T				MG270051	
<i>Ensifer fredii</i> USDA 191				MG270052	
<i>Ensifer xinjiangense</i> LMG 17930 ^T				MG270053	
<i>Ensifer arboris</i> LMG 14919 ^T				MG270054	
<i>Ensifer meliloti</i> 1021					PRJNA19 /CAC45290
<i>Ensifer adhaerens</i> A ^T	CP015880	HM997090	HM997099		
<i>Ensifer americanum</i> CFNEI 156 ^T	NR_025251	GU994065	DQ411943		
<i>Ensifer arboris</i> TTR 38 ^T	NR_037001	HM997093	DQ411946		
<i>Ensifer fredii</i> USDA 205 ^T	X67231	2731053067	AJ294379		
<i>Ensifer kostiensis</i> TTR 15 ^T	NR_104565	GU994063	DQ411947		
<i>Ensifer kummerowiae</i> AS 1.3046 ^T	NR_042720	GU994062	DQ411945		
<i>Ensifer medicae</i> USDA 1037 ^T	NR_104719	KR818746	AM182135		

Supplemental Tables and Figures

<i>Ensifer meliloti</i> USDA 1002 ^T	D14509	KR818748	AJ294382				
<i>Ensifer mexicanus</i> ITTG_R7 ^T	NR_115768	GU994064	DQ411951				
<i>Ensifer morelensis</i> Lc04 ^T	AY024335	EU170586	AJ505601				
<i>Ensifer numidicus</i> ORS 1407 ^T	AY500254	HM997092	HM346610				
<i>Ensifer psoraleae</i> CCBAU 65732 ^T	EU618039	EU618006	EU622106				
<i>Ensifer saheli</i> ORS 609 ^T	X68390	AF169589	AM182138				
<i>Ensifer sesbaniae</i> CCBAU 65729 ^T	JF834143	JX028809	JX028823				
<i>Ensifer sojae</i> CCBAU 05684 ^T	GU593061	2657683547	2657683956				
<i>Ensifer terangae</i> ATCC 51692 ^T	X68388	AF169590	AJ294383				
<i>Ensifer alkalisoli</i> YIC4027 ^T	KX668272	KR154013	KR153998				
<i>Ensifer glycinis</i> CCBAU 23380 ^T	KJ556384	KJ556511	KJ556427				
<i>Ensifer xinjiangense</i> CCBAU 110 ^T	EU152397	EU155088	DQ411944				
Bradyrhizobium accession numbers							
Strain	16S rDNA	<i>glnII</i>	<i>recA</i>	<i>nifH</i>	<i>nodC</i>	<i>ThrC</i>	Genome
<i>Bradyrhizobium</i> sp. BSN_1	MH086226	MH091838	MH091823	MH091851	MH091854		
<i>Bradyrhizobium</i> sp. BSN_2	MH086227	MH091832	MH091820	MH091847	MH091855		
<i>Bradyrhizobium</i> sp. BSN_4	MH086228	MH091831	MH091817		MH091871		
<i>Bradyrhizobium</i> sp. BSN_5	MH086229	MH091830	MH091818				
<i>Bradyrhizobium</i> sp. BSN_6	MH086230	MH091829	MH091819				
<i>Bradyrhizobium</i> sp. BSN_9	MH086232	MH091828	MH091821	MH091849	MH091858		
<i>Bradyrhizobium</i> sp. BSN_10	MH086233	MH091833	MH091822				
<i>Bradyrhizobium</i> sp. BSN_11				MH091848	MH091859		
<i>Bradyrhizobium</i> sp. BSN_3					MH091856		
<i>Bradyrhizobium</i> sp. BSN_8					MH091857		
<i>Bradyrhizobium</i> sp. BSN_12					MH091860		
<i>Bradyrhizobium</i> sp. BSN_7				MH091850			

<i>Bradyrhizobium</i> sp. BLN_1				MH091843	MH091861
<i>Bradyrhizobium</i> sp. BLN_10				MH091844	MH091869
<i>Bradyrhizobium</i> sp. BLN_2				MH091845	MH091862
<i>Bradyrhizobium</i> sp. BLN_9	MH086231	MH091834	MH091827		MH091868
<i>Bradyrhizobium</i> sp. BLN_3	MH086234	MH091837	MH091824	MH091846	MH091863
<i>Bradyrhizobium</i> sp. BLN_4	MH086235	MH091836	MH091825		MH091864
<i>Bradyrhizobium</i> sp. BLN_5					MH091865
<i>Bradyrhizobium</i> sp. BLN_6					MH091866
<i>Bradyrhizobium</i> sp. BLN_8					MH091867
<i>Bradyrhizobium</i> sp. BLN_11					MH091870
<i>Bradyrhizobium</i> sp. BLN_7	MH086236	MH091835	MH091826		
<i>Bradyrhizobium</i> sp. B2_4	MH086237	MH091841	MH091813	MH091852	MH091916
<i>Bradyrhizobium</i> sp. B2_9				MH091853	MH091915
<i>Bradyrhizobium</i> sp. B3_1	MH086238	MH091840	MH091816		MH091914
<i>Bradyrhizobium</i> sp. B2_2	MH086239	MH091842	MH091815		MH091917
<i>Bradyrhizobium</i> sp. B10_118	MH086240	MH091839	MH091814		MH091918
<i>Bradyrhizobium</i> sp. B3_2					MH091913
<i>Bradyrhizobium</i> sp. P.22 ^a					MH091905
<i>Bradyrhizobium diazoefficiens</i> USDA 110 ^T					MH091906
<i>Bradyrhizobium betae</i> LMG 21987 ^T					MH091907
<i>Bradyrhizobium yuanmingense</i> LMG 21827 ^T					MH091908
<i>Bradyrhizobium japonicum</i> USDA 6 ^T					MH091909
<i>Bradyrhizobium liaonigense</i> LMG 18230 ^T					MH091910
<i>Bradyrhizobium elkanii</i> USDA 76 ^T					MH091911
<i>Bradyrhizobium</i> sp. Lmj C					MH091912

Supplemental Tables and Figures

<i>Bradyrhizobium japonicum</i> USDA 6 ^T	MH091872	2512000449
<i>Bradyrhizobium japonicum</i> 22	MH091874	2529388226
<i>Bradyrhizobium</i> sp. CCGE-LA001	MH091875	2520791477
<i>Bradyrhizobium</i> sp. ORS 375	MH091876	2519431430
<i>Bradyrhizobium elkanii</i> USDA 76 ^T	MH091877	2517892959
<i>Bradyrhizobium japonicum</i> USDA 124	MH091878	2517101859
<i>Bradyrhizobium</i> genosp. SA-4 CB756	MH091879	2516908538
<i>Bradyrhizobium</i> sp. WSM4349	MH091880	2515626630
<i>Bradyrhizobium</i> sp. STM 3809	MH091881	2514442327
<i>Bradyrhizobium</i> sp. ORS285	MH091882	2514301231
<i>Bradyrhizobium</i> sp. S23321	MH091883	2514090323
<i>Bradyrhizobium</i> sp. WSM2793	MH091884	2514010950
<i>Bradyrhizobium</i> sp. WSM2254	MH091885	2513974572
<i>Bradyrhizobium japonicum</i> USDA 38	MH091886	2513939974
<i>Bradyrhizobium elkanii</i> USDA 3254	MH091887	2513917995
<i>Bradyrhizobium</i> sp. TV2a.2	MH091888	2513892825
<i>Bradyrhizobium japonicum</i> USDA 4	MH091889	2513876618
<i>Bradyrhizobium elkanii</i> USDA 94	MH091890	2513857999
<i>Bradyrhizobium</i> sp. USDA 3384	MH091891	2513848661
<i>Bradyrhizobium</i> sp. EC3.3	MH091892	2513717269
<i>Bradyrhizobium japonicum</i> USDA 135	MH091893	2513700548
<i>Bradyrhizobium elkanii</i> WSM1741	MH091894	2513691961
<i>Bradyrhizobium elkanii</i> WSM2783	MH091895	2513677858
<i>Bradyrhizobium elkanii</i> USDA 3259	MH091896	2513656810
<i>Bradyrhizobium japonicum</i> USDA 122	MH091897	2513646802
<i>Bradyrhizobium</i> sp. WSM3983	MH091898	2513637786
<i>Bradyrhizobium</i> sp. WSM1743	MH091899	2513625691
<i>Bradyrhizobium</i> sp. YR681	MH091900	2511399952

<i>Bradyrhizobium</i> sp. ARR65						MH091901	2509147435
<i>Bradyrhizobium</i> sp. WSM1253						MH091902	2508690969
<i>Bradyrhizobium</i> sp. WSM471						MH091903	2508543282
<i>Bradyrhizobium</i> sp. WSM1417						MH091904	2507509214
	16S rDNA	<i>glnII</i>	<i>recA</i>	<i>nifH</i>	<i>nodC</i>	Genome	
<i>Bradyrhizobium betae</i> PL7HG1 ^T	AY372184.1	FJ970431.1	FM253174.1				
<i>Bradyrhizobium canariense</i> BTA-1 ^T	AJ558025.1	AY386765.1	KF532942.1	640560650	AJ560653.1		
<i>Bradyrhizobium cytisi</i> CTAW11 ^T	EU561065.2	GU001594.1	LM994316.1	GU001618.1	EU597844.2		
<i>Bradyrhizobium daqingense</i> CCBAU 15774 ^T	HQ231274.1	HQ231301.1	LM994320.1	2596848783	HQ231326.1		
<i>Bradyrhizobium denitrificans</i> ATCC 43295 ^T	X66025.1	HM047121.1	EU665419.1	HM047125.1			
<i>Bradyrhizobium diazoefficiens</i> USDA 110 ^T	NC_004463.1	CP011360.1	CP011360.1	2688593839	2688593630	CP011360.1	
<i>Bradyrhizobium elkanii</i> USDA 76 ^T	U35000.3	AY599117.1	KF532941.1	2517890319	2517890664		
<i>Bradyrhizobium embrapense</i> SEMIA 6208	AY904773.1	GQ160500.1	HQ634899.1	2636528215	2636530411		
<i>Bradyrhizobium erythrophlei</i> CCBAU 53325	KF114645.1	KF114693.1	KF114669.1	KF114598.1	KF114576.1		
<i>Bradyrhizobium ferriligni</i> CCBAU 51502	KJ818096.1	KJ818099.1	KJ818112.1	KJ818108.1	KJ818109.1		
<i>Bradyrhizobium ganzhouense</i> RITF806 ^T	JQ796661.2	JX277110.1	JX277144.1	JX292065.1	JX292035.1		
<i>Bradyrhizobium guangdongense</i> CCBAU 51649 ^T	KC508867.1	KC509023.1	KC509269.1	KC509130.1			
<i>Bradyrhizobium huanghuaihaiense</i> CCBAU 23303 ^T	HQ231463.1	HQ231639.1	LM994321.1	2596983282	2596982408		
<i>Bradyrhizobium icense</i> LMTR 13 ^T	KF896156.1	CP016428.1	JX943615.1	KF896161.1	2723844490		
<i>Bradyrhizobium ingae</i> BR 10250 ^T	KF927043.1	KF927067.1	KF927061.1	KF927085.1	KF927054.1		
<i>Bradyrhizobium iriomotense</i> EK05 ^T	AB300992.1	AB300995.1	LM994319.1	AB300998.1	AB301000.1		
<i>Bradyrhizobium japonicum</i> USDA 6 ^T	U69638.3	HQ587875.1	AM168341.1	2513670276	2513670242		

Supplemental Tables and Figures

<i>Bradyrhizobium jicamae</i> PAC68 ^T	AY624134.1	FJ428204.1	LM994324.1	2733955171	AB573869.1
<i>Bradyrhizobium kavangense</i> 14-3 ^T	KP899562.1	KM378446.1	KM378399.1	KM378254.1	KT033402.1
<i>Bradyrhizobium lablabi</i> CCBAU 23086 ^T	GU433448.1	GU433498.1	LM994322.1	2733978064	2733974194
<i>Bradyrhizobium liaoningense</i> 2281 ^T	AF208513.1	AY386775.1	FM253180.1	EU818925.1	GU263466.1
<i>Bradyrhizobium lupini</i> USDA 3051 ^T	KM114861.1	KM114862.1	KM114866.1	LC107546.1	KM114864.1
<i>Bradyrhizobium manausense</i> BR3351 ^T	HQ641226.2	KF785986.1	KF785992.1	2647449812	2647453763
<i>Bradyrhizobium neotropale</i> BR 10247 ^T	KF927051.1	KJ661700.1	KJ661714.1	KJ661728.1	KJ661727.1
<i>Bradyrhizobium oligotrophicum</i> ATCC 43045 ^T	JQ619230.1	JQ619233.1	JQ619231.1	AP012603.1	
<i>Bradyrhizobium ottawaense</i> OO99 ^T	JN186270.1	HQ587750.1	HQ587287.1	JN186287.1	HQ587980.1
<i>Bradyrhizobium pachyrhizi</i> PAC48 ^T	AY624135.1	FJ428201.1	LM994323.1	2655285856	2655287255
<i>Bradyrhizobium paxllaeri</i> LMTR 21 ^T	AY923031.1	KF896169.1	JX943617.1	DQ085619.2	KF896160.1
<i>Bradyrhizobium retamae</i> Ro19 ^T	KC247085.1	KC247108.1	LM994318.1	KF806460.1	2733970641
<i>Bradyrhizobium rifense</i> CTAW71 ^T	EU561074.2	GU001604.1	LM994317.1	GU001627.1	EU597853.2
<i>Bradyrhizobium stylosanthis</i> BR 446 ^T	KU724142.1	KU724148.1	KU724163.1	KU724157.1	KU724160.1
<i>Bradyrhizobium subterraneum</i> 58 2-1 ^T	KP308152.1	KM378484.1	KM378397.1	KM378289.1	
<i>Bradyrhizobium tropiciagri</i> SEMIA 6148 ^T	AY904753.1	FJ391048.1	FJ391168.1	HQ259540.1	2641671636
<i>Bradyrhizobium valentinum</i> LmjM3 ^T	NR_125638.1	JX518575.1	JX518589.2	2733942381	2733941890
<i>Bradyrhizobium viridifuturi</i> SEMIA 690 ^T	FJ025107.1	KR149131.1	KR149140.1	2655418735	2655417808
<i>Bradyrhizobium yuanmingense</i> CCBAU 10071 ^T	AF193818.1	AY386780.1	AY591566.1	2617379895	2617379339
<i>Bradyrhizobium vignae</i> 7-2 ^T	NR_147716.1	KM378443.1	KM378374.1	KM378251.1	KT362339.1
<i>Bradyrhizobium guangxiense</i> CCBAU 53363 ^T	NR_145894.1	KC509033.1	KC509279.1	KC509140.1	
<i>Bradyrhizobium arachidis</i> CCBAU 051107 ^T	NR_117791.1	HM107251.1	KF962707.1	2657053881	2657052964
<i>Bosea thiooxidans</i> DSM 9653 ^T	NR_041994.1				

<i>Ensifer meliloti</i> 1021	PRJNA19	PRJNA19	PRJNA19
<i>Rhizobium leguminosarum</i> bv. <i>viciae</i> 3841		639650755	
<i>Neorhizobium galegae</i> HAMBI 540 ^T		2586004227	

***Serratia* accession numbers**

Strain	16S rDNA
<i>Serratia</i> sp. T5_3	MH086001
<i>Serratia</i> sp. T4_16	MH086002
<i>Serratia</i> sp. T1_1	MH086003
<i>Serratia</i> sp. T2_10	MH086004
<i>Serratia</i> sp. T1_12	MH086005
<i>Serratia</i> sp. T1_4	MH086006
<i>Serratia</i> sp. T1_6	MH086007
<i>Serratia</i> sp. T1_11	MH086008
<i>Serratia</i> sp. T1_13	MH086009
<i>Serratia</i> sp. T1_16	MH086010
<i>Serratia</i> sp. T1_17	MH086011
<i>Serratia</i> sp. T2_2	MH086012
<i>Serratia</i> sp. T4_8	MH086013
<i>Serratia</i> sp. T4_36	MH086014
<i>Serratia</i> sp. T5_7	MH086015
<i>Serratia</i> sp. B1_113	MH086016
<i>Serratia</i> sp. B1_115	MH086017
<i>Serratia</i> sp. B1_6	MH086018
<i>Serratia</i> sp. B1_10	MH086019
<i>Serratia</i> sp. B1_42	MH086020
<i>Serratia</i> sp. B1_62	MH086021
<i>Serratia</i> sp. M3_2	MH086022
<i>Serratia</i> sp. M3_16	MH086023

Supplemental Tables and Figures

<i>Serratia</i> sp. M2_3	MH086024
<i>Serratia aquatilis</i> 2015-2462-01 ^T	KT387999.1
<i>Serratia entomophila</i> DMS12358 ^T	NR_025338.1
<i>Serratia ficaria</i> DSM 4569 ^T	NR_041979.1
<i>Serratia fonticola</i> DSM 4576 ^T	NR_025339.1
<i>Serratia glossinae</i> C1 ^T	FJ790328.1
<i>Serratia grimesii</i> 30063 ^T	AJ233430.1
<i>Serratia liquefaciens</i> CIP 103238 ^T	AJ306725.1
<i>Serratia marcescens</i> DSM 30121 ^T	AJ233431.1
<i>Serratia marinorubra</i> JCM 1240 ^T	AB004751.1
<i>Serratia myotis</i> 12 ^T	KJ739884.1
<i>Serratia nematodiphila</i> DZ0503SBS1 ^T	EU036987.1
<i>Serratia odorifera</i> DSM 4582 ^T	AJ233432.1
<i>Serratia plymuthica</i> DSM 4540 ^T	AJ233433.1
<i>Serratia proteamaculans</i> DSM 4543 ^T	AJ233434.1
<i>Serratia quinivorans</i> DSM 4597 ^T	AJ233435.1
<i>Serratia rubidaea</i> JCM 1240 ^T	AB004751.1
<i>Serratia symbiotica</i> CWBI-2.3 ^{TS1}	GU394001.1
<i>Serratia ureilytica</i> NiVa 51 ^T	AJ854062.1
<i>Serratia vespertilionis</i> 52 ^T	KJ739885.1
<i>Serratia marcescens</i> subsp. <i>sakuensis</i> ^T	AB061685.1
<i>Xenorhabdus nematophila</i> DSM 3370 ^T	AY278674.1

Supplemental Table S2. *Neorhizobium*, *Ensifer*, *Bradyrhizobium* and *Serratia* genotypes.

<i>Neorhizobium</i> sp. <i>rpoB</i> genotypes from Tomejil soil				
N° of genotype	Genotype representative	N° of isolates	Isolates	Isolation
1	T4_1	2	T4_1	Soil
			T18_15	
2	T4_8	2	T4_8	
			T16_4	
3	T5_2	1	T5_2	
4	T25_27	4	T5_26	
			T25_28	
			T25_27	
			T22_7	
5	T6_1	2	T6_1	
			T17_6	
6	T6_21	3	T6_21	
			T22_11	
			T17_15	
7	T6_25	4	T8_5	
			T6_25	
			T6_23	
			T25_30	
8	T7_12	6	T7_12	
			T25_20	
			T23_26	
			T21_19	
			T21_1	
9	T7_7	1	T20_25	
			T7_7	
10	T7_11	1	T7_11	
11	T9_24	2	T9_24	
			T20_10	
12	T20_22	12	T5_27	
			T28_6	
			T25_19	
			T24_25	
			T20_22	
			T17_4	
			T17_26	
			T17_20	
			T17_14	

			T16_2
			T16_12
			T11_12
13	T13_2	3	T7_9
			T7_1
			T13_2
14	T16_9	1	T16_9
15	T21_15	1	T21_15
16	T22_47	1	T22_47
17	T23_12	1	T23_12
18	T24_19	1	T24_19
19	T25_7	2	T25_7
			T16_1
20	T25_13	3	T25_5
			T25_4
			T25_13

***Ensifer* sp. 16S rDNA genotypes from Tomejil and Motril soils**

N° of genotype	Genotype representative	N° of isolates	Isolates	Isolation
1	TMN_1	21	TMN_1	Nodule
			TMN_2	
			TMN_21	
			TMN_15	
			TMN_3	
			TMN_4	
			TMN_5	
			TMN_6	
			TMN_7	
			TMN_8	
			TMN_9	
			TMN_10	
			TMN_17	
			TMN_18	
			TMN_19	
			TMN_20	
			TMN_22	
			TMN_23	
			TMN_28	
			TMN_31	
			TMN_27	
2	TMN_12	1	TMN_12	
3	TMN_13	1	TMN_13	

4	TMN_14	1	TMN_14	Soil
5	TMN_25	1	TMN_25	
6	TMN_26	1	TMN_26	
1	MMN_1	4	MMN_1	
			MMN_2	
			MMN_10	
			MMN_8	
2	MMN_3	1	MMN_3	
3	MMN_5	1	MMN_5	
4	MMN_7	1	MMN_7	
5	MMN_6	1	MMN_6	
1	T_1_100	7	T_1_100	
			T_3_13	
			T_2_6	
			T_2_53	
			T_2_4	
			T_2_50	
			T1_77	
2	T_1_45	1	T_1_45	
3	T_2_8	1	T_2_8	
4	T1_77	1	T1_77	
5	T1_81	1	T1_81	
6	T_1_27	1	T_1_27	
1	M_5_5	7	M_4_29	
			M_4_50	
			M_5_5	
			M_5_38	
			M_5_45	
			M8_16	
			M8_18	
2	M7_2	2	M7_2	
			M7_3	
3	M5_60	2	M5_60	
			M5_50	
4	M_5_14	1	M_5_14	
5	M_4_45	1	M_4_45	
6	M_4_46	1	M_4_46	
Bradyrhizobium sp. 16S rDNA genotypes from Burguillos soil				
Nº of genotype	Genotype representative	Nº of isolates	Isolates	Isolation
1	BSN_5	4	BSN_5 BSN_12	Nodule

Supplemental Tables and Figures

			BSN_11	
			BSN_7	
			BLN_9	
			BLN_5	
2	BLN_9	6	BLN_2	
			BLN_11	
			BLN_1	
			BLN_6	
3	BSN_4	2	BSN_4	
			BSN_8	
4	BSN_6	1	BSN_6	
5	BSN_2	1	BSN_2	
6	BSN_9	1	BSN_9	
7	BSN_10	1	BSN_10	
8	BSN_1	1	BSN_1	
9	BLN_3	1	BLN_3	
10	BLN_4	1	BLN_4	
11	BLN_7	1	BLN_7	
1	B2_4	2	B3_2	Soil
			B2_4	
2	B3_1	2	B3_1	
			B2_9	
3	B2_2	1	B2_2	
4	B10_118	1	B10_118	
Bradyrhizobium sp. nodC genotypes from Burguillos soil				
Nº of genotype	Genotype representative	Nº of isolates	Isolates	Isolation
1	BSN_2	3	BSN_2	Nodule
			BSN_5	
			BSN_10	
2	BSN_3	1	BSN_3	
3	BSN_4	1	BSN_4	
4	BSN_1	1	BSN_1	
5	BSN_8	1	BSN_8	
6	BSN_9	1	BSN_9	
7	BSN_11	1	BSN_11	
8	BSN_12	1	BSN_12	
9	BLN_1	1	BLN_1	
10	BLN_2	1	BLN_2	
11	BLN_3	1	BLN_3	
12	BLN_4	1	BLN_4	
13	BLN_5	1	BLN_5	

14	BLN_6	1	BLN_6
15	BLN_8	1	BLN_8
16	BLN_9	1	BLN_9
17	BLN_10	1	BLN_10
18	BLN_11	1	BLN_11

***Bradyrhizobium* sp. *nifH* genotypes from Burguillos soil**

N° of genotype	Genotype representative	N° of isolates	Isolates	Isolation
1	BLN_1	7	BLN_1	Nodule
			BLN_4	
			BLN_5	
			BLN_6	
			BLN_7	
			BLN_9	
			BLN_11	
2	BLN_10		BLN_10	
3	BLN_2		BLN_2	
4	BLN_3		BLN_3	
5	BSN_2	6	BSN_2	
			BSN_3	
			BSN_4	
			BSN_5	
			BSN_6	
6	BSN_11		BSN_10	
			BSN_11	
			BSN_12	
7	BSN_9		BSN_9	
8	BSN_7		BSN_7	
9	BSN_1		BSN_1	
1	B2_9	1	B2_9	Soil
2	B2_4	1	B2_4	

***Serratia* 16S rDNA genotypes from Tomejil, Motril and Burguillos soil**

Soil	N° of genotype	Genotype representative	N° of isolates	Isolates
Tomejil	1	T_2_10	20	T_2_10
				T_2_6
				T_3_10
				T_3_11
				T_3_13
				T_3_19
				T_3_7
				T_3_18

				T_3_9
				T_2_9
				T_3_16
				T_3_5
				T_3_23
				T_3_21
				T_3_20
				T_3_17
				T_3_15
				T_3_14
				T_2_5
				T_3_12
				T_4_16
				T_4_28
				T_4_27
				T_4_13
				T_4_41
2	T_4_16	11		T_4_12
				T_4_35
				T_4_34
				T_4_39
				T_4_5
				T_4_22
				T_1_1
				T_1_18
				T_1_14
				T_1_10
3	T_1_1	9		T_1_5
				T_1_9
				T_1_3
				T_1_7
				T_1_2
				T_1_12
				T_1_8
				T_3_2
4	T_1_12	7		T_3_4
				T_2_12
				T_3_1
				T_1_15
				T_5_3
5	T_5_3	3		T_4_15

				T_5_1
	6	T_1_4	1	T_1_4
	7	T_1_6	1	T_1_6
	8	T_1_11	1	T_1_11
	9	T_1_13	1	T_1_13
	10	T_1_16	1	T_1_16
	11	T_1_17	1	T_1_17
	12	T_2_2	1	T_2_2
	13	T_4_8	1	T_4_8
	14	T_4_36	1	T_4_36
	15	T_5_7	1	T_5_7
Motril				M_3_2
	1	M_3_2	3	M_3_1
				M_1_1
	2	M_2_3	1	M_2_3
	3	M_3_16	1	M_3_16
Burguillos				B_1_115
				B_1_121
				B_1_23
				B_1_38
				B_1_40
	1	B_1_115	11	B_1_44
				B_1_64
				B_1_71
				B_1_94
				B_1_98
				B_1_89
				B_1_113
				B_1_114
	2	B_1_113	6	B_1_43
				B_1_45
				B_1_59
				B_1_72
	3	B_1_6	2	B_1_6
				B_1_28
	4	B_1_10	1	B_1_10
	5	B_1_42	1	B_1_42
	6	B_1_62	1	B_1_62

Supplemental Figure S1. Multiple alignment of *asmA* marker gene (465 pb) in *Ensifer* sp.

<i>Ensifer meliloti</i> 1021 (consensus)	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer meliloti</i> 2011	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer meliloti</i> RM017	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer meliloti</i> BL225C	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer meliloti</i> SM11	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer meliloti</i> Rm41	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer meliloti</i> GR4	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer meliloti</i> AK83	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer medicae</i> WSM419	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer fredii</i> NGR 234	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer meliloti</i> 1021	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer americanum</i> CFNBI156	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer fredii</i> USDA191	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer xinjiangense</i> LMG17930	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
<i>Ensifer arboris</i> LMG 14919	GCCGGCGCAGGGGGACGACCGCGGATGGCGGATGGCCAGGCTCTTGTGCGGGGGTGGCG
T1_27	*
T1_45	*
T1_71	*
T1_81	*
T1_100	*
T2_8	*
M4_29	*
M4_46	*
<i>Ensifer meliloti</i> 1021 (consensus)	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer meliloti</i> 2011	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer meliloti</i> RM017	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer meliloti</i> BL225C	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer meliloti</i> SM11	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer meliloti</i> Rm41	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer meliloti</i> GR4	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer meliloti</i> AK83	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer medicae</i> WSM419	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer fredii</i> NGR 234	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer meliloti</i> 1021	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer americanum</i> CFNBI156	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer fredii</i> USDA191	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer xinjiangense</i> LMG17930	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
<i>Ensifer arboris</i> LMG 14919	CCTGGATGTCGGGTGTCGTCGAGGAAGTCTGGCCGGTCCGTTGCACCTGACCGATGTC
T1_27	**
T1_45	**
T1_71	**
T1_81	**
T1_100	**
T2_8	**
M4_29	**
M4_46	**
<i>Ensifer meliloti</i> 1021 (consensus)	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer meliloti</i> 2011	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer meliloti</i> RM017	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer meliloti</i> BL225C	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer meliloti</i> SM11	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer meliloti</i> Rm41	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer meliloti</i> GR4	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer meliloti</i> AK83	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer medicae</i> WSM419	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer fredii</i> NGR 234	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer meliloti</i> 1021	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer americanum</i> CFNBI156	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer fredii</i> USDA191	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer xinjiangense</i> LMG17930	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer arboris</i> LMG 14919	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
T1_27	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
T1_45	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
T1_71	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
T1_81	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
T1_100	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
T2_8	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
M4_29	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
M4_46	GCGGCGGGGTGATGATCGATGGAAGCCGGGTTTCGCTCGACATCGGAGACAGCACCTAT
<i>Ensifer meliloti</i> 1021 (consensus)	GCGAATGGCAGGCTGAGCGGCCGCGCGTCTCGGAGAAAGGGTGGGGCAAGGCGGC
<i>Ensifer meliloti</i> 2011	GCGAATGGCAGGCTGAGCGGCCGCGCGTCTCGGAGAAAGGGTGGGGCAAGGCGGC
<i>Ensifer meliloti</i> RM017	GCGAATGGCAGGCTGAGCGGCCGCGCGTCTCGGAGAAAGGGTGGGGCAAGGCGGC
<i>Ensifer meliloti</i> BL225C	GCGAATGGCAGGCTGAGCGGCCGCGCGTCTCGGAGAAAGGGTGGGGCAAGGCGGC
<i>Ensifer meliloti</i> SM11	GCGAATGGCAGGCTGAGCGGCCGCGCGTCTCGGAGAAAGGGTGGGGCAAGGCGGC
<i>Ensifer meliloti</i> Rm41	GCGAATGGCAGGCTGAGCGGCCGCGCGTCTCGGAGAAAGGGTGGGGCAAGGCGGC
<i>Ensifer meliloti</i> GR4	GCGAATGGCAGGCTGAGCGGCCGCGCGTCTCGGAGAAAGGGTGGGGCAAGGCGGC

197

Supplemental Tables and Figures

T1_71	CTCTGGGAGGCCGGCGTCGCCGATGTCTTCGGCCGTGTGATTATTCGCTCACCAATGGCA
T1_81	CTCTGGGAGGCCGGCGTCGCCGATGTCTTCGGCCGTCTGATTATTCGCTCACCAATGGCA
T1_100	CTCTGGGAGGCCGGCGTCGCCGATGTCTTCGGCCGTCTGATTATTCGCTCACCAATGGCA
T2_8	CTCTGGGAGGCCGGCGTCGCCGATGTCTTCGGCCGTCTGATTATTCGCTCACCAATGGCA
M4_29	CTCTGGGAGACGGGGTCGCCGATGTCTTCGGCCGTCTGATTATTCGCTCACCAACGGCA
M4_46	CTCTGGGAGACGGGGATCGCCGATGTCTTCGGCCGTCTGATTATTCGCTCACCAACGGCA
	* * * * *
<i>Ensifer meliloti</i> 1021 (consensus)	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
<i>Ensifer meliloti</i> 2011	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
<i>Ensifer meliloti</i> RM017	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
<i>Ensifer meliloti</i> BL225C	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
<i>Ensifer meliloti</i> SM11	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
<i>Ensifer meliloti</i> Rm41	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
<i>Ensifer meliloti</i> GR4	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
<i>Ensifer meliloti</i> AK83	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
<i>Ensifer medicae</i> WSM419	GCCTCGCCGGTTTCGACGTTTCGCGCCTTCACCGA
<i>Ensifer fredii</i> NGR 234	GCATCGCCGGCTTCGATGCCCCGCTTTGCCGA
<i>Ensifer meliloti</i> 1021	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
<i>Ensifer americanum</i> CFNBI156	GCATCGCCGGCTTCGATGCCCCACGCCTTCGCCGA
<i>Ensifer fredii</i> USDA191	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
<i>Ensifer xinjiangense</i> LMG17930	GCATCGCCGGCTTCGACGCCCCACGCCTTTGCCGA
<i>Ensifer arboris</i> LMG 14919	ATCTCGTCGATATGGACATTCACGCCTTCACCGA
T1_27	GCCTCGCCGGTTTCGACGTTTCGCGCCTTCACCGA
T1_45	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
T1_71	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
T1_81	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
T1_100	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
T2_8	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
M4_29	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
M4_46	GGCTCGCCGATTTTCGACGTTACGCCTTCACCGA
	*** ** * ** * ***** ****

Supplemental Tables and Figures

<i>B.elkanii</i> WSM2783	GAGATCGAGTTCGAGGGCGCGACCGTCTACCGCGTCAACGGCTGATCGATGATTGCGGC
<i>B.elkanii</i> USDA 3259	GAGATCGAGTTCGAGGGCGCCACCGTCTATCGTGTCAACGGCTATATCGACGACTGCGGC
<i>B.japonicum</i> USDA 122	GAGATCGAGTTCGAGGGTGCAGCCGTCTACCGCGTCAACGGCTATATCGACGATTGCGGC
<i>B.sp.</i> WSM3983	GAGATCGAGTTCGAGGGCGCGACCGTCTACCGCGTCAACGGCTATATCGACGATTGCGGA
<i>B.sp.</i> WSM1743	GAGATCGAGTTCGAGGGCGCCACCGTCTACCGCGTCAACGGCTATATCGATGATTGCGGC
<i>B.sp.</i> YR681	GAGATCGAGTTCGAGGGCGCAACCGTCTACCGCGTCAACGGCTATATCGACGATTGCGGC
<i>B.sp.</i> ARR65	GAGATCGAGTTCGAGGGCGCGACCGTCTACCGCGTCAACGGCTGATCGATGATTGCGGC
<i>B.sp.</i> WSM1253	GAGATCGAGTTCAGGGCGCGACCGTCTATCGCGTCAACGGCTATATCGACGATTGCGGC
<i>B.sp.</i> WSM471	GAGATCGAGTTCGAGGGCGCCACCGTCTACCGCGTCAACGGTATATCGATGATTGCGGC
<i>B.sp.</i> WSM1417	GAGATCGAGTTCAGGGCGCGACCGTGTACCGCGTCAACGGCTATATCGACGATTGCGGC
<i>B.PZZA</i>	GAGATCGAGTTCGAGGGCGCGACCGTCTACCGCGTCAACGGCTATATCGACGATTGCGGC
<i>B.USDA</i> 110	GAGATCGAGTTCGAGGGCGCGACCGTCTACCGCGTCAACGGCTATATCGACGATTGCGGC
<i>B.betae</i> LMG21987	GAGATCGAGTTCGAGGGCGCCACCGTCTATCGCGTCAACGGCTATATCGACGATTGCGGC
<i>B.yuanmingense</i> LMG21827	GAGATCGAGTTCGAGGGCGCGACCGTCTACCGCGTCAACGGCTATATCGACGATTGCGGC
<i>B.japonicum</i> USDA 6	GAGATCGAGTTCAGGGCGCGACCGTCTATCGCGTCAACGGCTATATCGACGATTGCGGC
<i>B.Liaonigense</i> LMG18230	GAGATCGAGTTCGAGGGCGCGACCGTCTACCGCGTCAACGGCTATATCGATGATTGCGGC
<i>B.elkanii</i> USDA 76	GAGATCGAGTTCGAGGGCGCCACCGTGTATCGCGTCAACGGCTATATCGATGATTGCGGC
<i>B.Lmjc</i> m3	GAGATCGAGTTCGAGGGCGCCACCGTCTATCGCGTCAACGGTCTGATCGACGATTGCGGC
B3_2	GAAATCGAGTTCGAGGGCGCGACCGTCTATCGCGTCAACGGCTATATCGACGATTGCGGC
B3_1	GAGATCGAGTTCGAGGGCGCGACCGTCTACCGCGTCAACGGCTATATCGACGATTGCGGC
B2_9	GAGATCGAGTTCGAGGGCGCCACCGTCTACCGCGTCAACGGCTATATCGACGATTGCGGC
B2_4	GAGATCGAGTTCGAGGGCGCGACCGTCTACCGCGTCAATGGCTATATCGATGATTGCGGC
B2_2	GAGATCGAGTTCGAGGGCGCGACCGTCTATCGCGTCAACGGCTGATCGACGATTGCGGC
B10_118	GAGATCGAGTTCGAGGGCGCGACCGTCTACCGGGTCAACGGTTACATCGACGATTGCGGC
	** ***** * ***** ** ** *
<i>B.japonicum</i> USDA 6 (con.)	AAGATCGTCGCGCAAGGCAAGGCAAAAGTCGGCTGGTTCGACACGTCACGCTCAAGGAG
<i>B.japonicum</i> USDA 6	AAGATCGTCGCGCAAGGCAAGGCAAAAGTCGGCTGGTTCGACACGTCACGCTCAAGGAG
<i>B.japonicum</i> 22	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACGCTGAAGGAG
<i>B.sp.</i> CCGE-LA001	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACGCTGAAGGAG
<i>B.sp.</i> ORS 375	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.elkanii</i> USDA 76	AAGATCGTCGCGGAGGGAAAAGCGAAAGTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.japonicum</i> USDA 124	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACGCTGAAGGAG
<i>B.genosp.</i> SA-4 CB756	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACTTCGACGCTGAAGGAG
<i>B.sp.</i> WSM4349	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.sp.</i> STM 3809	AAGATCGTCGCGCAAGGCAAGGCCAAGGTTCGGCTGGTTGATACCTCGACGCTGAAGGAG
<i>B.sp.</i> ORS285	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.sp.</i> S23321	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACGTCACGCTGAAGGAG
<i>B.sp.</i> WSM2793	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.sp.</i> WSM2254	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACGTCACGCTGAAGGAG
<i>B.japonicum</i> USDA 38	AAAATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACGCTGAAGGAG
<i>B.elkanii</i> USDA 3254	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAAGAG
<i>B.sp.</i> TV2a.2	AAGATCGTCGCGGAGGGTAAAGGCCAAGGTTCGGCTGGTTCGATACCTCGACCTGAAGGAA
<i>B.japonicum</i> USDA 4	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACGCTAAAGGAG
<i>B.elkanii</i> USDA 94	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGATACCTCGACCTGAAGGAG
<i>B.sp.</i> USDA 3384	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACTTCGACGCTGAAGGAG
<i>B.sp.</i> EC3.3	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACGCTGAAGGAG
<i>B.japonicum</i> USDA 135	AAGATCGTCGCGGAGGGAAAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.sp.</i> WSM1741	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.elkanii</i> WSM2783	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACGTCGACCTGAAGGAA
<i>B.elkanii</i> USDA 3259	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAAGAG
<i>B.japonicum</i> USDA 122	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCCACGCTGAAGGAG
<i>B.sp.</i> WSM3983	AAGATCGTCGCGGAGGAAAAGCAAAAGTCGGCTGGTTCGACACCTCGACACTGAAGGAG
<i>B.sp.</i> WSM1743	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.sp.</i> YR681	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACGCTGAAGGAG
<i>B.sp.</i> ARR65	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGATACCTCCACCTGAAGGAG
<i>B.sp.</i> WSM1253	AAGATCGTCGCGGAGGGCAAGGCAAAAGTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.sp.</i> WSM471	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.sp.</i> WSM1417	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.PZZA</i>	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.USDA</i> 110	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCCACCTGAAGGAG
<i>B.betae</i> LMG21987	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGATACCTACGCTGAAGGAG
<i>B.yuanmingense</i> LMG21827	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.japonicum</i> USDA 6	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACGTCACGCTCAAGGAG
<i>B.Liaonigense</i> LMG18230	AAGATCGTCGCGGAGGGAAAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.elkanii</i> USDA 76	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACCTGAAGGAG
<i>B.Lmjc</i> m3	AAGATCGTCGCGGAGGGAAAAGGCCAAGGTTCGGCTGGTTCGATACCTCGACGCTGAAGGAG
B3_2	AAGATCGTCGCGGAGGGAAAAGGCCAAGGTTCGGCTGGTTCGATACCTCGACGCTGAAGGAG
B3_1	AAGATCGTTGGCGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACGCTGAAGGAG
B2_9	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCCACGCTGAAGGAG
B2_4	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACGCTGAAGGAG
B2_2	AAGATCGTCGCGGAGGGCAAGGCCAAGGTTCGGCTGGTTCGACACCTCGACGCTGAAGGAG
B10_118	AAGATCGTCGCGGAGGGAAAAGGCCAAGGTTCGGCTGGTTCGATACCTCGACCTGAAGGAG
	** ***** ** *

<i>B. japonicum</i> USDA 6 (con.)	CCGTACCGTATCGAGGGCAAGAAGACGATGGG
<i>B. japonicum</i> USDA 6	CCGTACCGTATCGAGGGCAAGAAGACGATGGG
<i>B. japonicum</i> 22	CCGTACCGCATCGAAGGCAAGAAGACGATGGG
<i>B.sp.</i> CCGE-LA001	CCGTACCGCATCGAGGGCAAGAAGACGATGGG
<i>B.sp.</i> ORS 375	CCGTACCGGATCGAGGGCAAGAAGACGATGGG
<i>B.elkanii</i> USDA 76	CCGTACCGGATCGAGGGCAAGAAGACGATGGG
<i>B. japonicum</i> USDA 124	CCGTACCGCATCGAAGGCAAGAAGACGATGGG
<i>B.genosp.</i> SA-4 CB756	CCGTACCGTATCGAAGGCAAGAAGACGATGGG
<i>B.sp.</i> WSM4349	CCGTACCGCATCGAAGGCAAGAAGACGATGGG
<i>B.sp.</i> STM 3809	CCTTACCGCATCGAGGGCAAGAAGACGATGGG
<i>B.sp.</i> ORS285	CCGTACCGCATCGAAGGCAAGAAGACGATGGG
<i>B.sp.</i> S23321	CCGTACCGTATCGAAGGCAAGAAGACGATGGG
<i>B.sp.</i> WSM2793	CCGTACCGCATCGAGGGCAAGAAGACGATGGG
<i>B.sp.</i> WSM2254	CCGTACCGCATCGAAGGCAAGAAGACGATGGG
<i>B. japonicum</i> USDA 38	CCGTACCGTATCGAGGGCAAGAAGACGATGGG
<i>B.elkanii</i> USDA 3254	CCGTACCGGATCGAGGGCAAGAAGACGATGGG
<i>B.sp.</i> TV2a.2	CCCTATCGGATCGAGGGCAAGAAGACGATGGG
<i>B. japonicum</i> USDA 4	CCGTACCGCATCGAAGGCAAGAAGACGATGGG
<i>B.elkanii</i> USDA 94	CCGTACCGGATCGAGGGCAAGAAGACGATGGG
<i>B.sp.</i> USDA 3384	CCGTACCGTATCGAAGGCAAGAAGACGATGGG
<i>B.sp.</i> EC3.3	CCGTACCGGATCGAGGGCAAGAAGACGATGGG
<i>B. japonicum</i> USDA 135	CCGTACCGCATCGAGGGCAAGAAGACGATGGG
<i>B.sp.</i> WSM1741	CCTTACCGGATCGAAGGCAAGAAGACGATGGG
<i>B.elkanii</i> WSM2783	CCGTACCGGATCGAGGGCAAGAAGACGATGGG
<i>B.elkanii</i> USDA 3259	CCGTACCGGATCGAGGGCAAGAAGACGATGGG
<i>B. japonicum</i> USDA 122	CCGTACCGCATCGAAGGCAAGAAGACGATGGG
<i>B.sp.</i> WSM3983	CCGTACCGCATCGAAGGCAAGAAGACGATGGG
<i>B.sp.</i> WSM1743	CCGTACCGCATCGAGGGCAAGAAGACGATGGG
<i>B.sp.</i> YR681	CCGTACCGAATCGAAGGCAAGAAGACGATGGG
<i>B.sp.</i> ARR65	CCATATCGGATCGAAGGCAAGAAGACGATGGG
<i>B.sp.</i> WSM1253	CCGTACCGGATCGAAGGCAAGAAGACGATGGG
<i>B.sp.</i> WSM471	CCGTACCGGATCGAAGGCAAGAAGACGATGGG
<i>B.sp.</i> WSM1417	CCGTACCGGATCGAAGGCAAGAAGACGATGGG
<i>B.betae</i> LMG21987	CCGTACCGCATCGAGGGCAAGAAGACGATGGG
<i>B.Lmj</i> c m3	CCTTACCGGATCGAAGGCAAGAAGACGATGGG
<i>B.elkanii</i> USDA 76	CGTACCGGATCGAGGGCAAGAAGACGATGGG
<i>B.Liaonigense</i> LMG18230	CCGTACCGCATCGAGGGCAAGAAGACGATGGG
<i>B.yuanmingense</i> LMG21827	CCGTACCGCATCGAGGGCAAGAAGACGATGGG
<i>B.UsDA</i> 110	CCGTACCGTATCGAAGGCAAGAAGACGATGGG
<i>B.PZZA</i>	CCGTACCGCATCGAGGGCAAGAAGACGATGGG
<i>B. japonicum</i> USDA 6	CCGTACCGTATCGAGGGCAAGAAGACGATGGG
B3_2	CCGTACCGCATCGAGGGCAAGAAGACGATGGG
B3_1	CCGTACCGCATCGAAGGCAAGAAGACGATGGG
B2_9	CCGTACCGCATCGAAGGCAAGAAGACGATGGG
B2_4	CCGTACCGCATCGAAGGCAAGAAGACGATGGG
B2_2	CCGTACCGGATCGAGGGCAAGAAGACGATGGG
B10_118	CCGTATCGGATCGAGGGCAAGAAGACGATGGG
	** ** ** ** ** *****