UNIVERSIDAD POLITÉCNICA DE MADRID

ESCUELA TÉCNICA SUPERIOR DE INGENIEROS DE
TELECOMUNICACIÓN

Deep Learning for assessing pulmonary
parenchyma disease in smoking-related injury

Ph.D. Thesis

DAVID BERMEJO PELÁEZ

2020

Biomedical Image Technologies Laboratory
Department of Electronic Engineering
Escuela Técnica Superior de Ingenieros de Telecomunicación
Universidad Politécnica de Madrid

PhD Thesis

Deep Learning for assessing pulmonary parenchyma disease in smoking-related injury

Author

David Bermejo Peláez

Advisors

María Jesús Ledesma Carbayo – Universidad Politécnica de Madrid, Madrid, Spain

Raúl San José Estépar – Harvard Medical School & Brigham and Women's Hospital, Boston,
USA

Copyright © 2020 David Bermejo Peláez

Agradecimientos

Porque no sé si hubiera acabado esta tesis sin todos ellos. Porque si lo hubiera hecho, estos años no hubieran sido los mismos. Gracias.

A mis directores de tesis María Jesús y Raúl, por su continuo soporte, ánimo y guía durante estos años. Por confiar tantas veces en mí más de lo que yo confiaba en mí mismo. Por sus ideas, por sus ganas de abrir horizontes.

Al que considero *mi* grupo, el grupo de Tecnologías de Imágenes Biomédicas (BIT). A todos los que lo forman, a Andrés, Chus, Juan, Juanjo, Eva, Lara, Felicia, Benito... A los itinerantes que han pasado también por allí durante estos años. Por hacer un grupo excepcional, tanto en el ámbito profesional como en el personal. Por vuestro apoyo, calidez, cercanía y en muchos casos amistad.

Al Applied Chest Imaging Laboratory (ACIL). A Raúl, de nuevo, George, y Sam porque sin vosotros esta tesis no hubiera sido ni parecida. Al resto del equipo, Gonzalo, Pietro, Jorge y Rubén. Por haberme acogido durante todas mis estancias como uno más desde el primer día. Por haberme hecho sentir que Boston es mi segunda ciudad.

A mi familia, por su continuo apoyo y paciencia desde siempre, especialmente en estos años. Por escucharme. Por aconsejarme en mis indecisiones. A mi tío Miguel, por su especial soporte, guía e interés en lo académico y profesional.

A Dios. Por todo.

David Bermejo Peláez

Junio 2020

Abstract

Tobacco smoking is one of the health risk factors most associated with population morbidity and mortality causing a variety of lung diseases that are responsible for the death of more than 8 million people each year. The inhalation of cigarette smoke can affect the lung in diverse ways and is directly associated with the development of lung parenchyma disorders including emphysema, characterized by airspace dilation and alveolar destruction, and interstitial or fibrotic changes caused by the inflammation and scarring of pulmonary parenchyma. These parenchymal injuries are often an irreversible process specially when they are in an advanced stage.

High resolution computed tomography (HRCT) plays an essential role in the diagnosis and assessment of disease prognosis and progression of lung diseases. In this thesis, we developed deep learning approaches to characterize parenchymal injury patterns on HRCT, which may facilitate disease understanding, serve as a strong basis to assess disease prognosis and progression, and facilitate automated diagnosis of lung diseases at early stages.

In this Ph.D. thesis, we described a novel optimally cost-effective convolutional neural network (CNN) to characterize emphysema subtypes which was subsequently extended by developing an ensemble of multiple CNN covering the lesions' multiscale representation designed to identify interstitial lung abnormalities that are found as early sign of fibrotic disorders in the lung parenchyma. Moreover, we proposed an innovative CNN segmentation architecture to identify paraseptal emphysema lesions, a particular subtype of emphysema with a unique notion of location, that leverages and exploits 3D contextual information while producing 2D annotations alleviating the need of having 3D labels which carries a high burden from experts. Finally, we designed a novel methodology that learns the spatial interdependence of parenchymal injuries patterns using a segmentation approach from an initial labelling based on local methods. This new paradigm relieves the need for delimited annotations to perform segmentation tasks.

Developed approaches have demonstrated that are able to identify subtype injury patterns which are related to different clinical outcomes. Therefore, these methods allow to advance in the understanding of both lung diseases and the clinical implications of the different parenchymal patterns.

Resumen

El consumo de tabaco es uno de los mayores factores de riesgo para la salud asociado a una alta morbilidad y mortalidad debido a que causan una variedad de enfermedades pulmonares responsables de la muerte de más de 8 millones de personas al año. El tabaco afecta a los pulmones de diversas maneras y está directamente relacionado con el desarrollo de desórdenes en el parénquima pulmonar como el enfisema pulmonar, caracterizado por la dilatación de los alveolos o la destrucción de sus paredes, así como cambios fibróticos causados por una inflamación y cicatrización del tejido pulmonar. Estos daños del tejido son usualmente procesos irreversibles, especialmente cuando ocurren en una etapa avanzada.

La Tomografía Axial Computarizada (TAC) juega un papel esencial en el diagnóstico y la evaluación de la progresión de las enfermedades pulmonares. En esta Tesis Doctoral se desarrollan diferentes enfoques basados en Deep Learning para la caracterización automática de daños en el parénquima pulmonar mediante imágenes de TAC que puedan facilitar y mejorar el entendimiento de las enfermedades, puedan servir como base para evaluar la progresión de la misma, y puedan facilitar el diagnóstico automático de las enfermedades pulmonares en estadios tempranos.

En esta Tesis Doctoral se propone una nueva y eficiente arquitectura de red neuronal convolucional (CNN, del inglés *Convolutional Neural Network*) especialmente diseñada para la caracterización de distintos subtipos de enfisema pulmonar. Además, esta nueva técnica propuesta ha sido extendida desarrollando un método que combina múltiples arquitecturas CNN que cubren distintas representaciones locales de las lesiones pulmonares para la identificación de anomalías intersticiales, consideradas como un signo temprano y que precede a la aparición de cambios fibróticos irreversibles en el parénquima pulmonar. Además, se propone una innovadora arquitectura de CNN para la segmentación de enfisema paraseptal, una lesión particular que tiene características únicas de localización. Esta arquitectura explota y aprovecha el contexto tridimensional alrededor de la lesión mientras que produce segmentaciones bidimensionales, de forma que elimina el hecho de necesitar anotaciones tridimensionales disminuyendo así la carga de expertos requerida para construir dichas etiquetas. Finalmente, se ha diseñado una nueva metodología para inferir y aprender las diferentes dependencias espaciales entre los distintos subtipos de parénquima pulmonar. Para ello se propone una técnica de segmentación a partir de un etiquetado inicial basado en métodos locales. De esta forma, además este nuevo paradigma elimina la necesidad de tener bases de datos de imágenes anotadas para llevar a cabo tareas de segmentación.

Los métodos desarrollados en esta Tesis Doctoral han demostrado que son capaces de identificar subtipos de parénquima pulmonar que están relacionados con diferentes resultados clínicos. Por tanto, estos métodos permiten el avance en una mejor comprensión de las enfermedades pulmonares así como de las implicaciones clínicas de los distintos patrones radiológicos presentes en el parénquima pulmonar.

Contents

1	Motivation and objectives	9
1.1	Motivation	10
1.2	Objectives	10
2	Clinical context and state of the art	13
2.1	Clinical context	14
2.1.1	Smoking-related lung injuries	14
2.1.2	Pathophysiology of parenchymal injuries	16
2.1.3	Quantitative imaging	22
2.2	State of the art in parenchymal injuries quantification	23
2.2.1	Lung density analysis	23
2.2.2	Texture and Machine learning-based analysis	25
3	Emphysema classification	29
3.1	Introduction	30
3.2	Methods	30
3.2.1	Multi-view Convolutional Network (2.5D CNN)	31
3.2.2	Comparative methods	32
3.3	Experiments and results	33
3.3.1	Dataset and evaluation setup	33
3.3.2	Validation	33
3.4	Conclusions	36
4	Parenchymal injuries classification	39
4.1	Introduction	40
4.1.1	Related work	40
4.2	Methods	42
4.2.1	Database	42
4.2.2	Multi-model ensemble	43
4.2.3	Overfitting prevention	45
4.3	Experiments and results	47
4.3.1	Experimental setup	47

4.3.2	Validation results	49
4.4	Conclusions	61
5	Paraseptal emphysema segmentation	65
5.1	Introduction	66
5.2	Methods	67
5.2.1	SR-Net	67
5.2.2	Comparative methods	69
5.3	Experiments and results	69
5.3.1	Dataset and evaluation approach	69
5.3.2	Results	70
5.4	Conclusions	72
6	Parenchymal injuries segmentation	73
6.1	Introduction	74
6.2	Methods	75
6.2.1	Domain transformation	75
6.2.2	SR-Net with domain transformation	76
6.3	Experiments and results	78
6.3.1	Training dataset	78
6.3.2	Visual validation	79
6.3.3	Clinical associations	81
6.4	Conclusions	82
7	Concluding remarks	85
	Publications	89
	Bibliography	90

1

Motivation and objectives

1.1 Motivation

Smoking is one of the health risk factors most associated with population morbidity and mortality. Tobacco is the only legal drug which, used in moderation and exactly as intended by manufacturers, causes severe harm to the consumers and even kills many of its users. The World Health Organization (WHO) has estimated that tobacco use is currently responsible for the death of more than 8 million people across the world each year. More than 7 million of those deaths were attributable to direct cigarette smoking, and around 1.2 million are the result of secondhand smoke.

Smoking related lung diseases are the respiratory manifestations of disease that are related to cigarette smoking. Inhalation of cigarette smoke can affect the lung in diverse ways, and is directly associated with the development of lung parenchyma disorders (Auerbach et al. 1974; Ryu et al. 2001).

Damage to the lungs is often an irreversible process, specially when it is in an advanced stage. Furthermore, many patients affected by lung diseases remain underdiagnosed. In this context, a precise and early diagnosis is highly important and remains critical to properly caring of these patients. This has become increasingly important specially for those patients with pulmonary fibrosis, a progressive and frequently fatal disease, but where new therapies that slow the disease progression have been introduced (Talmadge et al. 2014).

High resolution computed tomography (HRCT) plays an essential role in the diagnosis and assessment of disease prognosis and progression of lung diseases (Mai et al. 2017). Findings in HRCT images have a substantial effect on subsequent management decisions. However, visual subjective analysis for disease identification is time consuming, often insensitive to subtle changes and requires a high degree of expertise from radiologists. Objective identification of lung diseases based on HRCT images would be of enormous benefit, facilitating automated diagnostic support at early stages, disease understanding, and assessment of disease prognosis and progression.

New techniques in artificial intelligence, specifically the application of machine learning and deep learning to interpreting complex medical image data, have revolutionized the field by helping with objective diagnosis, detecting preliminary signs of disease, and predicting how the disease will develop.

1.2 Objectives

The main objective of this PhD Thesis is the **design, development and validation of completely automatic and objective methods to characterize and quantify pulmonary parenchymal injuries**. The most prominent hypothesis under the present work is that such methods

would support automated diagnostic and quantification of smoke-related lung diseases, specially at early stages. Additionally, the methods could facilitate disease understanding and become a potential source of information for disease prognosis.

This general objective will be addressed through the fulfillment of the following sub-objectives:

1. Design and implementation of local-based methods for the objective and automated quantification of lung parenchymal injuries. Lung parenchymal injuries have been traditionally defined as a semi-local damage affecting the pulmonary secondary lobule which is considered the fundamental unit of lung structure and function. This sub-objective aims at developing methods to identify and classify in an automatic manner different parenchymal injuries exploiting the local appearance of HRCT covering the secondary lobule extension.
2. Design and implementation of global-based methods for the objective and automated quantification of lung parenchymal injuries. In order to improve the identification of parenchymal lung injuries and to incorporate context and spatial information, we developed segmentation methods which uses global context. Getting precise annotations for training global-based methods carries a high burden from experts. We also proposed strategies to reduce the need of collecting such demanding reference data.
3. Clinically validate the proposed methods. To achieve clinical impact we exhaustive validated the proposed methods to test the association between quantitative measurements of parenchyma lung injuries and clinical and functional variables.

2

Clinical context and state of the art

2.1 Clinical context

2.1.1 Smoking-related lung injuries

Tobacco smoke is a toxic mixture containing around 5000 chemicals and considered as one of the most important causes of chemically mediated disorders in humans (Borgerding et al. 2005; Fowles et al. 2003; Thielen et al. 2008). Both direct toxicity and induced immune-mediated response result in a wide range of parenchymal abnormalities in the lung. These parenchymal injuries include both airspace dilation and alveolar destruction leading to a local decreases in lung density typically called emphysema, and inflammation and scarring leading to local increases in lung density, typically called as interstitial changes or fibrotic disorders (Houghton et al. 2008; Kumar et al. 2018; J. H. Ryu et al. 2001).

Many pathological and radiological findings coexist in the lung of a smoker. There is a increasing clinical, radiological and pathological recognition of the coexistence of emphysema and interstitial fibrosis, which appears to be at the opposite end of the spectrum of smoking-related lung diseases (Madan et al. 2016) (Figure 2.2). Combined pulmonary fibrosis and emphysema (CPFE) is a clinical entity characterize by the coexistence of both lung diseases (Cottin, Nunes, Brillet, et al. 2005). When both disorders appears in the same patient, emphysema usually appears in the upper lobe, while pulmonary fibrosis is located in the lower lobe (Figure 2.1). The coexistence of the two conditions is considered to have different pathophysiological characteristics. This overlapping spectrum warrants inclusion as a novel, distinct pulmonary manifestation within the spectrum of smoking-related disorders which have a distinct characteristics and prognosis compared to its individuals components (Cottin, Nunes, Mouthon, et al. 2011). The overall mortality of CPFE patients in comparison with those patients with lone fibrosis is still matter of debate (Bodlet et al. 2013; Kurashima et al. 2010; Mejia et al. 2009; Sugino et al. 2014). But CPFE has been related to pulmonary hypertension (Cottin, Nunes, Brillet, et al. 2005) and lung cancer (Kwak et al. 2014), which are associated with reduced survival in CPFE patients (Kumagai et al. 2014; Mejia et al. 2009; Sugino et al. 2014).

Computed tomography (CT) plays a critical diagnostic role, as it actually constitutes the main examination tool for establishing in vivo the diagnosis and evaluation of emphysema and interstitial lung diseases. CT also plays an important role in predicting the prognosis of the diseases. Quantitative CT has been established as the main tool for measuring the extent of those parenchymal disorders, also building a better understanding of the relationship between morphologic imaging findings and pulmonary function (Matsuoka et al. 2015b). Additionally, research use of CT has risen awareness that smokers can manifest early signs of pulmonary emphysema or interstitial lung abnormalities (ILA) many years before symptoms

Figure 2.1: **A-B** High resolution computed tomography (HRCT) scan of a patient with CPFE with different distribution of the zones of emphysema (upper zones) and fibrosis (basal zones). **C** Gross photo showing marked upper lobe emphysema with fibrosis in lower lobes. Sources (Tzilas et al. 2016) under CC BY 4.0 and (Inomata et al. 2014) under CC BY 2.0.

Figure 2.2: Spectrum of smoking-related lung diseases. Combined pulmonary fibrosis and emphysema disorder (CPEF) covers the overlapping area between emphysema and fibrosis.

appears. This findings present the possibility of studying smoking-related lung destruction prior to establishment of end-stage injury.

2.1.2 Pathophysiology of parenchymal injuries

Lungs are the essential organs in the respiratory system. Each lung is divided into lobes by fissures. The left lung is divided into two lobes (superior and inferior) and the right lung is divided into three lobes (superior, middle, and inferior) (see Figure 2.3). The lobes can be further divided into functional units called bronchopulmonary segments.

The trachea divides into two primary bronchi, which enter on its respective lung and divides into smaller branches called secondary bronchi. The secondary bronchi continue to branch forming tertiary bronchis, bronchioles, terminal bronchioles and respiratory bronchioles which have thin, gas-exchanged walls. Respiratory bronchioles transition into alveolar ducts, which open into numerous alveoli which forms alveolar sacs. The acinus is defined as the unit of the lung structure distal to the terminal bronchiole and that consists on alveolar ducts, alveolar sacs and alveoli (Berend et al. 1991).

The lung is composed of many anatomic structures smaller than a pulmonary lobe or segment. The secondary pulmonary lobule is considered the fundamental unit of lung structure and function. Secondary pulmonary lobules are irregularly polyhedral in shape and are approximately 1 to 2.5 cm on a side. Secondary lobules are supplied by a bronchiole and pulmonary artery branch and are delineated by connective tissue septa (Heitzman et al. 1969a; Heitzman et al. 1969b). Airways, pulmonary arteries and veins, as well as numerous components of the pulmonary interstitium are all normally observed at the level of the pulmonary secondary lobule. Several lung diseases, which have been shown to affect different components of the secondary lobule, can be studied by comparing the appearance of the normal secondary lobule with abnormal secondary lobules (Kawata et al. 2010; Webb 2006).

2.1.2.1 COPD: emphysema

Chronic obstructive pulmonary disease (COPD) is a general name for the chronic airflow limitation which is normally related to inflammatory processes in airways. COPD is developed most often as a result of cigarette smoking, although other factors, such as prolonged exposure to pollution, certain chemicals or genetic disorders (α_1 -antitrypsin deficiency) may affect to the development of the disease. The pathology of COPD cover numerous lesions in the airways, lung parenchyma and pulmonary vasculature, and these lesions can be correlated with changes in pulmonary function.

COPD encompass lesions of the lung parenchyma called emphysema. Emphysema is defined as the "abnormal permanent enlargement of the airspaces distal to the terminal

Figure 2.3: A Lung anatomy. B Pulmonary secondary lobule.

bronchioles accompanied by destruction of the alveolar wall without obvious fibrosis" (Snider et al. 1985). Enlargement of airspaces leads to hyperinflation of the lungs producing an increase in total lung capacity (TLC). As shown in Figure 2.4, in contrast to the normal lung, the emphysematous lung parenchyma shows a marked destruction of alveolar walls, leading to confluent alveoli and thus hyperinflation.

Figure 2.4: Smoking associated emphysema. **A** An area of emphysematous lung with severe effacement of alveolar walls. **B** Normal lung with intact alveolar walls. Source (Higham et al. 2019) under CC BY 4.0.

Emphysema can be morphologically classified according to the location of the lesions within the acinus. There are three commonly recognized types of emphysema: centrilobular, panlobular and paraseptal emphysema (Figures 2.5). CT is a reliable tool for demonstrating the pathology of emphysema, even in subtle changes within the pulmonary secondary lobules (Figure 2.6).

Figure 2.5: Morphologic classification of emphysema. Centrilobular emphysema: destruction and enlargement of respiratory bronchiole. Panlobular emphysema: uniform destruction of airspaces beyond the respiratory bronchiole. Paraseptal emphysema: destruction of subpleural areas.

Centrilobular emphysema The destruction affects the bronchioles in the central part of the secondary lobule, while the peripheral alveolar ducts and alveoli may be preserved. Centrilobular emphysema is most common in the upper lobes, and is most frequently

associated with smoking. CT in centrilobular emphysema shows evenly distributed centrilobular areas of low attenuation surrounded by normal tissue. As centrilobular emphysema becomes more severe, low attenuation areas become confluent.

Panlobular emphysema Shows destruction of the whole secondary lobule, with destruction of alveolar septa from the center to the periphery of the acinus. This type of emphysema is most common and severe in the lower lobes, and is most frequently associated with α_1 -AT deficiency. CT shows large low attenuation areas in panlobular emphysema. Early stages of this subtype are difficult to identify at CT.

Paraseptal emphysema Also known as distal acinar emphysema, is characterized by destruction of subpleural and peribronchovascular regions with intact interlobular septa, and is always located adjacent to the pleura and septal lines (Hansell et al. 2008). Paraseptal emphysema is usually limited in extent, located in the upper lobes and clinically asymptomatic. Paraseptal emphysema is characterized at CT by subpleural low attenuation areas with a well-defined wall, and appears along peripheral pleura and fissures.

Figure 2.6: **A** Illustrations of radiologic patterns in CT images of centrilobular, panlobular and paraseptal emphysema. Cases courtesy of A.Prof Frank Gaillard, Radiopaedia.org, rID: 9225. **B** Axial computed tomography images for all emphysema subtypes. Source (Smith et al. 2014).

2.1.2.2 Interstitial Lung Diseases

Interstitial lung diseases (ILD) form a complex and heterogeneous group of over 200 different disorders which primarily exert their effects on the collagen and elastic connective tissue found in the interstitium, but which may present airway or vascular manifestations as well. ILD may be progressive, slowly progressive or static in their course. Some of them are of unknown etiology, but some of the most common ILD are caused by exposure to inhaled dust and particles, and exposure to toxic drugs, although they may be also caused by infections.

ILD affecting distal part of the alveoli cause physiologic restrictions and decreased lung volumes while when the interstitium closer to the proximal aspect of the acinus near the bronchioles is involved cause physiologic obstruction but without impact on lung volumes. Normally, lung volumes, including vital capacity and total lung capacity (TLC) are reduced in ILD. In general, ILD produce dyspnea with rapid and shallow breathing, and in severe cases, mild cyanosis may be observed.

High resolution CT is considered the imaging technique of choice for the detection and characterization of ILD. The pattern of the radiographic abnormalities on HRCT can help identify specific ILD.

The most common abnormal findings seen in high resolution CT that may characterize ILD include reticulation, ground-glass opacities, consolidation, nodules, linear opacities and honeycombing (Hansell et al. 2008)(see Figure 2.7).

Reticulation Refers to a netlike pattern which is the result of thickened interlobular and intralobular septa. Is most commonly seen in the peripheral and basal lung zones.

Ground-glass pattern Are defined as areas of increased density and attenuation with visible vascular structures. Ground-glass opacities are very often limited in extent to a certain segment or lobe.

Consolidation Appears as an area of increased density and attenuation, but it can be differentiated from ground-glass opacities by the inability to visualize vascular structures. The feature of pulmonary tissue consolidation is the replacement of air in the alveoli with fluid, such as edema or blood, cells, solid matter, or other factors.

Nodules Are focal areas of increased attenuation. Nodules are assessed in size, appearance, density, number and distribution. Nodules can be mainly categorized into three categories: centrilobular, perilymphatic and random. Centrilobular nodules are characterized by their central location in the secondary pulmonary lobules (nodules are not present on the pleura). Perilymphatic nodules are located peribronchovascularly, subpleurally and on the interlobular septa. Random nodules occur diffusely throughout the lung parenchyma and do not show a predominant distribution.

Linear Opacities Arise based on thickening of pulmonary interstitium and can include subpleural line and linear scar patterns. Subpleural line refers to a thin curvilinear opacity lying adjacent and parallel to the pleural surface. Linear scar are linear opacities which run through the lung parenchyma and usually extend from a visceral pleural surface.

Honeycombing Represents the destruction of lung parenchyma in advanced stage of pulmonary fibrosis. The appearance is of clustered subpleural cystic structures with a well-defined walls composed of fibrous tissue. Predominate on peripheral and subpleural lung regions.

Figure 2.7: Axial slices of HRCT scans representing the typical ILD patterns. **A** shows basilar predominant reticular pattern. **B** Ground glass opacity in the left lung. **C** Honeycombing in the form of well-defined walled cysts in the right lung. **D** Subpleural curvilinear line in the basilar portion of the left lung. **E** Linear scar opacity in the left lung. **F** shows consolidation in the upper zone of the right lung. **G** Random macronodules on the left lung. **H** Perilymphatic micronodules adjacent to the fissure.

2.1.2.3 Subclinical Interstitial Lung Abnormalities

There is a growing acceptance that some forms of ILD, especially idiopathic pulmonary fibrosis (IPF) may be preceded by early or subtle radiographic findings seen on CT scans of the chest (Lederer et al. 2009; Washko, Hunninghake, Fernandez, Nishino, Okajima, Yamashiro, Ross, Estépar, et al. 2011). The visual presence of these findings, often termed interstitial lung abnormalities (ILA), which include subtle ground-glass opacities, reticulation, linear opacities, honeycombing and centrilobular nodularity, might therefore represent ILD at its earliest stage, and has been shown to be associated with reduced lung volume, decrements in exercise capacity, increased mortality and a genetic polymorphism that is also associated with IPF (Doyle et al. 2012; Hunninghake et al. 2013; Putman et al. 2016; Seibold et al. 2011; Washko, D. A. Lynch, Matsuoka, Ross, Umeoka, A. Diaz, Sciurba, Hunninghake, Estépar, et al. 2010).

The relationship between emphysema and interstitial lung abnormalities is an area of increased interest. Recent studies demonstrated a significant association between paraseptal emphysema and interstitial lung abnormalities (Araki et al. 2015). The combined presence of interstitial features and emphysema was also associated with worse physiologic measures, reduced exercise capacity, worse respiratory health-related quality of life, and higher mortality than was emphysema alone. In addition, interstitial features intensify the effect of emphysema on those outcomes (Ash, Harmouche, Ross, A. A. Diaz, Rahaghi, et al. 2018), suggesting that emphysema is more deleterious in patients with interstitial features than in those with emphysema alone.

The rapid growth in use of CT scanning, combined with the high mortality associated with IPF and the fact that its available treatments slow but do not reverse the disease process, has led to a growing need for methods to accurately detect early interstitial changes before they progress to end stage fibrosis (Ley et al. 2011; Richeldi et al. 2014).

2.1.3 Quantitative imaging

Medical imaging encompasses different imaging modalities and processes to image *in vivo* anatomy and physiological processes for a better prognostic, diagnostic and treatment purposes, playing an important role in improving medical care for patients and public health.

There is growing consensus among the respiratory community that although patient histories and lung function testing are the minimum required for clinical examinations, these tests alone are not sufficient for disease characterization. Therefore, the use of computed tomography (CT) imaging is increasingly used in clinical decision making for lung diseases.

There is ample evidence that HRCT provides diagnostic and prognostic information in COPD and ILD. In practice, qualitative or semi-quantitative analysis based on visual assess-

ment has an inherent degree of subjectivity and is limited by significant inter- and intra-observer variability (D. Lynch et al. 2005; Mascalchi et al. 2012). In addition, is unable to detect subtle disease, as well as is insensitive to longitudinal change.

In this context, computer-based methods for quantitative analysis of disease on HRCT could provide rapid and objective measurement of disease extent and change over time. Imaging-based biomarkers could provide objective diagnostic information of the disease, as well as could constitute a prognostic factor informing on the outcome of the disease. Imaging-based biomarkers could also provide predictive information on the benefit from treatment, playing an important role in the identification of treatment efficacy and the identification of patients that are most likely to benefit from a given therapy.

2.2 State of the art in parenchymal injuries quantification

Conventional pulmonary function tests (PFT), such as spirometry, are quantitative and reproducible techniques. However, although they are cheaper than CT and avoid the use of harmful ionizing radiations, PFT are only available at one location and only provide a global assessment of the lung damage. The regional information provided by CT, along with its capability to differentiate the components underlying obstructive or fibrotic lung disease, has placed HRCT as the gold standard modality for lung disease evaluation and quantification.

HRCT is a highly sensitive imaging tool for the assessment of macrostructural changes in lung parenchymal injuries. Quantification of CT density changes, overall lung histogram features, parenchymal texture features and other assessments by advanced algorithms including machine learning and deep learning approaches to image analysis have the potential to standardise and develop the role of HRCT in lung parenchymal injuries quantification.

2.2.1 Lung density analysis

Quantitative measures of lung density are based on the observation that air-filled areas in the lung appear as areas of low attenuation in CT images, while fibrotic and interstitial changes in pulmonary parenchyma appear as areas of high attenuation.

Low attenuation areas (LAA) is a measure of lung density, defined as the percentage of the voxels in the lung below a certain threshold (Figure 2.10). Densitometric analysis based on LAA is a widely accepted technique for emphysema quantification. Although numerous density thresholds have been investigated, including -970HU, -960HU, -950HU and -910HU (Dirksen et al. 1997; Madani et al. 2006; Muller et al. 1988), the -950HU threshold has shown a greater correlation with microscopic measurements of emphysema (Gevenois et al. 1996), and is the most commonly used.

Figure 2.8: Low attenuation areas (LAA) below -950HU (LAA_{950}) are highlighted in red.

High attenuation areas (HAA) is a measure of the percentage of the voxels in the lung above a certain threshold, or between two thresholds (low and high threshold). Incipient interstitial lung abnormalities have been quantified as the lung area between -250HU and -600HU (Kliment et al. 2015; Lederer et al. 2009), while when fibrotic lesions are quantified, measures based on areas above -200HU or -500HU are preferred (Tanizawa et al. 2015; Umakoshi et al. 2017).

Figure 2.9: High attenuation areas (HAA) below -500HU (HAA_{500}) are highlighted in red.

The percentile index (Perc) is a well-established density measurement for estimating emphysema, although it has also been used for ILD quantification. Perc is defined as the value in HU below which that given percentage of all voxels is distributed. When emphysema extent increases, the density histogram shifts towards lower HU values and the given Perc index decreases. Conversely, when areas affected by ILD increase, the density histogram shift towards greater HU values, resulting in an increment in the Perc index. For emphysema quantification, Perc at 15% ($Perc_{15}$) is a commonly used index (Heussel et al. 2009). $Perc_{85}$

represents the most informative density parameter to assess ILD (Ninaber et al. 2015).

Kurtosis and skewness represent the distortion or disparity deviation of a histogram when compared with a normal distribution. Skewness is a measure of the lack of symmetry of the histogram, while kurtosis is a measure to which the distribution is peaked relative to a normal distribution. These measurements have been used for the study of COPD (Yamashiro et al. 2011), although it has been used most for ILD quantification (Best et al. 2003; Hartley et al. 1994). In lung fibrosis, both kurtosis and skewness typically decrease.

Figure 2.10: Density histogram indicating the appearance in normal, emphysematous and fibrotic lung, and the deviation of densitometry indexes.

2.2.2 Texture and Machine learning-based analysis

The first texture-based method for quantification of parenchymal disorders used a combination of texture features, including gray-level histograms (GLH), gray-level co-occurrence matrices (GLCM) and run-Length encoding (RLE) features, to describe two-dimensional regions of interest (ROI), along with a linear discriminant (LD) analysis (Delorme et al. 1997). GLH, GLCM, RLE features have been widely used for many years (Chabat et al. 2003; Malone et al. 2004; Uchiyama et al. 2003; J. Wang et al. 2009b).

AMFM (adaptive multiple feature method) was firstly proposed for the objective assessment of global and regional changes in pulmonary parenchyma to classify emphysema, using a combination of different features describing the texture and structure in two-dimensional (2D) regions of interest (ROI), along with a Bayesian classifier (Uppaluri, Mitsa, et al. 1997). AMFM was rapidly extended for discriminating between multiple pulmonary diseases processes, including emphysema and fibrosis (Uppaluri, Hoffman, et al. 1999). Further work on

AMFM allowed three-dimensional (3D) assessment for the differentiation of parenchymal pathology associated with both emphysema and interstitial lung diseases (Xu et al. 2006).

Gabor filters have been also used along with a k-nearest-neighbors (KNN) classifier to classify diffuse lung opacities (Mitani et al. 2000), as have 2D fast Fourier transform (FFT) coefficients to represent the texture features and discriminate between normal and pathological tissue (Liu et al. 2001). Abnormality detection from patients with ILD was also performed extracting features to describe local image structure by means of a multi-scale filter bank and subsequently classified using various algorithms including LD classifier, quadratic discriminant (QD) classifier, support vector machine (SVM) and a KNN classifier (Sluimer et al. 2003).

Local binary patterns (LBP) and joint LBP and intensity histograms are used for characterizing regions of interest (ROI), along with a KNN classifier to distinguish between normal and 2 types of emphysema (Sorensen et al. 2010a). A combination of LBP with other intensity, shape and context features has been used to describe the appearance of ground glass nodules (Jacobs et al. 2011).

Simpler features and first order spatial statistics, such as kernel density estimation of local histogram (LH) has been also proposed to discriminate emphysema subtypes, rather than higher order spatial statistics such as LBP (Castaldi et al. 2013; Mendoza et al. 2012). Local histogram-based measures have been also recently used for the objective identification and quantification of both subclinical ILA and pulmonary fibrosis (Ash, Harmouche, Vallejo, et al. 2017; Ash, Harmouche, Ross, A. A. Diaz, Hunninghake, et al. 2017).

More recent approaches have proposed the use of features learned directly from the data, instead of hand-crafted ones. Some of these methods are based on unsupervised learning algorithms to construct a set of learned features such as restricted Boltzmann machine (RBM) for ILD classification (Li et al. 2013), k-means algorithm for emphysema discrimination (Gangeh et al. 2010), and the application and optimization of the sparse representation (SR) for the classification of various pulmonary patterns (Zhao et al. 2015). Data-driven textural analysis (DTA) is able to use clustering analysis to learn consistent patterns and generate a dictionary of ILD elements (Humphries, Yagihashi, et al. 2017).

Multiple deep learning approaches have been used to prognosticate and characterize disease. Convolutional Neural Networks (CNN) have been trained for a number of purposes, including characterization of disease severity and prediction of clinical outcomes (Gonzalez, Ash, et al. 2018), pulmonary artery-vein separation (Nardelli et al. 2018), biomarker regression (Gonzalez, Washko, et al. 2018) or pulmonary fissure detection (Gerard et al. 2019).

For ILD characterization, there have been proposed several CNN-based methods for auto-

matic classification of patients with fibrotic lung disease (Walsh et al. 2018), for classifying the image-level label of ILD types (C. Wang et al. 2019) and for classifying whole lung slices (Gao et al. 2018). On the other hand, ILD radiographic subtype classification has been addressed by proposing specific CNN architectures (Anthimopoulos et al. 2016; Christodoulidis et al. 2017) or by using well-known and standard models such as AlexNet and GoogleNet (Shin et al. 2016).

Emphysema classification and quantification have also been addressed using CNN (Hatt et al. 2018; Karabulut et al. 2015; Pei 2015), as have the integration of hand-crafted features into a CNN (V. Srivastava et al. 2019). Variations from standard CNN architectures have been proposed, including the incorporation of multi-scale information into the CNN to take full consideration of the characteristics of different emphysema (Peng et al. 2018), as well as the combination of CNN and recurrent neural networks (RNN) capable of learning dependencies in sequence data (Humphries, Notary, et al. 2018).

3

Emphysema classification

The association between tobacco smoking and the development of emphysema has been well documented. Although cigarette smoke is not the only cause of emphysema, approximately 20% of smokers develop emphysema. In this chapter we propose an automatic method to identify and classify different subtypes of emphysema in HRCT images. This method could be directly used for emphysema subtype quantification.

3.1 Introduction

Chronic Obstructive Pulmonary Disease (COPD) is the fourth leading cause of death worldwide, and it is estimated to be the third one in a few years, which has led to an increasing interest of population studies that allow a better understanding of its pathophysiology. COPD is defined as persistent obstruction of the airways. It can be divided into two main phenotypes: chronic bronchitis and pulmonary emphysema that causes an overall decrease in the lung elasticity affecting the lung tissue. The progression of emphysema has traditionally been evaluated by spirometric pulmonary function tests (PFTs), although nowadays it is considered as a non-specific parameter and non-sensitive to very early stages of the disease.

Densitometric analysis in chest CT is widely accepted as an alternative measurement of pulmonary emphysema more specific and sensitive than PFTs for *in vivo* studies (Cavigli et al. 2009). This method is based on the choice of a Hounsfield threshold inside the lung mask to discriminate between emphysematous and non-emphysematous tissue. Although a densitometric approach may be able to quantify the disease extension, it may not be able to classify it into different subtypes. That is why other methods based on texture information (Depeursinge et al. 2007; Y. S. Park et al. 2008; Uppaluri, MITSA, et al. 1997) or local histograms (Castaldi et al. 2013) have been proposed to carry out an emphysema subtype classification.

Recently, two approaches have been published to tackle the parenchyma patterns classification problem employing CNNs (Anthimopoulos et al. 2016; Christodoulidis et al. 2017). In (Anthimopoulos et al. 2016), a specific CNN to classify 2D patches into interstitial lung patterns was proposed, whereas (Christodoulidis et al. 2017) used the same CNN proposed in (Anthimopoulos et al. 2016), but pre-trained with a variety of texture data.

In this chapter we aim to compare these previously proposed CNN architectures in the emphysema subtyping problem with respect to 3D, 2.5D (multi-view) and 2D simpler architectures.

3.2 Methods

In this chapter, we proposed a CNN for objective and automated classification of emphysema patterns on chest CT images, considering a total of six radiographic tissue patterns: normal parenchyma (NP) and five emphysematous subtypes (paraseptal (PS), panlobular (PL) and mild, moderate and severe centrilobular (CL1, CL2, CL3) emphysema).

Deep architectures usually include hundreds of thousands of parameters, imposing considerable computation and a large number of training samples. In this chapter, we propose a relative simple architecture with a reduced number of parameters that tackle the emphysema classification problem considering a 2.5-dimensional representation of the input data. To

Figure 3.1: Proposed multi-view CNN (2.5D CNN) for emphysema classification. 2D and 3D architectures are the same as the 2.5D network, except for the input dimensionality, and the convolutional operations in the 3D case. Filter size and pooling size in convolutional (CONV) and max-pooling (MAX) layers specify their size on each dimension (2D or 3D depending on the architecture).

justify the usage of the proposed architecture, we compared it to different 2D- and 3D- CNNs, as well as the CNN proposed in (Anthimopoulos et al. 2016). We also compared the proposed method with a state-of-the-art work for emphysema subtypes discrimination that uses local intensity probability distribution functions subsequently classified with a KNN classifier (Castaldi et al. 2013).

3.2.1 Multi-view Convolutional Network (2.5D CNN)

3.2.1.1 Architecture

The proposed multi-view CNN architecture is shown in Figure 3.1, and it is composed of 4 convolutional, 3 max-pooling layers and 3 fully-connected layers. The input of the network consists of a multi-view (2.5D) representation of pulmonary segment tissue under study. Therefore, the input is comprised of three 2D patches of size 31x31 pixels, corresponding to the axial, sagittal and coronal views.

Emphysematous patterns in CT images are mainly characterized by local texture patterns. To capture these local patterns and to reduce the complexity of the network, and hence the number of parameters, we use relatively small kernels in the convolutional layers and reduce the number of pooling operations through the network compared to standard convolutional networks.

The network training is based on an optimization problem that minimizes the loss function. In this work we use the Stochastic Gradient Descent (SDG) method updated with Nesterov momentum (Bengio et al. 2013) to minimize the categorical cross entropy.

3.2.1.2 Overfitting prevention

Convolutional Neural Networks easily overfit the training data. In this work we apply 4 different techniques to prevent and reduce the overfitting. First, we have applied a regularization of the loss function. L2 regularization penalizes the square magnitude of the parameters (w) in the loss function by adding the term $1/2\lambda w^2$, where λ is the regularization strength. This method penalizes sharp changes in the parameters preferring soft ones. The second technique employed is data augmentation. We apply data augmentation using 7 different spatial transformations by rotating 90, -90 and 180 degrees, flipping along horizontal and vertical axes and combining both transformations over the training dataset. Another prevention technique is early stopping, where the training stage is stopped before overfitting process begins. Finally, we have applied dropout to prevent the overfitting, by randomly dropping units with a given probability ($p=0,5$) from the network during training.

3.2.2 Comparative methods

3.2.2.1 2D and 3D CNN

For comparative purposes, we also designed the 2D and 3D versions of the proposed architecture. Both networks have the same configuration as the one proposed with the only difference that the convolutional operations in the 3D CNN are made in three dimensions.

For the 2D CNN, the input consist of an image patch of size 31x31 pixels in the axial plane, whereas for the 3D CNN, the input corresponds to a 3D segment of size 31x31x31 pixels (see Figure 3.1).

3.2.2.2 State-of-the-art CNN

To justify the usage of the proposed CNN we also re-implemented the CNN proposed in (Anthimopoulos et al. 2016), where the authors focused on detecting patterns of interstitial lung diseases from 2D patches. The input of the CNN is a 32x32 image patch which is convolved by a series of 5 convolutional layers with 2x2 kernels, and followed by one average pooling with size equal to the size of the final features maps. Finally, the classification stage is comprised of 3 fully-connected layers.

The authors proposed the use of LeakyReLU as a non-linear activation function for both convolutional and fully-connected layers, and Adam optimizer for minimizing the categorical cross-entropy during training.

Table 3.1: Description of the dataset and distribution along different emphysema patterns.

Lung pattern	NP	PL	PS	CL1	CL2	CL3
Num. Samples	447	250	152	223	299	182

3.3 Experiments and results

3.3.1 Dataset and evaluation setup

267 CT scans were selected from the COPDGene study. CT scans were acquired with equipment from 3 vendors and a total of 9 different models. Through these scans, an experienced pulmonologist manually placed 1553 points corresponding to six different radiographic patterns (see Table 3.1). This database has been previously used for automated emphysema classification in (Mendoza et al. 2012). Of all of these points, 390 were randomly selected for testing (around 25%), while the remaining points were randomly split using a 10-fold cross validation scheme leading to the training and validation sets. The final evaluation of the proposed method as well as a comparison with other CNNs and state-of-the-art methods were carried out on the test set. Training sets was used for training the algorithms, while the validation sets were used to hyper-parameter tuning of the proposed architecture.

To evaluate the methods, we consider the sensitivity (SN), specificity (SP), geometric mean (GM) and balanced accuracy (BA), defined as:

$$\begin{aligned}
 SN &= \frac{TP}{TP + FN} \\
 SP &= \frac{TN}{TP + FP} \\
 GM &= \sqrt{TP * TN} \\
 BA &= \frac{SN + SP}{2}
 \end{aligned} \tag{3.1}$$

, where TP, FP, TN and FN stand for true positives, false positives, true negatives and false negatives rates respectively.

3.3.2 Validation

The proposed multi-view CNN, as well as other CNNs are trained on Regions of Interest (ROIs) extracted around the manually labeled points. The size of the ROIs are selected in accordance with each network architecture as described in Sections 2.1.1 and 2.2.1.

The only pre-processing performed on the dataset was subtracting the mean and dividing by the standard deviation computed on the training and validation sets to normalize the image ROIs.

The framework used in this work to train the CNNs is based on Theano and Lasagne

Table 3.2: Comparison of the proposed CNN (bold) with other CNNs architectures. Results are computed on the independent test set. Evaluation metrics (SN, SP, GM, BA), the number of parameters involved and the time needed to train each CNN are reported.

Method	2D	2.5D	3D	Lung-CNN (Anthimopoulos et al. 2016)
SN(%)	78.52	81.78	82.52	77.20
SP(%)	96.79	97.34	97.10	96.45
GM(%)	87.18	89.22	89.51	86.33
BA(%)	87.66	89.56	89.81	86.86
#Params	68918	69494	157558	467426
Time(s)	19.68	20.48	240.4	137.3

libraries using a PC with GPU NVIDIA TITAN X Pascal 12GB, CPU Intel Core i7 3.6 GHz and 32GB of RAM.

3.3.2.1 Comparison to other CNNs

Table 3.2 provides a comparison of the proposed multi-view CNN with other CNNs, including the 2D and 3D version of the latter, as well as the network proposed in (Anthimopoulos et al. 2016). All the networks were trained with the same samples, and tested in the same test set. As derived from the Table 3.2, the proposed 2.5D CNN achieves analogous results to those obtained using the 3D CNN, while keeping the same small number of parameters and the same training time as in the 2D CNN case.

The proposed method also outperforms the CNN proposed in (Anthimopoulos et al. 2016), and considerably reduces the computational cost by a factor around 6 to 7 in terms of number of parameters involved in the architecture and training time.

3.3.2.2 Comparison to Local Histogram

As an additional comparative study, we also compared our proposed method against a reference method in emphysema classification. This method has been previously detailed in (Castaldi et al. 2013). Briefly, the local intensity distribution of a two-dimensional ROI of size 31x31 pixels is computed using a kernel density estimator (KDE), and a k nearest neighbors (kNN) classifier assigns each ROI local histogram to its corresponding emphysema pattern based on a majority consensus from the k nearest training samples.

Our proposed method outperforms the results achieved by (Castaldi et al. 2013), which obtained a sensitivity (SN) of 44.75%, specificity (SP) of 88.39%, geometric mean (GM) of 60.65% and balanced accuracy (BA) of 66.57%. Although the training time of (Castaldi et al. 2013) is lower than the time needed to train the proposed CNN, the classification time is much higher due to the need to compute the KDE features.

True label	NP	97.0	0.0	0.0	3.0	0.0	0.0
	PS	1.5	95.5	1.5	0.0	1.5	0.0
	PL	0.0	2.6	65.8	0.0	0.0	31.6
	CL1	1.6	0.0	0.0	82.0	16.4	0.0
	CL2	0.0	3.7	0.0	6.2	85.2	4.9
	CL3	2.4	2.4	4.8	4.8	21.4	64.3
		NP	PS	PL	CL1	CL2	CL3
		Predicted label					

Figure 3.2: Confusion matrix of the proposed method.

Figure 3.3: ROC analysis for the proposed CNN and state-of-the-art approaches. The ROC curves are obtained by macro-averaging over all classes. The area under the curve (AUC) are also reported.

3.3.2.3 Analysis of the method's performance

Figure 4.6 shows the confusion matrix of the proposed 2.5D CNN. As can be seen, misclassifications generally occurred within the centrilobular emphysema classes, due to their common nature that differ only by the disease stage. Confusion error also occurred between severe centrilobular (CL3) and panlobular (PL) emphysema, since both classes involve a large parenchyma destruction.

Figure 3.3 shows the macro-average ROC curves and the area under the curve (AUC) for the proposed CNN and state-of-the-art methods, where it can be seen that the proposed method achieved the highest AUC.

For a further validation, we performed full-lung classification on a severe emphysema CT

Figure 3.4: Full-lung classification results for a severe emphysema case.

scan (Figure 4.10). The classification was carried out at a sampling grid of $5 \times 5 \times 5$ pixels in each axial image slice. The rest of the voxels were classified using nearestneighbor interpolation. An expert evaluator confirmed the good agreement of the resultant classification. However, it should be noted that although the identification of paraseptal emphysema obtained a high performance in the test set, the proposed methodology usually results in PSE over-estimation in the vicinity of high-intensity areas (i.e. in the proximity of the pleura). This suggests that the identification of PSE lesions require further investigations and additional work needs to be done to properly identify this particular type of emphysema.

3.4 Conclusions

In this chapter, we present a relatively simple Convolutional Neural Network (CNN) architecture for automated emphysema classification in CT images, considering six different tissue classes, including normal parenchyma and five emphysematous patterns. The proposed architecture, which is composed of 4 convolutional and 3 pooling layers, is considerably simpler than other deeper state-of-the-art architectures proposed in the literature that are specific for lung pattern classification (Anthimopoulos et al. 2016), reducing the computational cost by a factor around six to seven, and outperforming the results.

Additionally, we showed that the 2.5D proposed approach is optimally cost effective, since it is able to reflect 3D information and achieve similar results to those obtained with 3D CNNs, and have similar computation cost compared to 2D approaches.

To train and test the proposed method, we used 1553 manually labelled tissue samples corresponding to six different tissue classes. Visual inspection of full-lung classification and results in the test set (sensitivity of 81.78%, specificity of 97.34%) prove the potential of the method for emphysema detection and classification.

4

Parenchymal injuries classification

Cigarette smoking is not only associated with the development of emphysema but more generally with a variety of effects on the lung parenchyma. In this chapter, we profoundly extended the method presented in Chapter 3 proposing a robust method for the automatic identification and classification of early parenchymal lung disease including interstitial lung disease at an early stage as well as different subtypes of emphysema.

4.1 Introduction

Chronic cigarette smoke exposure is associated with a variety of effects on the lung parenchyma. While there are other, more uncommon manifestations, generally, these effects can be characterized as those associated with destruction of the parenchyma (Auerbach et al. 1974; Ryu et al. 2001) leading to a local decreases in lung density typically called emphysema, and those associated with inflammation and scar formation leading to local increases in lung density, typically called as interstitial changes. In fact, these manifestations can co-exist even within the same individual.

Interstitial Lung Diseases (ILD) are a heterogeneous group of more than 200 lung disorders that largely affect the lung parenchyma but which may present airway or vascular manifestations as well. There is a growing acceptance that some forms of ILD, especially idiopathic pulmonary fibrosis (IPF), may be preceded by interstitial lung abnormalities (ILA), which are early or subtle radiographic findings seen on computed tomography (CT) scans of the chest (Lederer et al. 2009; Washko, Hunninghake, Fernandez, Nishino, Okajima, Yamashiro, Ross, Estépar, et al. 2011). The visual presence of ILA has been shown to be associated with reduced lung volume, increased mortality and a genetic polymorphism that is also associated with IPF (Doyle et al. 2012; Hunninghake et al. 2013; Putman et al. 2016; Seibold et al. 2011; Washko, D. A. Lynch, Matsuoka, Ross, Umeoka, A. Diaz, Sciurba, Hunninghake, Estépar, et al. 2010). The rapid growth in use of CT scanning, combined with the high mortality associated with IPF and the fact that its available treatments slow but do not reverse the disease process, has led to a growing need for methods to accurately detect early interstitial changes before they progress to end stage fibrosis (Ley et al. 2011; Richeldi et al. 2014). While visual CT analysis remains the mainstay of clinical imaging interpretation, there has been an increasing recognition of the potential role of objective analysis techniques to quantify abnormalities on CT and characterize disease subtypes (Ash, Harmouche, Ross, A. A. Diaz, Hunninghake, et al. 2017).

In this chapter we propose a method for objective and automated identification and classification of ILA and emphysematous patterns on chest CT scans, based on Convolutional Neural Networks (CNNs), considering a total of eight different radiographic patterns: normal parenchyma, five interstitial patterns (ground-glass, reticular, nodular, linear scar, subpleural line) and two emphysematous patterns (centrilobular and paraseptal) as depicted in Fig. 4.1.

4.1.1 Related work

There have been several approaches to computer-aided analysis and automated identification and classification of ILD using CT images. For example, in patients with IPF, densitometric measures, such as the skewness, kurtosis and the mean of the histogram of distribution of

Figure 4.1: Examples of patches centered on selected points along different ILA patterns, including normal parenchyma (NP), ground-glass (GG), reticular (RETIC), nodular (NOD), linear scar (LINSCL), subpleural line (SUBPL), paraseptal emphysema (PS) and centrilobular emphysema (CL).

lung density, have been found to be associated with outcomes such as pulmonary function and transplant free survival (Matsuoka et al. 2015a; Shin et al. 2011). Hand-crafted texture features such as gray level co-occurrence matrices (GLCM), run length matrices (RLM), and fractal analysis, have been also widely used (Lim et al. 2011; S. O. Park et al. 2011; J. Wang et al. 2009a) for classification of patterns of interstitial change as have local binary patterns (LBP) (Sørensen et al. 2010) and local histogram-based measures (Ash, Harmouche, Vallejo, et al. 2017; Ash, Harmouche, Ross, A. A. Diaz, Hunninghake, et al. 2017). For example, Park et al proposed a CAD scheme to detect, but not classify, early ILD using low-dose CT images based on hand-crafted features such as RLM and histogram features on 30 patients (S. C. Park et al. 2011).

There have also been proposed approaches to learn the features directly from the data. Some of these methods are based on unsupervised learning algorithms such as restricted Boltzmann machine (RBM) (Li et al. 2013) or k-means and k-SVD to construct a set of learned features (Gangeh et al. 2010; Zhao et al. 2015). However, the majority of the proposed methods for learning features from the data are supervised algorithms based on Convolutional Neural Networks (CNN). There have been proposed CNN-based methods for automatic classification of patients with fibrotic lung disease (Walsh et al. 2018) and classifying the image-level label of ILD types (C. Wang et al. 2019). There has also been an attempt to classify in an holistic manner whole lung slices by using a pre-trained and fine-tuned AlexNet architecture (Gao et al. 2018). However, a holistic classification of a whole slice as a unique label does not consider multiple

tissue subtypes in the same slice thus it only provides a rough quantification method of the disease. Regarding ILD radiographic subtype classification, other recent notable approaches include the use of a variety of CNNs. These include CNNs to classify 2D patches as interstitial patterns with or without pre-training using a variety of texture datasets (Anthimopoulos et al. 2016; Christodoulidis et al. 2017; Kim et al. 2018) as well as the use of the well-known architectures such as AlexNet and GoogleNet pre-trained on ImageNet (Shin et al. 2016).

One potential approach for improving the performance of all of these techniques is to combine multiple classifiers into an ensemble. While this has not been attempted for ILD classification, others have used this approach for the identification of other lung structures such as nodules. These efforts have included both the combination of multiple CNN with the same configuration, each of them having as input a different nodule view (Setio et al. 2016) and the combination of MLP, KNN and SVM to classify hand-crafted features for nodule diagnosis (Farahani et al. 2015).

In this work we propose the first deep learning-based method to identify and classify radiographic patterns of ILA, which likely represents early or subtle ILD in some cases, that implies the characterization of 8 different parenchymal features types. We propose a methodology based on an ensemble of various 2D, 2.5D and 3D CNN specific architectures to tackle subtle parenchymal patterns, including multi-context and multi-stage architectures, proving the superiority with respect to previous methods specifically designed to address late-stage interstitial diseases revealing the need of specific designs and research to tackle ILA properly. Additionally, this work precisely defines how to perform and optimize the ensemble of different CNN architectures which has rarely been addressed.

4.2 Methods

4.2.1 Database

The CT scans used in this study for both training and evaluating the proposed method were acquired as part of the COPDGene study, a previously described (Regan et al. 2010) multi center study designed to identify genetic and epidemiologic factors associated with COPD. The COPDGene study was approved by Partners Human Research Committee (Protocol Number 2007-P-000554/2) and institutional review boards at all study sites, and written informed consent was obtained from all subjects. All research in this study was performed in accordance with relevant guidelines and regulations.

The scans utilized for this study were acquired at full inspiration and were obtained with a total of 9 CT scanner models from 3 manufactures. From each CT scan, images were acquired using two different reconstruction kernels, namely a soft filter (B35, Smooth Recon) and a

Table 4.1: Description of the database

ILA radiographic pattern	Num. Points	Num. Scans
Normal parenchyma	23696	161
Ground-glass	137	9
Reticular	5409	58
Nodular	116	7
Linear scar	195	8
Paraseptal emphysema	3845	43
Centrilobular emphysema	3613	51
Subpleural line	413	19

shaper one (B50, Sharp Recon). Two pulmonologists manually placed a total of 37427 training points in the scans of 208 randomly selected individuals. These points were placed throughout the lungs of the participants and included the following parenchymal features types: normal parenchyma, interstitial features including ground glass, reticular, nodular, linear scar and sub pleural line, and emphysematous features including centrilobular and paraseptal emphysema (Table 4.1). This list of features was felt to represent the majority of parenchymal tissue types in the cohort and all of the feature types were included, so that, during the classification process, the entire lung volume could be classified as a particular subtype. That is, no volume was classified as other or unclassifiable. Of note, normal parenchyma was labeled both distant from and adjacent to the other tissue subtypes, but only in areas that were clearly visually normal. Additionally, no pan-lobular emphysema was identified in the training cohort, likely due to the lack of alpha-1-antitrypsin disease, and thus this feature was excluded.

4.2.2 Multi-model ensemble

In order to address the problem of radiographic ILA pattern classification from CT images, we designed a multi-model ensemble of deep convolutional neural networks. Ensemble strategies allow combining the individual predictions of a set of classifiers following some defined criterion such as weighted averaging or majority voting rule. The combination of multiple independent models usually performs better than the independent models separately (Rokach 2010). Individual networks work well on finding locally optimal solutions of the true function that maps the input image space to the output label space. Ensemble methods show their advantage by starting the optimal solution search of the true function from different locally optimal points. The ensemble algorithm can then combine different information from each individual model to construct a better representation of the input therefore enabling a better mapping function.

In the proposed ensemble method, each individual network is trained from scratch using the same training samples. Then, the output probability distribution responses of the networks are combined using a weighted averaging to form the overall output of the ensemble:

$$Y(x) = \sum_{i=1}^n w_i P_i(x) \quad (4.1)$$

Where x is the input image and w_i is the weight assigned to the individual response of the i th network, $P_i(x)$. We use a heuristic search of the optimal values of the weights applied to the individual networks. The weights are randomly initialized and iteratively evolve so that the overall performance of the resulting ensemble improves. We solve this optimization problem by minimizing the overall error of the ensemble using the Tree-structured Parzen Estimator (TPE) (Bergstra et al. 2011), an improved version of Sequential Model-based Algorithm Configuration (SMAC). Briefly, TPE constructs the surrogate function $p(Y|w)$, a probabilistic model of the objective function which maps the hyperparameters w to a probability of the score Y , by applying the Bayes' theorem. Instead of modeling the surrogate $p(Y|w)$ directly, TPE models $p(w|Y)$ and $p(Y)$, where $p(w|Y)$ is modeled by two non-parametric density functions based on observations w .

The final predicted label is based on the optimal decision rule, where the predicted class is that which has the highest probability.

The proposed ensemble is comprised of seven Convolutional Neural Networks: three 2D CNNs, two 2.5D CNNs and two 3D CNN, as detailed in Figure 4.2.

Radiographic ILA patterns are mainly characterized by local texture patterns in CT images. In order to capture these local texture patterns, our baseline architecture uses relative small kernels in convolutional layers in order not to introduce non-local information in the receptive field. We also reduced the number of pooling layers through the network compared to standard CNNs to avoid spatial information loss. In view of these considerations, and considering that shallow CNNs do not have enough discriminative power while too deep architectures are computationally expensive to train and can easily overfit the training set, our baseline architecture (BCNN^{2D}) for the classification of ILA radiographic patterns in HRCT images is a deeper variant of the well-known LeNet configuration, where each convolutional layer with 5x5 kernels are replaced with two layers with 3x3 kernels. We also introduced a batch normalization layer after each pooling operation to accelerate the network training and improve the overall performance.

In order to introduce architectures that exploit contextual information into the ensemble, we designed the multi-stage and multi-context versions of the baseline architecture

(MSTAGE-CNN^{2D} and MCONTEXT-CNN^{2D} respectively). Multi-stage networks exploit contextual information by branching middle-level features from the CNN and concatenating them with the last features before being fed into the classification step, so that the network has richer representation of the input image with different scales of the receptive field. This configuration combines, at the classification step, high-level features from the latter stage with global information, and low-level features that represent local structures from the image so that the loss of spatial relationship between global features is avoided. Multi-context networks also add a secondary path to the architecture that parallel processes a larger scale input image so that the network simultaneously learns features from the input image at multiple scales.

To take advantage of spatial information, we designed the respective 2,5D and 3D versions of the baseline and multi-stage networks previously presented (yielding the BCNN^{2.5D}, MSTAGE-CNN^{2.5D}, BCNN^{3D} and MSTAGE-CNN^{3D} configurations), though the 2,5D and 3D multi-context versions were discarded due to their inadequate performance and convergence. 2,5D networks explore two-and-a-half dimensional (2,5D) representation of the image, learning features and capturing information from the three orthogonal planes, namely axial, sagittal and coronal slices. It has been shown that a two-and-a-half dimensional representation of pulmonary segments is optimally cost effective and has been claimed to be more suitable compared to more complex architectures (see Chapter ??). A natural extension of 2,5D architectures are 3D CNNs which have the ability to model and capture volumetric contextual information around the voxel to classify.

All the architectures of the CNNs that comprise the proposed ensemble are described in details in Figure 4.2.

The training of the CNNs is based on an optimization problem that minimizes a loss function. In this work we use Adam optimizer to minimize the L2-regularized categorical cross entropy.

4.2.3 Overfitting prevention

Deep architectures, which involve a large number of parameters, easily overfit the training data. In this work we apply four different techniques to prevent and reduce the overfitting.

In order to increase the number of training samples and to consider variability through samples from the same class, we used two approaches to augment the training data by modeling both geometric and physiological transformations. Geometric transformations on training samples belonging to the classes with less training data (i.e. ground glass, nodular, linear scar and subpleural line) were used, combining rotation, horizontal and vertical shifts and flips and shearing. Additionally, in all samples of the training data, we also performed a domain-specific data augmentation to model the differences of tissue density depending

Figure 4.2: Architectures of the CNNs included in the proposed ensemble. Each network is fed with a 2D, 2.5D or 3D ROI extracted from a sample point.

on the level of inspiration, which is frequently needed when estimating pulmonary diseases in CT images (Camiciottoli et al. 2013). In this work, we implemented a new approach to augment the training data modeling variations in lung density and volume due to differences on inspiration levels among patients in order to be robust and relatively independent of the respiratory state of the lungs at acquisition time. Our approach is based on the so-called sponge model of the lung (Stoel et al. 2008) which assumes mass preservation over the lung respiratory cycle causing that a proportional decrease in lung volume would yield an equally proportional increase in density. We have modeled random lung volume variations of ± 100 milliliters (mL) $—\mathcal{U}(-1000, 1000)—$, corresponding to mild fluctuations in inspiratory lung volume.

On the other hand, we have also applied a regularization of the loss function for overfitting prevention. L2 regularization penalizes the square magnitude of the parameters (w) in the loss function by adding the term $1/2\lambda w^2$, where λ is the regularization strength. This method penalizes sharp changes in the parameters, preferring soft ones. Dropout technique (N. Srivastava et al. 2014) was also used to prevent the overfitting by randomly dropping units with a given probability p from the network during training. Finally, we also employed early stopping technique, where the training stage ends before overfitting process begins. The training finishes when the error on the validation set (validation loss) does not decrease for a preset iteration epochs (patience).

4.3 Experiments and results

4.3.1 Experimental setup

4.3.1.1 Implementation details

All individual networks from the proposed ensemble method were trained on Regions of Interest (ROIs) extracted around the manually labeled points. The size of the ROIs was selected in accordance with each network architecture, always ensuring the coverage of a whole secondary pulmonary lobe (functional and anatomical unit of the lung), leading to ROIs of 30 mm^2 mean size for the 48×48 axial patches. As derived from Figure 4.2, we extracted two-dimensional ROIs from the axial plane for 2D-CNNs, while two-and-a-half-dimensional ROIs (from axial, sagittal and coronal planes) and three-dimensional ROIs were extracted for 2.5D-CNNs and 3D-CNNs respectively. Of the two available images per scan, the dataset was built with ROIs extracted from the CT images reconstructed with the B50 filter.

The only pre-processing performed on the dataset was normalizing the attenuation by subtracting the mean and dividing by the standard deviation tissue density computed on training and validation sets to normalize the image ROIs.

The framework used to implement the proposed method was based on Theano and Lasagne libraries using a PC with GPU GeForce GTX TITAN X Pascal 12GB, CPU Intel Core i7 3.6 GHz and 32GB of RAM.

4.3.1.2 Evaluation

To evaluate all proposed and state-of-the-art methods for the classification of ILA patterns, we constructed a train-validation-test scheme. For radiographic tissue classes with 800 or more training points, we randomly selected 600 samples (maximum of 75%) to be part of the training set, while for those classes with less than 800 points (i.e. ground glass, nodular, linear scar and subpleural line), 75% of the points were randomly separated for the training set and augmented with geometric transformations until all classes had 600 points in order to build an equally distributed training set. These training data were then split using a 10-fold cross validation scheme leading to create the final training and validation datasets, and all the CNNs were trained and evaluated 10 times according to the 10-fold cross validation split. For all tissue classes, the remaining samples not considered in this selection were defined as the test set (minimum of 25%). The training of the models was carried out on the training sets, and the validation sets were used for hyper-parameter tuning. The final performance evaluations of the proposed and state-of-the-art methods were carried out on the test sets.

Additionally, and for further validation, we performed an additional experiment with a train-test split at patient level, creating a completely independent test set without overlap between samples from the same CT scan in both the training and the new independent test sets. This experiment allows us to check the repeatability of our results at the subject level assuming that disease presence within a subject is somehow correlated between lung regions. From the 254 CT scans described in Database section, we randomly selected 5 patients for testing ensuring that all eight tissue classes under study were represented, and the rest of the scans were used for training, leading to 4800 training points and 2330 samples for testing.

We employed five point-metrics to evaluate and compare the proposed and baseline classification methods. To deal with the skewed class distribution of the test dataset and to equalize the importance of each class, we computed the point-metric for each class, and estimated their unweighted average. We considered the sensitivity (SN), specificity (SP), geometric mean (GM) and balanced accuracy (BA), defined as:

$$SN = \frac{TP}{TP + FN} \quad SP = \frac{TN}{TN + FP} \quad GM = \sqrt{TP * TN} \quad BA = \frac{SN + SP}{2} \quad (4.2)$$

where TP, FP, TN and FN denote the number of true positives, false positives, true negatives and false negatives respectively.

Table 4.2: Fine-tuned hyperparameters

Hyper-parameter	Value
Optimizer	Adam updates $\eta=0.001, \beta_1=0.9, \beta_2=0.999, \epsilon=10e-08$
L2 regularization strength	0.0025
Patience (early stopping)	30
Dropout rate	0.5
Batch size	128

4.3.2 Validation results

In the following subsections we describe a set of experiments performed to validate the proposed method. First, we compared the proposed ensemble to the individual networks. Second, we evaluated the generalizability of the method when it is applied to data with characteristics different than those of the training set. Next, we performed an overall analysis of the system's performance with an independent test dataset. Finally we performed a comparison of our techniques with previous studies, as well as an experiment to investigate the performance of our method when it is applied to the full lungs with sliding ROIs.

4.3.2.1 Ensemble of models

We first compared the performance of individual networks to the proposed multi-model ensemble. To identify the optimal architectures of individual networks we experimented with different configurations, resulting in the architectures shown in Figure 4.2. To optimize the training procedure we also experimented with different hyper-parameters, but for purposes of comparative evaluation, we used the same parameter set for all individual CNNs of the ensemble. In Table 4.2 we summarize these hyper-parameters tuned via cross-validation. The weights w assigned to the individual response of each network of the ensemble (Eq. 4.1) were also optimized via cross validation.

Figure 4.3 compares the proposed ensemble with the individual networks performance by giving the unweighted average along all classes for each evaluation metric. It can be seen that the ensemble achieved a significantly (two-sample t-test: p -value<0.001) better performance compared to isolated models by combining its individual responses, outperforming in terms of all considered metrics the individual CNNs, and reducing the standard deviation of the mean of each metric.

Table 4.3 demonstrates the effect of using different strategies to aggregate the individual networks to build the ensemble, including weighting and non-weighting methods. For weighting methods, we used different optimization algorithms to optimize the weights. Minimizing

Figure 4.3: Classification performance of each individual models (gray) and combined in the ensemble (orange). The two-samples t-test to assess the statistical significance of the improvement in performance of the ensemble compared to the individual networks are represented with blue asterisks (***)two-sample t-test: p -value<0.001). SN: sensitivity, SP: specificity, GM: geometric mean, BA: balanced accuracy.

the overall error of the ensemble using cross-validation, we concluded that all the methods yielded roughly the same performance, with TPE providing the best results, followed by Bayesian Optimization (BO) and Random Search (RS). When Majority Voting rule (MVR) was used to combine the individual responses it also obtained a higher performance compared to individual models, but all weighting methods scored better than the latter.

As shown in Table 4.4, there was considerable variation between the learned weights of each of the individual networks using the different optimization algorithms, although all performed similar (Table 4.3). This may be due to the fact that all algorithms find different but equivalent local minima and close to the same global minima.

4.3.2.2 Evaluation of generalization capability

Because CNNs learn features from their training data and it is important to determine if they are generalizable to data with characteristics that are different than those of the training dataset. For CT imaging data this may include diverse acquisition protocols including different CT scans vendors, reconstruction kernels, and other imaging characteristics.

The database used for training the individual networks from the ensemble is composed of

Table 4.3: Classification performance using different aggregation strategies to build the ensemble

Ensemble strategy		SN(%)	SP(%)	GM(%)	BA(%)
Weighting method	RS	92.39	98.91	95.60	95.65
		(± 0.55)	(± 0.10)	(± 0.33)	(± 0.32)
	TPE	92.51	98.93	95.67	95.72
		(± 0.62)	(± 0.10)	(± 0.37)	(± 0.36)
	BO	92.47	98.93	95.64	95.70
		(± 0.58)	(± 0.09)	(± 0.34)	(± 0.33)
Non-weighting method	MVR	91.81	98.83	95.25	95.32
		(± 0.87)	(± 0.12)	(± 0.51)	(± 0.49)

SN: sensitivity; SP: specificity; GM: geometric mean;

BA: balanced accuracy.

Table 4.4: Optimized weights

Weights	Optimization Algorithm		
	RS	TPE	BO
w_1 : BCNN ^{2D}	0.468	0.682	0.745
w_2 : MSTAGE-CNN ^{2D}	0.635	0.979	1.000
w_3 : MCONTEXT-CNN ^{2D}	0.285	0.367	0.447
w_4 : BCNN ^{3D}	0.630	0.781	0.849
w_5 : MSTAGE-CNN ^{3D}	0.690	0.98	1.000
w_6 : BCNN ^{2.5D}	0.145	0.373	0.389
w_7 : MSTAGE-CNN ^{2.5D}	0.645	0.902	1.000

Figure 4.4: Evaluation of the proposed ensemble optimized with TPE when trained and tested on datasets with images reconstructed with different kernels (B35 and B50), with respect to the evaluation metrics SN (sensitivity), SP (specificity), GM (geometric mean) and BA (balanced accuracy).

different scans from diverse vendors and models, which leads to a non-homogeneous dataset, and likely improves generalizability. We sought to further explore the generalizability of our approach, specifically relating to reconstruction kernel, by taking the advantage of the fact that we were able to extract two different ROIs corresponding to two different reconstruction kernels for each point in the dataset (B35 and B50 filters).

As shown in Figure 4.4, we achieved the higher performance when training and testing were carried out on images reconstructed with the B50 filter, corresponding to the sharp reconstruction kernel. These results are in accordance with the clinical routine, where the use of sharper images are preferred when performing visual identification and quantification of interstitial diseases. Additionally, it is also shown the generalization capability of the proposed method when the networks are trained and tested on images reconstructed with different kernels.

4.3.2.3 Analysis of the ensemble performance

Figure 4.6 shows the normalized confusion matrix of the proposed ensemble optimized with TPE. As shown, the misclassifications generally occurred within interstitial group, where misclassified reticular samples were mainly confused with subpleural line and nodular patterns. Misclassifications also occurred between normal parenchyma and centrilobular emphysema. Detailed numerical outcomes of each class are presented in Table 4.5.

In addition, an evaluation of the proposed method on a subject independent test set with respect to the training set showed a sensitivity (SN) of 88.9%, specificity (SP) of 98.4%, geometric mean (GM) of 93.5% and balanced accuracy (BA) of 93.61%. The results of our

Figure 4.5: Activation maps generated by BCNN2D network. Correct label appears above the original image patch. Predicted label appears above the activation maps. Activation maps corresponding to normal parenchyma (NP) focused on the entire image patch, almost uniformly, as there is not any radiological sign of disease. Activation maps corresponding to linear scar (LINS), centrilobular emphysema (CL), subpleural line (SUBPL) and nodular pattern (NOD) focused on those areas that define each disease subtype and lesion location. For parastatal emphysema (PSE) regions the network focused on both low attenuation areas representing emphysema and pleural regions.

solution when training using each sample independently or using only samples from the same subjects are comparable. Our interpretation is that the lung damage is a process that affects the secondary lobule and is somehow independent across different secondary lobules. Therefore training and testing could be done using samples from the same subject as long as they belong to different secondary lobules as it is the case in our training set.

Understanding and interpreting the results obtained with deep learning techniques is of paramount importance. Grad-CAM technique is a valuable manner to better understand and interpret what trained deep models have learned (Selvaraju et al. 2017). We mapped the activation maps corresponding to one of our proposed two-dimensional networks (BCNN^{2D}). Activation maps highlight the regions for each sample that most influence the predictions. We observed that the activation maps consistently select the location of the lesions (Figure 4.5).

Table 4.5: Detailed performance of the proposed method

Group	Class	#Samples		Results			
		CV Train	Test	SN(%)	SP(%)	GM(%)	BA(%)
HEALTHY	Normal parenchyma (NP)	600	23096	0.8937	0.9649	0.9286	0.9293
INTERSTITIAL	Ground glass (GG)	600	34	0.9118	0.9907	0.9504	0.9512
	Reticular (RETIC)	600	4809	0.8064	0.9501	0.8753	0.8782
	Nodular (NOD)	600	30	0.9667	0.9977	0.9820	0.9822
	Linear scar (LINSCL)	600	48	0.9375	0.9936	0.9652	0.9656
	Subpleural line (SUBPL)	600	103	0.8544	0.9907	0.9200	0.9225
EMPHYSEMA	Paraseptal emphysema (PS)	600	3245	0.9405	0.9861	0.9630	0.9633
	Centrilobular emphysema (CL)	600	3013	0.9220	0.9810	0.9510	0.9515
	Avg/Total	4800	34378	0.9041	0.9818	0.9420	0.9430

SN: sensitivity; SP: specificity; GM: geometric mean; BA: balanced accuracy.

True label	NP	89.4	0.5	6.1	0.5	0.0	0.3	1.3	1.9
	GG	2.9	91.2	5.9	0.0	0.0	0.0	0.0	0.0
	RETIC	3.9	4.1	80.6	4.0	1.3	2.9	1.9	1.3
	SUBPL	0.0	0.0	11.7	85.4	1.0	1.0	1.0	0.0
	NOD	0.0	0.0	3.3	0.0	96.7	0.0	0.0	0.0
	LINSCL	0.0	0.0	4.2	0.0	2.1	93.8	0.0	0.0
	PS	1.0	0.0	1.3	0.1	0.3	0.3	94.1	3.1
	CL	5.8	0.0	0.3	0.0	0.0	0.2	1.6	92.2
		NP	GG	RETIC	SUBPL	NOD	LINSCL	PS	CL
		Predicted label							

Figure 4.6: Normalized confusion matrix of the proposed ensemble optimized with TPE for the eight tissue classes under study.

4.3.2.4 Comparison with the state-of-the-art methods

Table 4.6 compares the performance of the state-of-the-art methods to the proposed ensemble of CNNs (ECNN). The first row corresponds to a method based on local histogram hand-crafted features (LH) (Ash, Harmouche, Ross, A. A. Diaz, Hunninghake, et al. 2017), while the rest correspond to methods that use CNNs for feature extraction and classification, including the well-known GoogleNet (Szegedy et al. 2015) (57 convolutional layers) and VGG-16-Net (Simonyan et al. 2014) (13 convolutional layers) networks. Both networks were trained from scratch on our dataset, and pretrained on ImageNet and fine-tuned on our data (GoogleNet-P and VGG-P). Also shown the results of the CNN proposed by Anthimopoulos et al (Anthimopoulos et al. 2016), which focused on detecting patterns of interstitial lung

Figure 4.7: ROC analysis for the proposed ensemble and eight state-of-the-art methods. The ROC curves represent the macro-average over all classes. The area under the curve (AUC) values are also given for each ROC. VGG-P and GoogleNet-P stands for the pretrained versions on ImageNet of VGG-Net and GoogleNet architectures. VGG-LF and VGG-EF stands for the ensemble of VGG networks using late and early fusion respectively.

diseases from 2D patches, as well as those obtained with the architecture proposed by Kim GB et al (Kim et al. 2018), which considers an image patch of 20×20 pixels. We independently implemented all of the methods and trained and tested them using the same data and framework. Table 4.6 also presents the performance of all individual networks that constitute the ensemble (last seven rows).

Additionally, and for a more extended comparison, we have also implemented two different ensembles of VGG-16 networks. The ensembles are composed by nine VGG networks, each of them having as an input a single image patch including the seven axial images which comprise the 3D volume as well as the sagittal and coronal views. We used the VGG networks pretrained on ImageNet as a feature extractor. We then combine the features using both early and late fusion. For the early fusion (VGG-EF), we fused the features with a single multilayer perceptron trained on our own dataset. Conversely, for the late fusion (VGG-LF), the information is fused at the decision level, where the final fully connected layers of the nine VGG networks were fine tuned on our data independently and the predictions were subsequently fused using a soft voting rule.

In general, our proposed ensemble method, outperformed the other techniques. For example, our approach had a 28% higher BA than the method that uses hand-crafted features. In addition, compared with the rest of the methods based on CNNs, our ensemble method outperformed the best of them by 7.6% in terms of BA. We also found that very deep CNNs such as GoogleNet easily overfit the data when they are not trained in very large datasets,

yielding poor performance on the test set. The ensembles of VGG networks which process multi-view and multi-dimensional data do not outperformed the proposed method. The macro-average ROC curves for each method are shown in Figure 4.7 and demonstrated that the proposed method achieved the highest area under the curve (AUC). Finally, for comparison, in Figure 4.8, the confusion matrices of the proposed method, the individual networks that compose the ensemble and state-of-the-art are all shown.

Figure 4.8: Normalized confusion matrices of the proposed ensemble, individual networks that make up the ensemble and state-of-the-art methods.

4.3.2.5 Visual evaluation

In order to complete the validation from a clinical point of view and to extend the analysis from isolated ROIs, we performed a visual evaluation of the proposed method. For these analyses, we computed the full-lung classification of twenty CT scans from COPDGene cohort at different stages of disease severity and ILA patterns.

Full-lung classification was carried out at a fixed sampling grid with spacing $5 \times 5 \times 5$ voxels, and the rest of the voxels were classified using a nearest-neighbor interpolator. The time needed for the proposed method to process an average-sized entire scan (700 slices) using a sampling grid of $10 \times 10 \times 10$ voxels was 3,5 minutes, while the corresponding time needed when a sampling grid of $5 \times 5 \times 5$ is used was 25 minutes, using the previously described hardware.

For visual scoring, two experts inspected the classification results of twenty full lung classifications performed by two methods: the proposed ensemble of CNNs, and, for comparison,

Table 4.6: Comparison of the proposed ensemble with state-of-the-art methods. The first group of methods (first seven rows) presents the results of the state-of-the-art and the proposed ensemble methods. The second group presents the results of the individual models.

Method	SN(%)	SP(%)	GM(%)	BA(%)
LH (Ash, Harmouche, Ross, A. A. Diaz, Hunninghake, et al. 2017)	0.6888	0.9376	0.7976	0.8132
VGG	0.6241	0.9319	0.7565	0.7780
VGG-P	0.6553	0.9703	0.7727	0.8128
GoogleNet-P	0.6930	0.9264	0.7943	0.8097
GoogleNet	0.6256	0.9341	0.7608	0.7798
VGG-EF	0.7762	0.9633	0.8647	0.8697
VGG-LT	0.8148	0.9678	0.8880	0.8913
(Kim et al. 2018)	0.3759	0.9027	0.5825	0.6393
(Anthimopoulos et al. 2016)	0.7687	0.9657	0.8567	0.8672
ECNN	0.9041	0.9818	0.9420	0.9430
BCNN ^{2D}	0.8615	0.9730	0.9152	0.9172
MSTAGE-CNN ^{2D}	0.8552	0.9740	0.9125	0.9146
MCONTEXT-CNN ^{2D}	0.8621	0.9766	0.9171	0.9194
BCNN ^{2.5D}	0.8426	0.9748	0.9051	0.9087
MSTAGE-CNN ^{2.5D}	0.8776	0.9754	0.9246	0.9265
BCNN ^{3D}	0.8271	0.9650	0.8917	0.8960
MSTAGE-CNN ^{3D}	0.8335	0.9727	0.8993	0.9031

SN: sensitivity; SP: specificity; GM: geometric mean; BA: balanced accuracy

a local histogram-based method (Ash, Harmouche, Ross, A. A. Diaz, Hunninghake, et al. 2017), which has previously shown to identify interstitial patterns with both clinical and genetic associations (Ash, Harmouche, Putman, et al. 2017). The scoring was done in a blinded manner, and based on the consensus of the two reviewers as to the goodness of the method to classify each of the eight interstitial. Additionally, the performance in terms of the spatial consistency of the resulting labeled lung and the classification precision on the hilum, which is an area prone to errors due to its anatomical complexity, were assessed. Visual scoring of the results of the subtypes classification was done using a range from 0 to 10 according to the degree of under- and over-estimation of the extent of the tissue class, where 0 means significant under estimation, 5 means neither under estimation nor over estimation of the subtype, and 10 represents a high over estimation. Visual scores for spatial consistency and precision on the hilum range from 5 to 10, where 5 means high or good spatial consistency and hilar precision and 10 low spatial consistency and hilar precision.

Table 4.7 and Figure 4.9 summarize the visual scores for both methods. Table 4.7 reports the Mean Absolute Error (MAE), defined as the distance to 5, for all the inspected aspects

(subtypes and overall considerations), and the mean of overestimation (scores > 5) and underestimation (scores < 5) for all subtypes. Additionally, for reference, the Mean Percentage Volume (MPV) of each subtype under study was measured for the twenty cases. For each case the subtype percentage volume was estimated as the mean classified volume provided by both LH and ECNN methods. Although for some subtypes the local-histogram based method exhibits a better behaviour in terms of extent definition, the proposed ensemble of CNNs is significantly better than those obtained with the local histogram-based method in terms of Mean MAE for all the subtypes (two-sample t-test, p -value < 0.001). It can be also observed that for the subtypes with higher presence in our test population (defined as $MPV \geq 5\%$) the ECNN performs better than the local-histogram approach.

As can be seen in Figure 4.10, qualitative review of the technique's performance suggests that while the overall performance of the approach was excellent, there were a few systematic errors. These include the classification of linear bronchovascular bundles as linear scar (LINS), the misclassification of proximal bronchovascular bundles viewed in cross-section as nodular (NOD), and the identification of motion artifact related to the fissures and diaphragm as ground glass (GG), all of which are errors that can be easily removed via masking. Reticular (RETIC) areas were also sometimes misclassified as Nodular (NOD) pattern. This effect was previously described, and is also observed in the confusion matrix (Figure 4.6).

Figure 4.9: Visual score for twenty full-lung classification results obtained with the proposed ECNN and LH methods. *Spatial Consistency; **Hilar Precision

Figure 4.10: Automated full lung classification results of two CT scans with different ILA subtypes and stage of disease severity. The first row of the upper figure shows several ground truth areas.

4.3.2.6 Clinical validation

For further clinical validation we wanted to test if the proposed method is directly associated with the visually determined score of the presence of ILA. We hypothesized that the proposed method can determine the binary classification of a patient presenting ILA using as benchmark the visual identification of ILA.

The visual assessment of CT scans for ILA has been carried out following the previously

Table 4.7: Visual assessment of the proposed ensemble and local histogram-based method. LH stands for Local Histogram. ECNN stands for the proposed Ensemble of Convolutional Networks method.

	MPV (<i>mean±sd</i>)	MAE		Over.		Under.	
		LH	ECNN	LH	ECNN	LH	ECNN
NP	73.2±10.9	1.25	0.80	6.00	5.60	3.89	4.37
RETIC	10.0±5.6	2.45	0.80	7.45	5.77	5.00	4.75
PS	3.2±1.6	0.70	0.70	5.07	5.50	4.32	4.46
CLE	6.5±9.2	1.45	0.80	6.33	5.33	4.17	4.23
SUBPL	1.5±0.6	0.75	0.25	5.75	5.21	5.00	4.94
GG	1.2±1.0	0.40	0.50	5.40	5.07	5.00	4.52
NOD	2.5±3.8	0.00	1.15	5.00	6.15	5.00	5.00
LINSC	1.9±0.9	1.40	0.70	6.37	5.70	4.78	5.00
Sp. Cons.*	-	2.6	1.10	-	-	-	-
H. Prec.**	-	2.35	0.85	-	-	-	-
Mean	-	1.33	0.76	5.92	5.54	4.64	4.66

*Spatial Consistency; **Hilar Precision

MPV: Mean Percentage Volume, MAE: Mean Absolute Error, Over.: Overestimation, Under.: Underestimation

described procedure (Washko, Hunninghake, Fernandez, Nishino, Okajima, Yamashiro, Ross, Estepar, et al. 2011). All individual's CT scans were visually analyzed and determined to have ILA if there were nondependent ground-glass or reticular abnormalities, diffuse centrilobular nodularity, nonemphysematous cysts, honeycombing, or traction bronchiectasis affecting more than 5% of any lung zone.

We defined a quantitative ILA score as the percentage of the lung area affected by any interstitial pattern, namely ground glass, reticular, nodular, linear scar and sub pleural line where only the voxels with greater than a determined certainty threshold were considered. The certainty of a voxel was defined based on the probability given by the proposed ensemble of CNNs and associated to the predicted label for this voxel (see example in Figure 4.11a-c).

Receiver operating characteristic curves and the area under the curve were generated using logistic regression via cross-validation for the prediction of those individuals with visually identified presence of ILA.

CT scans from 114 subjects, that were not included in the training set, were evaluated (52 with visually defined ILA and 62 without ILA). Overall the proposed method had an area under the receiver operating characteristic (ROC) curve of 0.863 ± 0.067 for the detection of visually defined ILA when using a certainty threshold of 95%. For lower certainty thresholds the classifier performance slightly decreased as shown in Figure 4.11d.

Figure 4.11: **A** Original image; **B** Certainty map given by the proposed method for non-ILA labeled voxels (blue) and ILA labeled voxels (yellow); **C** Lung area affected by interstitial patterns with certainty greater than 95%; **D** ROC curves using several thresholds for the detection of visually defined ILA.

4.4 Conclusions

In this chapter, we proposed a novel method to detect and classify radiographic patterns of ILD at an early stage in CT images, considering eight radiographic classes of lung tissue, including normal tissue, five interstitial features subtypes and two emphysematous classes. Specifically, we proposed a multi-model ensemble of deep CNNs. The ensemble is comprised of seven CNNs, incorporating 2D, 2.5D and 3D networks. To this end, each individual network was trained from scratch on our database, and then the outputs of the networks are summed up in a weighted manner and combined to form the overall output of the ensemble. Both

the CNNs and the weights of the ensemble are trained via cross-validation. The resulting ensemble achieved a higher performance compared to each of the individual models, proving the potential of combining the responses of various classifiers.

The augmentation in learning enabled by utilizing different CNN architectures has been scarcely used in medical imaging but shows promising results (A. Kumar et al. 2017). Our proposal combines deeper and shallower representations with different input dimensionality. The optimized ensemble weights show that each of the single models contributes to the final classification. Although there is not a single local minima for the ensemble, the different optimization approaches produce comparable results in terms of accuracy. This might indicate some degree of redundancy in the prediction provided by each ensemble. Further studies could include the minimum set of information that is needed to achieve similar performance.

Previous studies have shown the advantage of using 2.5D and 3D architectures encoding more spatial contextual information, and therefore producing representations with higher discrimination capability (Dou et al. 2017; Roth, Lu, et al. 2016). Our study proves that the combination of different input data representations (2D, 2.5D and 3D) is important, given that the proposed ensemble improves the performance with respect to the isolated models.

Our study suggests that the proposed method can be employed to identify radiographic patterns of ILD at an early stage. In our test studies, which involve a large dataset of 34378 samples, the performance of our method showed greater area under the ROC curve (AUC) than state-of-the-art architectures commonly used by the computer vision community. Our method also outperforms two specific architectures of CNN to detect ILD patterns (Anthimopoulos et al. 2016; Kim et al. 2018), as well as a simpler approach to objectively identified ILA based on density histogram features (Ash, Harmouche, Ross, A. A. Diaz, Hunninghake, et al. 2017).

Additionally, we have shown the generalization capability of our method to be applied to large cohorts of patients with data reconstructed with different kernels. We performed intra-kernel and inter-kernel studies in which the training and test images were reconstructed with the same and different kernels respectively, and to the best of our knowledge, this is the first work that comprises images with different reconstruction algorithms and that has studied and verified the generalization capability of a method for automated classification of lung tissue.

We also performed a quantitative experiment where the method was tested on a completely independent test set separating the training and testing dataset at the patient level. This experiment allowed us to ensure the repeatability of results and generalization of the

proposed method.

To confirm the clinical feasibility of the proposed method we performed two experiments including a visual evaluation of the proposed method and a clinical validation where we tested the association between the method and the visually determined score of the presence of ILA. For the visual evaluation, two experts visually evaluated and scored by consensus twenty full-lung classification results. The results of the clinical visual assessment confirmed the statistically significant (p -value<0.001) superior performance of the proposed method against previous work based on local histogram features that has recently shown to provide insightful clinical and genetic associations (Ash, Harmouche, Putman, et al. 2017). Our visual evaluation highlights the difficulty of assessing interstitial subtypes with low prevalence. It is worth noting that the subtypes for which our method obtained lower accuracy in terms of visual scoring, are the ones that correspond to the cases with the lowest number of training and testing samples that may influence the ability of their CNN approach to learn their unique characteristics. We expect that the inclusion of more training data for these subtypes could alleviate this effect.

Regarding the ability of the method for detecting visually defined ILA, the experiment carried out on 114 subjects showed that estimation of ILA patterns in high certainty areas seems to better predict the visual assessment of ILA, suggesting the importance of the prediction probability.

In summary, we have demonstrated that the proposed ensemble of CNNs, in spite of the intrinsic difficulty of the problem due to the subtle nature of the disease, is able to identify radiographic patterns of early parenchymal lung disease that have been previously shown to be associated with prognosis and clinical outcomes. It has also been shown that our method can identify individuals with interstitial abnormalities, but further study is needed to determine the association between the response of the proposed method and other clinical outcomes including mortality.

5

Paraseptal emphysema segmentation

Emphysema subtype classification has been traditionally addressed by local-based methods which exploits the local appearance, intensity and texture of HRCT images. However, as outlined in Chapter 3, the automatic detection of paraseptal emphysema represents a unique challenge and requires global and contextual information within the lung in addition to intensity and texture information. Thus, local-based methods would not exploit the intrinsic nature of paraseptal emphysema lesions and are not able to properly characterize it. In this chapter, we present a novel method to identify paraseptal emphysema lesions exploiting contextual information and considering the specific nature of this particular subtype of emphysema.

5.1 Introduction

Pulmonary emphysema is characterized by abnormal permanent enlargement of airspaces distal to the terminal bronchiole, accompanied by the destruction of their walls, and without obvious fibrosis (Snider et al. 1985). Emphysema is divided into three major subtypes: centrilobular, panlobular and paraseptal emphysema.

Paraseptal emphysema (PSE), also known as distal acinar emphysema, is characterized by subpleural and peribronchovascular regions of low attenuation separated by intact interlobular septa (Hansell et al. 2008). PSE is a relatively unexplored emphysema subtype with a very distinct clinical manifestation that is often underrecognized clinically. However, recent studies demonstrated a significant association between paraseptal emphysema (PSE) and interstitial lung abnormalities (ILA) (Araki et al. 2015), which the latter have been associated with clinical outcomes, including mortality (Washko, D. A. Lynch, Matsuoka, Ross, Umeoka, A. Diaz, Sciurba, Hunninghake, Estepar, et al. 2010).

Densitometric analysis in computed tomography (CT) is commonly used and widely accepted to identify and quantify the extent of pulmonary emphysema. This method discriminates emphysematous and non-emphysematous tissue by thresholding the intensity levels in the CT image within the lung region.

A densitometric approach disregards the information present in the morphology of the emphysema subtypes. Thus, several methods have been previously proposed to carry out emphysema subtype classification exploiting local texture and intensity patterns (Sorensen et al. 2010b). The justification for using a local-based approach is usually founded on the pseudo-independence between adjacent regions of interest (ROI) that correspond to the secondary pulmonary lobule (smallest subunit of the lung). Although this approach is valid for detecting centrilobular and panlobular subtypes, it has been observed that is not able to resolve the identification of PSE properly. Given its particular structure, PSE is not only defined as a local pattern but it is a context-based lesion with a notion of location, always appearing adjacent to a pleural surface, as well as lobar septal walls, and which may have a considerable extension. This characteristics make the PSE identification problem consistent with a segmentation approach.

In this chapter, we present the first technique to particularly address paraseptal emphysema identification by proposing the Slice-Recovery architecture (*SR-Net*) that exploits 3D information to perform segmentation of PSE lesions on 2D slices of CT images.

5.2 Methods

3D manual annotation carries a high burden from experts, and when they make 2D annotations they usually use 3D information to place value on the spatial context around the region to segment. In this work, we propose the new Slice-Recovery network (*SR-Net*) for automatic segmentation of paraseptal emphysema (PSE) on CT images. The proposed CNN architecture, fully described in Section 5.2.1.2, leverages 3D information for 2D segmentation of PSE in CT axial images. We hypothesize that incorporating 3D context information can improve 2D segmentation results in medical imaging by capturing volumetric information around the lesion.

5.2.1 SR-Net

5.2.1.1 Volume extraction

The input of the proposed network consists on a 3D volume of size $384 \times 384 \times 8$ voxels, where the central axial slice corresponds to the target 2D slice to be segmented. This input volume is further pre-processed by segmenting and masking only the lung region.

5.2.1.2 CNN architecture

The proposed SR-Net is illustrated in Figure 5.1 and its innovative dense efficient architecture decodes 2D segmentation maps from a 3D encoded volume through a Slice-Recovery layer. The encoding pathway of the proposed CNN is composed by encoder blocks. Each encoder block consists of a convolutional dense block with an iterative concatenation of previous feature maps. Dense blocks introduced in (Huang et al. 2017) have been previously used to deal with the problem of semantic segmentation (Jégou et al. 2017). These dense blocks are subsequently processed by an ENet-style block which combines max pooling and strided convolutions to avoid representation bottlenecks (Paszke et al. 2016). It has been shown that a purely 3D version of the proposed architecture outperforms state-of-the-art segmentation CNNs such as U-Net (Ronneberger et al. 2015) when segmenting 3D pulmonary arteries in CTA images (López-Linares Román et al. 2018).

On the other hand, in the decoding pathway, transposed convolutional layers (deconvolution) and skip-connections are subsequently processed by a dense block. Skip-connections of the SR-Net are designed to pass higher resolution feature maps to the decoding path. These connections are based on a Slice-Recovery (SR) layer. We designed the SR layer to select the higher resolution feature maps that correspond to the original target 2D image to be segmented (highlighted in red in Figure 5.1). Thus, the decoding path can generate a 2D segmentation map based only on the higher resolution features of the slice under study rather than the whole input 3D volume. Deconvolution operation in the decoding path is performed

Figure 5.1: **A** SR-Net architecture which generates a 2D segmentation map given a 3D input volume. **B** Example of a dense block of 2 layers ($N=2$) (green), ENET-style block (blue) and Up-convolutional block (yellow).

to generate 2D segmentation maps from contracted features extracted by the encoding path from the 3D volume by fixing the third dimension of the 3D convolutional kernels to be 1. The decoding path ends with a $1 \times 1 \times 1$ convolution with a sigmoid activation function to generate the final probability map.

The network were trained on data augmented with random non-rigid distortions. In this work we used Adam Stochastic Optimization (with an initial learning rate $1e-4$ and plateau learning rate decay with a factor of 0.2 when the validation loss is not improved after 5 epochs) to minimize a class-weighted Dice-based loss function to consider class imbalance.

The network was implemented using Keras with TensorFlow, and built in a PC with GPU NVIDIA TITAN X Pascal 12GB, CPU Intel Core i7 3.6 GHz and 32GB of RAM.

5.2.2 Comparative methods

To prove the potential of incorporating 3D context to the network for 2D segmentation problems, we compared the proposed 3D-to-2D SR-Net to a purely 2D version of the latter (2D-Net). Additionally, we compared the proposed method to the UNet (Ronneberger et al. 2015), a strong baseline method in medical image segmentation problems, and to the converted 3D-to-2D version of the UNet following the proposed SR-Net approach (SR-UNet). We therefore prove the possibility and potential of converting previously designed architectures using the proposed SR-Net paradigm. Additionally, we will compare the proposed method to the local-based approach proposed in Chapter 3.

5.3 Experiments and results

5.3.1 Dataset and evaluation approach

An experienced radiologist manually segmented PSE lesions on 664 2D axial images coming from 111 CT scans from the COPDGene study. COPDGene centers obtained approval from their Institutional Review Boards and all subjects provided written informed consent. Of all of these subjects, 89 were randomly selected for training and validation, while the remaining 22 subjects were used for testing. Both training (55%) and validation (20%) sets comprised 498 axial slices, while the independent test set was composed by 166 images (25%). The evaluation of the proposed method as a well as the comparison to other methods were carried out on the test set. Training and validation sets were used to hyperparameter tuning of the proposed architecture.

Segmentation accuracy was evaluated using Dice Similarity Coefficient (DSC), Jaccard

Similarity Coefficient (JSC), and Mean Accuracy (MA), defined as:

$$\begin{aligned} \text{DSC} &= \frac{2 * \text{TP}}{2 * \text{TP} + \text{FP} + \text{FN}} \\ \text{JSC} &= \frac{\text{TP}}{\text{TP} + \text{FN} + \text{FP}} \\ \text{MA} &= \frac{\text{TNR} + \text{TPR}}{2} \end{aligned} \tag{5.1}$$

where TP, FP, TN and FN denote the number of true positives, false positives, true negatives and false negatives respectively. TNR and TPR denote true negative and true positive rates.

The lung area affected with PSE can vary substantially from one patient to another, ranging from 0.23 cm² up to 134 cm². Overlap ratio measures have been shown unsuitable for comparing segmentation accuracy on objects that differ in size. To compensate for this drawback, all reported metrics correspond to a weighted average of the previous metrics according to the area of the lesions, thus giving a higher weight to those lesions with a greater extension. These weighted metrics are consistent with the overall measure used to report disease extension that would be percentage of volume affected by PSE.

5.3.2 Results

The evaluation metrics are computed between the manually segmented images and the thresholded probability maps generated by the CNN. The threshold was chosen so that it maximizes the performance on the validation set.

An overview of the results are shown in Table 5.1, providing a comparison of the proposed SR-Net with a purely 2D version of the architecture as well as with UNet and a converted version of UNet using the proposed SR-Net approach. All the networks were trained with the same training samples, and tested with the same test set. Table 5.1 proves the potential of incorporating volumetric 3D context information to the network for a 2D segmentation problem. We can observe how both converted 3D-to-2D versions of the baseline architectures outperform their 2D versions by 22% and 5.1% respectively on terms of mean DSC.

Figure 5.2 shows the segmentation results of 5 cases selected from the test set with different PSE extension and in the presence of other emphysema subtypes, even with an advanced stage.

Results shown in Table 5.1 and Figure 5.2 also prove how a local-based approach fails at properly characterize PSE.

Figure 5.2: Comparison of the segmentation results for the proposed method, UNet and a the local-based approach presented in Chapter 3 for three cases from the test set with different emphysema extension.

Table 5.1: Overview of the results obtained with the proposed method and comparative methods (weighted mean $\pm$ std). Results are computed on the independent test set. DSC: Dice Similarity Coefficient, JSC: Jaccard Similarity Coefficient, MA: Mean Accuracy.

	DSC	JSC	MA
2D-UNet	0.492 ($\pm$ 0.047)	0.355 ($\pm$ 0.044)	0.713 ($\pm$ 0.020)
SR-UNet	0.712 ($\pm$ 0.013)	0.563 ($\pm$ 0.016)	0.844 ($\pm$ 0.007)
2D-Net	0.715 ($\pm$ 0.019)	0.572 ($\pm$ 0.021)	0.864 ($\pm$ 0.006)
SR-Net	0.764 ($\pm$ 0.014)	0.630 ($\pm$ 0.017)	0.908 ($\pm$ 0.004)
Local-based (Ch.3)	0.448 ($\pm$ 0.014)	0.295 ($\pm$ 0.008)	0.730 ($\pm$ 0.003)

5.4 Conclusions

In this chapter, we present a dense efficient 3D-to-2D CNN architecture (referred as SR-Net) for automatic paraseptal emphysema (PSE) segmentation in axial 2D images leveraging and exploiting contextual 3D information. Additionally, we have shown the benefit of this approach when applied to previous state-of-the-art CNN architectures.

Manual annotation of regions in 3D is a non-trivial task which demands an intensive and time-consuming labour from experts. On the other hand, when experts annotate 2D images belonging to a 3D volume, they usually use 3D information to reach optimal 2D annotations placing value on contextual information around the region to segment. Considering these premises, we proposed an approach based on 2D annotations but incorporating spatial 3D information.

It has been shown that previous local-based methods to emphysema subtype identification, achieve remarkable accuracy in characterizing and identifying centrilobular and panlobular emphysema subtypes, but are not able to properly characterize PSE. The proposed method tackles PSE identification considering the specific nature of PSE, consistent with a context-based segmentation approach.

The results, which were obtained on an independent test set, demonstrated that the proposed SR-Net outperforms state-of-the-art methods when segmenting PSE lesions on CT images.

6

Parenchymal injuries segmentation

Although local-based methods achieve remarkable accuracy in the automatic identification of parenchymal lung injuries, they still have the problem of spatial inconsistency when are extended from isolated local regions and applied for full lung classification. In this chapter we present a method to identify parenchymal lung injuries which produces a more spatially-consistent solution compared to the one obtained with a local-based method.

6.1 Introduction

Parenchymal lung injuries have been shown to affect different components of the secondary pulmonary lobule (Bergin et al. 1988). This has led to propose different methods to assess parenchymal lung injuries identification using a local-based approach by analyzing the appearance of HRCT local patches covering the secondary lobule. These local approaches have been shown to be able to identify radiographic patterns associated to both interstitial and emphysematous parenchymal lung injuries. However, it is known that lung injuries are not only defined at a local level, but performing a proper identification implies considering context and spatial information. For example, it is known that reticulation is most commonly seen in the peripheral and basal lung zones, ground glass opacities are usually limited in extent to a pulmonary lobe, honeycombing predominate on peripheral and subpleural lung regions and paraseptal emphysema always appears along peripheral pleura and fissures. Thus, consistency in terms of both neighboring tissues and the spatial position has to be considered.

Local-based approaches do not consider this context and spatial information. In general, when are applied for full lung classification, these methods label each voxel of the lung image by analyzing the appearance of local image patches centered on a given voxel. This procedure usually gives rise to a spatial inconsistency in the resulting label map.

Attempts have been made to reduce the spatial inconsistency induced by the use of local-based methods for image segmentation. These approaches are mainly focused on the post processing of the output generated by the local methods. Formulating constraints among the final label map by using conditional random fields (CRF) (Cai et al. 2016; Christ et al. 2016; Gao et al. 2016) or Markov random fields (MRF) (Shakeri et al. 2016) to refine the results can improve the final segmentation and reduce the spatial inconsistency by applying a spatial regularization to the resulting label map. Gaussian filtering of the resulting probability map has also been proposed to smooth the resulting label map and thus gaining spatial consistency (Roth, Farag, et al. 2015).

Global approaches such as segmentation methods address the spatial inconsistency problem. However, ground truth generation for segmentation methods by manually delimiting regions is a non-trivial task which demands an intensive and time-consuming labour from experts, requiring much more resources than a local labeling based on patches or points. Additionally, although parenchymal injuries are related with its spatial context, it is complicated making ground truth annotations in the form of delineated regions, as clear boundaries delimiting the lesions are hard to define, with the exception of paraseptal emphysema lesions.

We propose a novel approach to identify parenchymal lung injuries using a segmentation

method which uses a spatial global context, without the need of delimited annotations. The proposed approach is a two stage solution, where we first train a local-based method using local annotations. We then use the result of this first stage to bootstrap and train the final segmentation method which will gain spatial consistency compared to the local method.

The proposed approach may be applied to any problem where a solution with spatial consistency is needed, but segmentation ground truth is unavailable or can not be generated. This approach performs a domain transformation from a locally-defined problem to a segmentation solution.

6.2 Methods

6.2.1 Domain transformation

The proposed approach consists on the use of a segmentation architecture to produce a spatially-consistent solution for the problem of lung parenchymal injuries identification. We propose a novel approach to use a locally-defined training set to generate a new database which is consistent to a segmentation problem, and thus allowing to train a segmentation architecture to produce higher spatially-consistent solutions.

This approach is divided into two steps, where the revised output of a previously trained local method produces the ground truth for the final segmentation method. First we trained a local-based method using the original locally-defined ground truth. Once the local method is trained, we deployed the local method on a set of cases using a sliding window procedure and aggregated the results, constituting a new dataset. This generated dataset will represent the ground truth for the segmentation method (see Figure 6.1).

We hypothesize that the segmentation network is able to learn the intrinsic patterns of each parenchymal injury class as well as its spatial dependencies producing segmentation results based on a sub-optimal ground truth which is generated relying on the results obtained with a preceding method originally trained with a locally-defined dataset.

It should be noted that the proposed approach is completely agnostic to the local method used for generating the new ground truth.

6.2.1.1 Ground truth generation

As the output of the local method may contain certain level of error, it is necessary to carry out a post-processing of this output before it becomes the final ground truth of the segmentation method.

In order to show the capacity of the proposed approach to generate more consistent results regardless of the underlying local method, we will apply the domain transformation approach to two different case studies using two different local methods. The first one corresponds to a

local method based on local histogram hand-crafted features (LH) (Ash, Harmouche, Ross, A. A. Diaz, Hunninghake, et al. 2017). The second one corresponds to an ensemble strategy of various CNN which were trained on a locally-defined dataset and is fully described on Chapter 4. Both methods were trained to identify radiographic patterns of early parenchymal lung disease, including normal parenchyma (NP), incipient fibrotic pattern (ILA), centrilobular emphysema (CLE) and paraseptal emphysema (PSE).

As previously discussed on Chapter 5, local-based methods achieve remarkable accuracy in characterizing and identifying CLE subtype, but are not able to properly characterize PSE lesions. In order not to introduce this error in the generated ground truths and propagate it through the segmentation methods, for both local-based results, all voxels classified by the local method as paraseptal emphysema were eliminated. With the purpose of having PSE representation in the generated ground truths, we used the segmentation architecture presented in Chapter 5 to generate PSE annotations. It is important to note that as we introduced PSE annotations to the new ground truth, any other kind of information may be included at this point. Fig 6.1 shows a general example of the process of generating the new ground truth by processing the output of a local-based method, with the inclusion of different information which comes from new resources other than the local method. Fig 6.2 shows the particular example of the generation of the new ground truth compatible with a segmentation approach given the result of the local method based on an ensemble of CNN.

Furthermore, we reduced the error of the result provided by the ensemble of CNN during this post-processing step by only keeping those voxels in the resulting output from this local method with greater than a determined certainty threshold. The certainty of a voxel was defined based on the probability given by the ensemble of the CNN and associated to the predicted label for this voxel. Voxels with lower certainty than the determined threshold were labeled as undefined regions, as we do not have enough confidence to assign a valid label to those uncertain regions. The certainty threshold was set to be 75% reaching a balance between certainty and size of the resulting ground truth. A very high certainty threshold reduces the noise of the generated ground truth while the size (number of voxels) of the new dataset is also considerably reduced. Note that this processing could not be performed on the results generated by the LH local method since the output of this approach is not built upon probability distributions and therefore all the labeled voxels were considered.

6.2.2 SR-Net with domain transformation

The network used in this study is a multi-class version of the Slice-Recovery network (SR-Net) presented in Chapter 5 which has been demonstrated that outperforms state-of-the-art methods for segmentation purposes.

Figure 6.1: General example of the process to generate the ground truth dataset for the proposed domain transformation methodology.

Figure 6.2: Generation of the segmentation ground truth based on the result given by the ensemble of local CNN. The output is processed by incorporating PSE labels from the PSE segmentation method and eliminating low certainty labels from the local method.

The network was trained on both datasets generated by the local methods which identify normal tissue, centrilobular emphysema and interstitial lung abnormalities, as well as on PSE annotations generated by a previously trained segmentation algorithm. We used Adam Stochastic Optimization (with an initial learning rate of $1e-4$ and plateau learning rate decay with a factor of 0.2 when the validation loss is not improved after 5 epochs).

To tackle representation imbalance between classes we used a class-weighted Dice score as loss function. The loss function has also been designed to ensure that non-annotated regions (regions labeled as undefined when the dataset generated by the ensemble of CNN was used) are not considered as they are uncertain areas without a valid label. Hence, the loss for an image x with output $\hat{y}$ can be defined as:

$$L(x, \hat{y}) = \sum_{i=1}^C w_i \frac{2(\sum_j y_j \hat{y}_j)}{\sum_j y_j + \sum_j \hat{y}_j}; j \in (\text{UNDF} \neq 0) \quad (6.1)$$

where y is true label, C is the number of classes, w_i is the weight for class i which is inversely proportional to the class support, and j represents the pixels of certainty areas (i.e. pixels not labeled as undefined, UNDF).

6.3 Experiments and results

6.3.1 Training dataset

Both local methods used for the generation of the new ground truth datasets were previously trained on a locally-defined dataset comprised by 37427 manually placed training points coming from 208 CT scans acquired as part of the COPDGene study (see more details in Section

4.2.1). These points included the following parenchymal feature types: normal parenchyma, interstitial features including ground glass, reticular, nodular, linear scar and sub pleural line, and emphysematous features including centrilobular and paraseptal emphysema.

Segmentation methods were trained on 2432 axial slices from the same 208 CT scans and were each local method was deployed. Its corresponding PSE annotations were additionally incorporated for each axial slice. All interstitial lung features were aggregated so that each voxel in the lung field were labeled as normal, paraseptal emphysema, centrilobular emphysema, interstitial, or undefined.

6.3.2 Visual validation

In order to examine the benefits of the proposed approach we performed a visual evaluation of the results obtained with both segmentation methods and compared to the ones obtained with its corresponding local methods by extending the experiment carried out in Section 4.3.2.5. All methods, i.e. locally defined ensemble of CNN (L-ECNN, described on chapter 4), locally-defined LH method (L-LH)(Ash, Harmouche, Ross, A. A. Diaz, Hunninghake, et al. 2017), segmentation method trained with labels produced by ECNN method (S-ECNN) and segmentation method trained on LH labels (S-LH), were deployed on twenty CT scans completely independent from the ones used for training the methods and two experts visually inspected their results and scored the goodness of the methods to identify each of the four parenchymal subtypes: normal parenchyma (NP), interstitial subtype (ILA), paraseptal emphysema (PSE) and centrilobular emphysema (CL). Visual scoring of the results was done using a range from 0 to 10 according to the degree of under- and over-estimation of the extent of the tissue class, where 0 means significant under-estimation, 5 means neither under- nor over-estimation of the subtype, and 10 represents a high over-estimation.

Figure 6.3 and Table 6.1 summarize the visual evaluation for all methods. Table 6.1 reports the Mean Absolute Error (MAE), defined as the distance to 5 (neither under- nor over-estimation) and the mean of overestimation (scores greater than or equal to 5) and underestimation (scores less than or equal to 5) for all the inspected parenchymal subtypes. The results prove the potential of the proposed approach which is able to considerably reduce the error by regularizing the final result. It can be seen that the underestimation of normal tissue is reduced using the segmentation approach compared to the local methods. On the other hand, it can also be seen that the overestimation of ILA subtype is considerably reduced. In general terms the proposed approach is able to reduce the mean absolute error by a factor of approximately 30% when the ECNN baseline method is used, while when LH method is employed, the error is reduced by a factor of approximately 55%.

These results prove that the proposed approach can be applied to increase the perfor-

mance of any type of local method. Moreover, it can be seen that the increment in the performance is even more clear when LH method is used which takes a greater advantage of the benefit of this approach in comparison to the CNN-based method. However, it can be noted that the CNN-based method continued to show the best performance.

Table 6.1: Visual evaluation of all methods, including both segmentation methods using the proposed approach (S-ECNN, S-LH) and its corresponding local methods (L-ECNN, L-LH).

	Method	NP	ILA	CL	PSE	Mean
MAE	S-ECNN	0,50	0,85	0,70	0,35	0,60
	L-ECNN	0,80	0,90	0,60	0,95	0,81
	S-LH	0,65	1,48	0,60	0,45	0,80
	L-LH	1,25	2,26	1,45	0,70	1,42
Over.	S-ECNN	5,07	5,63	5,25	5,00	5,24
	L-ECNN	5,44	6,06	5,21	5,95	5,67
	S-LH	5,08	6,29	4,71	5,21	5,32
	L-LH	6,00	7,26	6,36	5,07	6,17
Under.	S-ECNN	4,78	5,00	4,69	4,89	4,84
	L-ECNN	4,29	4,42	4,47	5,00	4,55
	S-LH	4,61	5,00	4,71	5,00	4,83
	L-LH	3,89	-	4,17	4,32	4,13

MAE: Mean Absolute Error, *Over.*: Overestimation, *Under.*: Underestimation

Figure 6.3: Visual scoring for twenty cases comparing the results obtained with the proposed segmentation approach (S-ECNN, S-LH) and its corresponding local method (L-ECNN, L-LH).

For illustrative purposes, Figure 6.4 shows the result of two cases at different stage of disease severity and with different parenchymal injuries subtypes obtained with L-ECNN and S-ECNN.

Figure 6.4: Automatic identification and classification of different parenchymal injuries subtypes with the proposed S-ECNN and its corresponding local method (L-ECNN).

6.3.3 Clinical associations

For further validation, we wanted to compare the results between the two described methods (S-ECNN, S-LH) which use the proposed regularization approach. Our hypothesis is that the segmentation method based on an ensemble of CNN (S-ECNN) is able to identify lung parenchymal regions which are more descriptive and related to pulmonary function and clinical outcomes.

Multivariable regression models were used to evaluate the descriptiveness of parenchymal subtype specific characteristics for S-CNN and S-LH methods and its association with visual diagnosis of ILA and clinical outcomes including pulmonary function test measures such as FEV1PP and FVCP (FEV1 and FVC expressed as percentage of predicted value respectively), and exercise capacity metrics such as the six-minute walk distance test (6MWD). In particular, simple CT-based subtype specific characteristics including ILA volume percentage, mean density, standard deviation and skewness of density in ILA regions, and mean density and standard deviation of density in normal parenchyma regions were included as covariates to assess the descriptive capacity of the methods.

The analysis were performed on a population of 7497 smokers with and without spirometric COPD.

Figure 6.5 shows the receiver operating characteristic (ROC) curves for the association between the CT-based measures and the visual diagnosis of ILA using a logistic regression model. Table 6.2 summarizes the results for the association of CT-based metrics and the

Figure 6.5: ROC curves for describing the association between CT-based metrics on regions delineated by S-CNN and S-LH methods and the visual ILA diagnosis.

clinical outcomes using a linear regression model. As derived from the figure and the table, simple subtype characteristics such as ILA volume and histogram-based features for S-CNN method are able to better determine both the visual ILA diagnosis and the associations with clinical outcomes. The area under the ROC curve for S-CNN method is considerably higher than the one obtained with S-LH (experiencing an increment of 12.5%) for determining the visual ILA diagnosis. On the other hand, the total explained variance for all the clinical outcomes under study using the linear regression models are higher for the S-CNN, which also presents lower values of AIC (Akaike information criterion) for all models, when using the proposed covariates.

Therefore, these results along with the ones presented in Table 6.3 may indicate that subtype characteristics of regions detected by the S-CNN method are more explanatory and have a greater clinical value compared to S-LH, indicating that the method better delineates the radiological characteristics of the pulmonary parenchyma disease. As derived from Table 6.3 tissue heterogeneity in ILA regions appears to be an important and informative densitometric characteristic with a high association with clinical outcomes. Further research is necessary to better understand why factors contributing to the different clinical outcome differ depending on the model.

6.4 Conclusions

In this chapter, we present a novel methodology for pulmonary parenchymal injuries identification using a segmentation approach which produces a spatially consistent result. This

Table 6.2: Comparison of the association of subtype specific characteristics of regions detected by S-CNN and S-LH methods with FEV1PP, FVCP and 6MWD clinical outcomes.

	FEV1PP		FVCP		6MWD	
	R^2	AIC	R^2	AIC	R^2	AIC
S-CNN	0.331	67510	0.140	72633	0.172	109558
S-LH	0.260	68258	0.112	72907	0.144	109811

Table 6.3: Associations of CT-based subtype specific measures and clinical outcomes.

	Visual ILA diagnosis				FEV1PP			
	S-ECNN		S-LH		S-ECNN		S-LH	
	Coef.	p value	Coef.	p value	Coef.	p value	Coef.	p value
ILAvp	0,100	0,000	0,069	0,000	0,061	0,229	-0,565	0,000
HUMean_{ILA}	-0,001	0,802	0,002	0,648	0,112	0,000	0,046	0,008
HUStd_{ILA}	0,011	0,000	-0,007	0,053	0,156	0,000	0,138	0,000
HUSkewness_{ILA}	-0,553	0,002	0,190	0,108	6,777	0,000	-2,829	0,000
HUMean_{NP}	-0,010	0,001	-0,002	0,434	0,195	0,000	0,313	0,000
HUStd_{NP}	-0,002	0,604	0,005	0,069	-0,666	0,000	-0,659	0,000

	FVCP				6MWD			
	S-ECNN		S-LH		S-ECNN		S-LH	
	Coef.	p value	Coef.	p value	Coef.	p value	Coef.	p value
ILAvp	-0,084	0,018	-0,359	0,000	-0,961	0,252	-11,329	0,000
HUMean_{ILA}	0,080	0,000	0,127	0,000	2,576	0,000	3,196	0,000
HUStd_{ILA}	0,032	0,021	-0,0321	0,017	-0,454	0,152	-1,393	0,000
HUSkewness_{ILA}	5,737	0,000	-0,748	0,149	150,343	0,000	-3,316	0,776
HUMean_{NP}	-0,047	0,000	-0,064	0,000	-1,949	0,000	-1,265	0,000
HUStd_{NP}	-0,184	0,000	-0,156	0,000	-3,353	0,000	-3,709	0,000

ILAvp: ILA volume percentage

HUMean_{ILA}, HUStd_{ILA}, HUSkewness_{ILA}: mean density, standard deviation and skewness of density (HU) in ILA regions

HUMean_{NP}, HUStd_{NP}: mean density and standard deviation of density (HU) in normal parenchyma regions

approach does not need manual annotations of regions which requires an intensive labour from experts, since it is based on the result of a previously deployed local method. This chapter also shows that the proposed paradigm can be also used with weakly annotation data. It should be also noted that this method can be combined with traditional training data for segmentation algorithms and can be seen as a new manner of data augmentation.

In order to show the capacity of the proposed approach regardless of the underlying local method, we applied it to two different local methods, leading to the so-called S-CNN and S-LH methods. We compared all reference local and segmentation methods and two experts visually evaluated and scored by consensus all their results. The results proved the potential of the methodology and indicate that it is agnostic to the underlying local method used, being able to be used with any previously proposed local method for lung parenchyma characterization (Castaldi et al. 2013; Mendoza et al. 2012; Sorensen et al. 2010b).

For further evaluation, and to study the clinical feasibility of the final proposed S-CNN method which uses the proposed paradigm together with the locally defined ensemble of CNN, we studied the descriptiveness of specific subtype characteristics of regions detected by both S-CNN and S-LH methods. The results, obtained on a population of 7497 subjects, indicated that S-CNN method is able to extract imaging biomarkers which are more informative and are more related with different clinical outcomes than the ones obtained with S-LH.

7

Concluding remarks

Discussion and contributions

High resolution computed tomography (HRCT) scans plays a fundamental role in the diagnosis and assessment of pulmonary parenchyma disease in smoking-related injury, specially supporting an early diagnosis and thus affecting on subsequent management decisions.

The previous chapters describe new methods for automatic assessment and quantification of smoking-related injuries of pulmonary parenchyma, namely obstructive diseases such as emphysema and restrictive diseases including interstitial lung diseases and its early and subtle precursors findings in HRCT images.

Chapter 3 described a method for automatic emphysema identification and classification considering six different tissue classes proposing a optimally cost effective 2,5-dimensional approach, being able to reflect 3D information with similar computation cost compared to standard 2D approaches.

Based on the results obtained in Chapter 3 showing the capability of convolutional networks to automatic classify tissue classes, we hypothesized that a CNN-based method could perform an automatic and objective identification and classification of smoking-related parenchymal injuries covering a wide disease spectrum including interstitial lung abnormalities (ILA) patterns, which are subtle and early findings in HRCT which have been shown to be associated with clinical outcomes including increased mortality, as well as emphysema subtypes. In this way we profoundly extended the method, and Chapter 4 presented a robust method to detect eight radiographic classes of lung tissue, including normal tissue, five interstitial features subtypes and two emphysematous classes by a means of an ensemble of CNNs. The proposal combined deeper and shallower architectures with different input dimensionality, proving the importance and benefit of combining different input data representation (2D, 2,5D and 3D). This method showed a high generalization capability when applied to large cohorts of patients with scans reconstructed with different kernels, and demonstrated the ability to directly identify those individuals with visually identified presence of ILA.

However, the methods presented in these chapters were built based on a local-based approach. Although demonstrated to be optimal to detect most of lung tissue classes, we showed that local methods could not properly characterize the specific nature of paraseptal emphysema (PSE). Chapter 5 proposed a novel context-based segmentation method for tackling PSE identification. We proposed an innovative 3D-to-2D architecture which generates 2D annotations leveraging and exploiting 3D contextual information, without the need of having 3D labels which carries a high cognitive burden from experts and is time consuming. We additionally showed the possibility and potential of converting previously designed segmentation architectures using the paradigm proposed in this chapter.

Although Chapter 4 presented a robust method with remarkable accuracy for the identification of parenchymal injuries, it also arose that local methods still have the problem of spatial inconsistency and do not leverage the context and spatial information that characterize lung parenchymal subtypes. Chapter 6 presented a novel methodology to identify parenchymal lung injuries using a segmentation approach that uses a spatial context. The proposed paradigm does not require delimited annotation data which considerably reduces the burden from experts and opens the possibility to apply segmentation methods even when annotation training databases are not available. This chapter additionally proved that the specific subtype characteristics of those regions detected by the proposed method have a valuable source of knowledge and are related to different clinical outcomes.

Future lines should consider the application of all proposed methods for the identification of pulmonary parenchymal subtypes for the assessment of disease prognosis and progression. The proposed parenchymal subtyping methods enable the study of the evolution of lung diseases, specially ILD, by studying not only the progression of the disease extent during different time points, but also measuring the local transition between different parenchyma subtypes for the different lung regions. This probably requires a new methodology which finds the alignment of CT scans acquired at different time points, and that, alongside the proposed methods in the thesis, will be able to quantify the transitions over time of lung regions between different parenchymal subtypes. Additionally, the proposed methods enable the study of the predictive capability of disease prognosis and progression for each parenchyma subtype. In this direction, an extension of the segmentation algorithm proposed in Chapter 6 which will identify different ILA subtypes is also desirable. Finally, Chapter 5 which proposes a novel method for paraseptal emphysema segmentation opens up new possibilities for the study and understanding of this particular subtype of emphysema, which is relatively unexplored, often underrecognized clinically and that little is known about its pathology and pathogenesis.

Publications

Journal articles

Bermejo-Peláez, D., Ash, S. Y., Washko, G. R., San José Estépar, R., and Ledesma-Carbayo, M. J. (2020). “Classification of Interstitial Lung Abnormality Patterns with an Ensemble of Deep Convolutional Neural Networks”. In: *Scientific Reports* 10.1, pages 1–15. ISSN: 2045-2322. DOI: 10.1038/s41598-019-56989-5.

Jimenez-Carretero, D., Bermejo-Peláez, D., Nardelli, P., Fraga, P., Fraile, E., San José Estépar, R., and Ledesma-Carbayo, M. J. (2019). “A graph-cut approach for pulmonary artery-vein segmentation in noncontrast CT images”. In: *Medical Image Analysis* 52, pages 144–159. ISSN: 1361-8423. DOI: 10.1016/j.media.2018.11.011.

Nardelli, P., Jimenez-Carretero, D., Bermejo-Pelaez, D., Washko, G. R., Rahaghi, F. N., Ledesma-Carbayo, M. J., and San José Estépar, R. (2018). “Pulmonary Artery-Vein Classification in CT Images Using Deep Learning”. In: *IEEE Transactions on Medical Imaging* 37.11, pages 2428–2440. ISSN: 1558-254X. DOI: 10.1109/TMI.2018.2833385.

Conference articles

Bermejo-Peláez, D., Okajima, Y., Washko, G. R., Ledesma-Carbayo, M. J., and Estépar, R. S. J. (2019). “A SR-NET 3D-to-2D Architecture For Paraseptal Emphysema Segmentation”. In: *2019 IEEE 16th International Symposium on Biomedical Imaging (ISBI 2019)*, pages 303–306. DOI: 10.1109/ISBI.2019.8759184.

Bermejo-Peláez, D., Estepar, R. S. J., and Ledesma-Carbayo, M. J. (2018). “Emphysema classification using a multi-view convolutional network”. In: *2018 IEEE 15th International Symposium on Biomedical Imaging (ISBI 2018)*, pages 519–522. DOI: 10.1109/ISBI.2018.8363629.

- Gimenez Ucendo, M., Bermejo-Peláez, D., Ortuño Fisac, J. E., McVeigh, E. R., and Ledesma-Carbayo, M. J. (2018). "Desarrollo de una herramienta para la detección automática del plano valvular mitral mediante algoritmos de deep learning". In: *Actas XXXVI Congreso Anual de la Sociedad Española de Ingeniería Biomédica*, pages 33–36.
- Bermejo-Peláez, D., Estepar, R. S. J., and Ledesma-Carbayo, M. J. (2017). "Emphysema detection and classification using a multi-scale deep convolutional network". In: *CARS 2017-Computer Assisted Radiology and Surgery Proceedings of the 31st International Congress and Exhibition*. Volume 12, pages 141–143. DOI: 10.1007/s11548-017-1588-3.
- Nardelli, P., Jimenez-Carretero, D., Bermejo-Peláez, D., Ledesma-Carbayo, M. J., Rahaghi, F. N., and Estépar, R. S. J. (2017). "Deep-learning strategy for pulmonary artery-vein classification of non-contrast CT images". In: *2017 IEEE 14th International Symposium on Biomedical Imaging (ISBI 2017)*, pages 384–387. DOI: 10.1109/ISBI.2017.7950543.
- Bermejo-Peláez, D., Estepar, R. S. J., and Ledesma-Carbayo, M. J. (2016a). "Detección y clasificación de enfisema pulmonar en imágenes de TAC mediante Redes Neuronales Convolucionales Multiescala". In: *Actas XXXIV Congreso Anual de la Sociedad Española de Ingeniería Biomédica*, pages 370–373.
- Bermejo-Peláez, D., Estepar, R. S. J., and Ledesma-Carbayo, M. J. (2016b). "Publicly available tool for emphysema subtyping quantification in computed tomography images". In:
- Bermejo-Peláez, D., Estepar, R. S. J., España-Palomares, S., Peces-Barba, G., Cosío, B. G., Agustí, A., Ruiz-Cabello, J., and Ledesma-Carbayo, M. J. (2015). "Cuantificación de enfisema pulmonar a partir de imágenes de TAC para la determinación de calcio coronario". In: *Actas XXXIII Congreso Anual de la Sociedad Española de Ingeniería Biomédica*, pages 285–288.

Bibliography

- Anthimopoulos, M., Christodoulidis, S., Ebner, L., Christe, A., and Mougiakakou, S. (2016). "Lung Pattern Classification for Interstitial Lung Diseases Using a Deep Convolutional Neural Network". In: *IEEE Trans Med Imaging* 35.5, pages 1207–1216. DOI: 10.1109/TMI.2016.2535865.
- Araki, T., Nishino, M., Zazueta, O. E., Gao, W., Dupuis, J., Okajima, Y., Latourelle, J. C., Rosas, I. O., Murakami, T., O'Connor, G. T., Washko, G. R., Hunninghake, G. M., and Hatabu, H. (July 2015). "Paraseptal emphysema: Prevalence and distribution on CT and association with interstitial lung abnormalities". In: *European Journal of Radiology* 84.7, pages 1413–1418. DOI: 10.1016/j.ejrad.2015.03.010.
- Ash, S. Y., Harmouche, R., Vallejo, D. L. L., Villalba, J. A., Ostridge, K., Gunville, R., Come, C. E., Onieva, J. O., Ross, J. C., Hunninghake, G. M., El-Chemaly, S. Y., Doyle, T. J., Nardelli, P., Sanchez-Ferrero, G. V., Goldberg, H. J., Rosas, I. O., Estepar, R. S. J., and Washko, G. R. (2017). "Densitometric and local histogram based analysis of computed tomography images in patients with idiopathic pulmonary fibrosis". In: *Respir Res* 18.1. DOI: 10.1186/s12931-017-0527-8.
- Ash, S. Y., Harmouche, R., Ross, J. C., Diaz, A. A., Hunninghake, G. M., Putman, R. K., Onieva, J., Martinez, F. J., Choi, A. M., Lynch, D. A., Hatabu, H., Rosas, I. O., Estepar, R. S. J., and Washko, G. R. (2017). "The Objective Identification and Quantification of Interstitial Lung Abnormalities in Smokers". In: *Acad Radiol* 24.8, pages 941–946.
- Ash, S. Y., Harmouche, R., Putman, R. K., Ross, J. C., Diaz, A. A., Hunninghake, G. M., Onieva, J. O., Martinez, F. J., Choi, A. M., Lynch, D. A., Hatabu, H., Rosas, I. O., Estepar, R. S. J., and Washko, G. R. (2017). "Clinical and Genetic Associations of Objectively Identified Interstitial Changes in Smokers". In: *Chest*.

- Ash, S. Y., Harmouche, R., Ross, J. C., Diaz, A. A., Rahaghi, F. N., Sanchez-Ferrero, G. V., Putman, R. K., Hunninghake, G. M., Onieva, J. O., Martinez, F. J., Choi, A. M., Bowler, R. P., Lynch, D. A., Hatabu, H., Bhatt, S. P., Dransfield, M. T., Wells, J. M., Rosas, I. O., Estepar, R. S. J., Washko, G. R., and Investigators, C. (Aug. 2018). "Interstitial Features at Chest CT Enhance the Deleterious Effects of Emphysema in the COPD Gene Cohort". In: *Radiology* 288.2, pages 600–609. DOI: 10.1148/radiol.2018172688.
- Auerbach, O., Garfinke, L., and Hammond, E. C. (1974 1974). "Relation of Smoking and Age to Findings in Lung Parenchyma - Microscopic Study". In: *Chest* 65.1, pages 29–35. DOI: 10.1378/chest.65.1.29.
- Bengio, Y., Boulanger-Lewandowski, N., and Pascanu, R. (2013). "Advances in Optimizing Recurrent Networks". In: *IEEE ICASSP*, pages 8624–8628.
- Berend, N., Rynell, A., and Ward, H. (Feb. 1991). "Structure of a Human Pulmonary Acinus". In: *Thorax* 46.2, pages 117–121.
- Bergin, C., Roggli, V., Coblentz, C., and Chiles, C. (July 1988). "The Secondary Pulmonary Lobule - Normal and Abnormal CT Appearances". In: *American Journal of Roentgenology* 151.1, pages 21–25. DOI: 10.2214/ajr.151.1.21.
- Bergstra, J., Bardenet, R., Bengio, Y., and Kégl, B. (2011). "Algorithms for hyper-parameter optimization". In: *Adv Neural Inf Process Syst*, pages 2546–2554.
- Best, A. C., Lynch, A. M., Bozic, C. M., Miller, D., Grunwald, G. K., and Lynch, D. A. (Aug. 2003). "Quantitative CT indexes in idiopathic pulmonary fibrosis: Relationship with physiologic impairment". In: *Radiology* 228.2, pages 407–414. DOI: 10.1148/radiol.2282020274.
- Bodlet, A., Maury, G., Jamart, J., and Dahlqvist, C. (Nov. 2013). "Influence of radiological emphysema on lung function test in idiopathic pulmonary fibrosis". In: *Respiratory medicine* 107.11, pages 1781–1788. DOI: 10.1016/j.rmed.2013.08.039.
- Borgerding, M. and Klus, H. (July 2005). "Analysis of complex mixtures - Cigarette smoke". In: *Experimental and Toxicologic Pathology* 57, pages 43–73.
- Cai, J., Lu, L., Zhang, Z., Xing, F., Yang, L., and Yin, Q. (2016-Oct 2016). "Pancreas Segmentation in MRI using Graph-Based Decision Fusion on Convolutional Neural Networks." In: *Medical image computing and computer-assisted intervention : MICCAI ...Internationa-*

- tional Conference on Medical Image Computing and Computer-Assisted Intervention* 9901, pages 442–450. DOI: 10.1007/978-3-319-46723-8_51.
- Camiciottoli, G., Diciotti, S., Bartolucci, M., Orlandi, I., Bigazzi, F., Matucci-Cerinic, M., Pistolesi, M., and Mascalchi, M. (2013). “Whole-Lung Volume and Density in Spirometrically-Gated Inspiratory and Expiratory CT in Systemic Sclerosis: Correlation with Static Volumes at Pulmonary Function Tests”. In: *Sarcoidosis Vasc Diffuse Lung Dis* 30.1, pages 17–27.
- Castaldi, P. J., Estepar, R. S. J., Mendoza, C. S., Hersh, C. P., Laird, N., Crapo, J. D., Lynch, D. A., Silverman, E. K., and Washko, G. R. (2013). “Distinct Quantitative Computed Tomography Emphysema Patterns Are Associated with Physiology and Function in Smokers”. In: *Am J Resp and Crit Care* 188.9, pages 1083–1090.
- Cavigli, E., Camiciottoli, G., Diciotti, S., Orlandi, I., Spinelli, C., Meoni, E., Grassi, L., Farfalla, C., Pistolesi, M., Falaschi, F., and Mascalchi, M. (2009). “Whole-lung densitometry versus visual assessment of emphysema”. In: *Eur Radiol* 19.7, pages 1686–1692.
- Chabat, F., Yang, G., and Hansell, D. (Sept. 2003). “Obstructive lung diseases: Texture classification for differentiation at CT”. In: *Radiology* 228.3, pages 871–877. DOI: 10.1148/radiol.2283020505.
- Christ, P. F., Elshaer, M. E. A., Ettlinger, F., Tatavarty, S., Bickel, M., Bilic, P., Rempfler, M., Armbruster, M., Hofmann, F., D’Anastasi, M., Sommer, W. H., Ahmadi, S. .-A., and Menze, B. H. (2016 2016). “Automatic liver and lesion segmentation in CT using cascaded fully convolutional neural networks and 3D conditional random fields”. In: *Medical Image Computing and Computer-Assisted Intervention - MICCAI 2016*, page 23. DOI: 10.1007/978-3-319-46723-8_48.
- Christodoulidis, S., Anthimopoulos, M., Ebner, L., Christe, A., and Mougiakakou, S. (2017). “Multisource Transfer Learning with Convolutional Neural Networks for Lung Pattern Analysis”. In: *IEEE J Biomed Health* 21.1, pages 76–84.
- Cottin, V., Nunes, H., Brillet, P. Y., Delaval, P., Devouassoux, G., Tillie-Leblond, I., Israel-Biet, D., Court-Fortune, I., Valeyre, D., Cordier, J. F., and GERMOP (Oct. 2005). “Combined pulmonary fibrosis and emphysema: a distinct underrecognised entity”. In: *European Respiratory Journal* 26.4, pages 586–593. DOI: 10.1183/09031936.05.00021005.

- Cottin, V., Nunes, H., Mouthon, L., Gamondes, D., Lazor, R., Hachulla, E., Revel, D., Valeyre, D., Cordier, J.-F., and Orphelines, G. E. R. M. (Jan. 2011). "Combined Pulmonary Fibrosis and Emphysema Syndrome in Connective Tissue Disease". In: *Arthritis and Rheumatism* 63.1, pages 295–304. DOI: 10.1002/art.30077.
- Delorme, S., KellerReichenbecher, M. A., Zuna, I., Schlegel, W., and vanKaick, G. (Sept. 1997). "Usual interstitial pneumonia - Quantitative assessment of high-resolution computed tomography findings by computer-assisted texture-based image analysis". In: *Investigative radiology* 32.9, pages 566–574. DOI: 10.1097/00004424-199709000-00009.
- Depeursinge, A., Sage, D., Hidki, A., Platon, A., Poletti, P.-A., Unser, M., and Mueller, H. (2007). "Lung tissue classification using wavelet frames". In: *IEEE EMBC*.
- Dirksen, A., Friis, M., Olesen, K. P., Skovgaard, L. T., and Sorensen, K. (Sept. 1997). "Progress of emphysema in severe alpha(1)-antitrypsin deficiency as assessed by annual CT". In: *Acta radiologica* 38.5, pages 826–832.
- Dou, Q., Chen, H., Yu, L., Qin, J., and Heng, P.-A. (2017). "Multilevel Contextual 3-D CNNs for False Positive Reduction in Pulmonary Nodule Detection". In: *IEEE Trans Biomed Eng* 64.7, pages 1558–1567.
- Doyle, T. J., Hunninghake, G. M., and Rosas, I. O. (2012). "Subclinical interstitial lung disease: Why you should care". In: *Am J Respir Crit Care Med* 185.11, pages 1147–1153. DOI: 10.1164/rccm.201108-1420PP.
- Farahani, F. V., Ahmadi, A., and Zarandi, M. H. F. (2015). "Lung nodule diagnosis from CT images based on ensemble learning". In: *2015 IEEE Conf Compu Inte Bioinform Comp Bio (CIBCB)*, pages 1–7.
- Fowles, J. and Dybing, E. (Dec. 2003). "Application of toxicological risk assessment principles to the chemical constituents of cigarette smoke". In: *Tobacco control* 12.4, pages 424–430.
- Gangeh, M. J., Sørensen, L., Shaker, S. B., Kamel, M. S., Bruijne, M. D., and Loog, M. (2010 2010). "A Texton-Based Approach for the Classification of Lung Parenchyma in CT Images". In: *Lect Notes Comput Sc - MICCAI 2010* 6363, pages 595–602.
- Gao, M., Bagci, U., Lu, L., Wu, A., Buty, M., Shin, H.-C., Roth, H., Papadakis, G. Z., Depeursinge, A., Summers, R. M., Xu, Z., and Mollura, D. J. (2018). "Holistic classification of CT atten-

- uation patterns for interstitial lung diseases via deep convolutional neural networks". In: *Comput Methods Biomech Biomed Eng Imaging Vis* 6.1, pages 1–6. DOI: 10.1080/21681163.2015.1124249.
- Gao, M., Xu, Z., Lu, L., Wu, A., Nogues, I., Summers, R. M., and Mollura, D. J. (2016 2016). "Segmentation Label Propagation using Deep Convolutional Neural Networks and Dense Conditional Random Field". In: *2016 IEEE 13th International Symposium on Biomedical Imaging (ISBI)*, pages 1265–1268. DOI: 10.1109/ISBI.2016.7493497.
- Gerard, S. E., Patton, T. J., Christensen, G. E., Bayouth, J. E., and Reinhardt, J. M. (Jan. 2019). "FissureNet: A Deep Learning Approach For Pulmonary Fissure Detection in CT Images". In: *IEEE Trans Med Imaging* 38.1, pages 156–166. DOI: 10.1109/TMI.2018.2858202.
- Gevenois, P. A., DeVuyst, P., deMaertelaer, V., Zanen, J., Jacobvitz, D., Cosio, M. G., and Yernault, J. C. (July 1996). "Comparison of computed density and microscopic morphometry in pulmonary emphysema". In: *American Journal of Respiratory and Critical Care Medicine* 154.1, pages 187–192. DOI: 10.1164/ajrccm.154.1.8680679.
- Gonzalez, G., Ash, S. Y., Vegas-Sanchez-Ferrero, G., Onieva, J. O., Rahaghi, F. N., Ross, J. C., Diaz, A., Estepar, R. S. J., Washko, G. R., COPDGene, and Investigator, E. (Jan. 2018). "Disease Staging and Prognosis in Smokers Using Deep Learning in Chest Computed Tomography". In: *Am J Respir Crit Care Med* 197.2, pages 193–203. DOI: 10.1164/rccm.201705-08600C.
- Gonzalez, G., Washko, G. R., and Estepar, R. S. J. (2018 2018). "Deep Learning for Biomarker Regression. Application to Osteoporosis and Emphysema on Chest CT Scans". In: *Proc SPIE Int Soc Opt Eng* 10574, UNSP 105741H. DOI: 10.1117/12.2293455.
- Hansell, D. M., Bankier, A. A., MacMahon, H., McLoud, T. C., Mueller, N. L., and Remy, J. (2008). "Fleischner Society: Glossary of terms for thoracic imaging". In: *Radiology* 246.3, pages 697–722.
- Hartley, P. G., Galvin, J. R., Hunninghake, G. W., Merchant, J. A., Yagla, S. J., Speakman, S. B., and Schwartz, D. A. (Jan. 1994). "High-Resolution Ct-Derived Measures of Lung Density are Valid Indexes of Interstitial Lung-Disease". In: *Journal of applied physiology* 76.1, pages 271–277.
- Hatt, C., Galban, C., Labaki, W., Kazerooni, E., Lynch, D., and Han, M. (2018 2018). "Convolutional Neural Network Based COPD and Emphysema Classifications Are Predictive of

- Lung Cancer Diagnosis". In: *Image Analysis for Moving Organ, Breast, and Thoracic Images* 11040, pages 302–309. DOI: 10.1007/978-3-030-00946-5_30.
- Heitzman, E. R., Markarian, B., Berger, I., and Dailey, E. (1969a). "Secondary Pulmonary Lobule - A Practical Concept for Interpretation of Chest Radiographs. I. Roentgen Anatomy of Normal Secondary Pulmonary Lobule". In: *Radiology* 93.3, pages 507–+.
- (1969b). "Secondary Pulmonary Lobule - A Practical Concept for Interpretation of Chest Radiographs. II. Application of the Anatomic Concept to an Understanding of Roentgen Pattern in Disease States". In: *Radiology* 93.3, pages 513–+.
- Heussel, C. P., Herth, F. J. F., Kappes, J., Hantusch, R., Hartlieb, S., Weinheimer, O., Kauczor, H. U., and Eberhardt, R. (Oct. 2009). "Fully automatic quantitative assessment of emphysema in computed tomography: comparison with pulmonary function testing and normal values". In: *European radiology* 19.10, pages 2391–2402. DOI: 10.1007/s00330-009-1437-z.
- Higham, A., Quinn, A. M., Cancado, J. E. D., and Singh, D. (Mar. 2019). "The pathology of small airways disease in COPD: historical aspects and future directions". In: *Respiratory Research* 20, page 49. DOI: 10.1186/s12931-019-1017-y.
- Houghton, A. M., Mouded, M., and Shapiro, S. D. (Oct. 2008). "Common origins of lung cancer and COPD". In: *Nature medicine* 14.10, pages 1023–1024. DOI: 10.1038/nm1008-1023.
- Huang, G., Liu, Z., Maaten, L. v. d., and Weinberger, K. Q. (2017). "Densely Connected Convolutional Networks". In: *2017 IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, pages 2261–2269. DOI: 10.1109/CVPR.2017.243.
- Humphries, S. M., Notary, A. M., Centeno, J. P., and Lynch, D. A. (2018). "Automatic Classification of Centrilobular Emphysema on CT Using Deep Learning: Comparison with Visual Scoring". In: *Image Analysis for Moving Organ, Breast, and Thoracic Images* 11040, pages 319–325. DOI: 10.1007/978-3-030-00946-5_32.
- Humphries, S. M., Yagihashi, K., Huckleberry, J., Rho, B.-H., Schroeder, J. D., Strand, M., Schwarz, M. I., Flaherty, K. R., Kazerooni, E. A., Beek, E. J. R. van, and Lynch, D. A. (Oct. 2017). "Idiopathic Pulmonary Fibrosis: Data-driven Textural Analysis of Extent of Fibrosis at Baseline and 15-Month Follow-up". In: *Radiology* 285.1, pages 270–278. DOI: 10.1148/radiol.2017161177.

- Hunninghake, G. M., Hatabu, H., Okajima, Y., Gao, W., Dupuis, J., Latourelle, J. C., Nishino, M., Araki, T., Zazueta, O. E., Kurugol, S., Ross, J. C., Estepar, R. S. J., Murphy, E., Steele, M. P., Loyd, J. E., Schwarz, M. I., Fingerlin, T. E., Rosas, I. O., Washko, G. R., O'Connor, G. T., and Schwartz, D. A. (June 2013). "MUC5B Promoter Polymorphism and Interstitial Lung Abnormalities". In: *N Engl J Med* 368.23, pages 2192–2200. DOI: 10.1056/NEJMoa1216076.
- Inomata, M., Ikushima, S., Awano, N., Kondoh, K., Satake, K., Masuo, M., Kusunoki, Y., Moriya, A., Kamiya, H., Ando, T., Yanagawa, N., Kumasaka, T., Ogura, T., Sakai, F., Azuma, A., Gemma, A., and Takemura, T. (2014). "An autopsy study of combined pulmonary fibrosis and emphysema: correlations among clinical, radiological, and pathological features". In: *BMC Pulmonary Medicine* 14, page 104.
- Jacobs, C., Sanchez, C. I., Saur, S. C., Twellmann, T., Jong, P. A. de, and Ginneken, B. van (2011). "Computer-Aided Detection of Ground Glass Nodules in Thoracic CT Images Using Shape, Intensity and Context Features". In: *Medical Image Computing and Computer-Assisted Intervention, Miccai 2011, Pt Iii* 6893, pages 207–+.
- Jégou, S., Drozdal, M., Vázquez, D., Romero, A., and Bengio, Y. (2017). "The One Hundred Layers Tiramisu: Fully Convolutional DenseNets for Semantic Segmentation". In: *2017 IEEE Conference on Computer Vision and Pattern Recognition Workshops (CVPRW)*, pages 1175–1183.
- Karabulut, E. M. and Ibrikci, T. (2015). *Emphysema Discrimination from Raw HRCT Images by Convolutional Neural Networks*, page 708.
- Kawata, Y., Kageyama, K., Niki, N., Umetani, K., Yada, K., Ohmatsu, H., Moriyama, N., and Itoh, H. (2010). "Microstructural analysis of secondary pulmonary lobule imaged by synchrotron radiation micro CT using offset scan mode". In: *Medical Imaging 2010: Biomedical Applications in Molecular, Structural, and Functional Imaging* 7626, UNSP 762610.
- Kim, G. B., Jung, K.-H., Lee, Y., Kim, H.-J., Kim, N., Jun, S., Seo, J. B., and Lynch, D. A. (Aug. 2018). "Comparison of Shallow and Deep Learning Methods on Classifying the Regional Pattern of Diffuse Lung Disease". In: *J Digit Imaging* 31.4, pages 415–424. DOI: 10.1007/s10278-017-0028-9.
- Kliment, C. R., Araki, T., Doyle, T. J., Gao, W., Dupuis, J., Latourelle, J. C., Zazueta, O. E., Fernandez, I. E., Nishino, M., Okajima, Y., Ross, J. C., Estepar, R. S. J., Diaz, A. A., Lederer,

- D. J., Schwartz, D. A., Silverman, E. K., Rosas, I. O., Washko, G. R., O'Connor, G. T., Hatabu, H., and Hunninghake, G. M. (Oct. 2015). "A comparison of visual and quantitative methods to identify interstitial lung abnormalities". In: *Bmc Pulmonary Medicine* 15, page 134. DOI: 10.1186/s12890-015-0124-x.
- Kumagai, S., Marumo, S., Yamanashi, K., Tokuno, J., Ueda, Y., Shoji, T., Nishimura, T., Huang, C.-I., and Fukui, M. (Dec. 2014). "Prognostic significance of combined pulmonary fibrosis and emphysema in patients with resected non-small-cell lung cancer: a retrospective cohort study". In: *European Journal of Cardio-Thoracic Surgery* 46.6, E113–E119. DOI: 10.1093/ejcts/ezu384.
- Kumar, A., Kim, J., Lyndon, D., Fulham, M., and Feng, D. (2017). "An Ensemble of Fine-Tuned Convolutional Neural Networks for Medical Image Classification". In: *IEEE Trans Biomed Eng* 21.1, pages 31–40.
- Kumar, Cherian, Vassallo, Yi, and Ryu (Aug. 2018). "Current Concepts in Pathogenesis, Diagnosis, and Management of Smoking-Related Interstitial Lung Diseases". In: *Chest* 154.2, pages 394–408. DOI: 10.1016/j.chest.2017.11.023.
- Kurashima, K., Takayanagi, N., Tsuchiya, N., Kanauchi, T., Ueda, M., Hoshi, T., Miyahara, Y., and Sugita, Y. (July 2010). "The effect of emphysema on lung function and survival in patients with idiopathic pulmonary fibrosis". In: *Respirology* 15.5, pages 843–848. DOI: 10.1111/j.1440-1843.2010.01778.x.
- Kwak, N., Park, C.-M., Lee, J., Park, Y. S., Lee, S.-M., Yim, J.-J., Yoo, C.-G., Kim, Y. W., Han, S. K., and Lee, C.-H. (Mar. 2014). "Lung cancer risk among patients with combined pulmonary fibrosis and emphysema". In: *Respiratory medicine* 108.3, pages 524–530. DOI: 10.1016/j.rmed.2013.11.013.
- Lederer, D. J., Enright, P. L., Kawut, S. M., Hoffman, E. A., Hunninghake, G., Beek, E. J. R. V., Austin, J. H. M., Jiang, R., Lovasi, G. S., and Barr, R. G. (2009). "Cigarette smoking is associated with subclinical parenchymal lung disease: The Multi-Ethnic Study of Atherosclerosis MESA-lung study". In: *Am J Respir Crit Care Med* 180.5, pages 407–414. DOI: 10.1164/rccm.200812-19660C.

- Ley, B., Collard, H. R., and Jr., T. E. K. (Feb. 2011). "Clinical Course and Prediction of Survival in Idiopathic Pulmonary Fibrosis". In: *Am J Respir Crit Care Med* 183.4, pages 431–440. DOI: 10.1164/rccm.201006-0894CI.
- Li, Q., Cai, W., and Feng, D. D. (2013). "Lung image patch classification with automatic feature learning". In: *Conf Proc IEEE Eng Med Biol Soc*, pages 6079–6082. DOI: 10.1109/EMBC.2013.6610939.
- Lim, J., Kim, N., Seo, J. B., Lee, Y. K., Lee, Y., and Kang, S. .-.-H. (2011). "Regional context-sensitive support vector machine classifier to improve automated identification of regional patterns of diffuse interstitial lung disease". In: *J Digit Imaging* 24.6, pages 1133–1140. DOI: 10.1007/s10278-011-9367-0.
- Liu, C.-T., Tai, P.-L., Chen, A. Y. .-.-, Peng, C.-H., Lee, T., and Wang, J.-S. (2001). "A content-based CT lung image retrieval system for assisting differential diagnosis images collection". In: *IEEE International Conference on Multimedia and Expo, 2001. ICME 2001*. Pages 174–177. DOI: 10.1109/ICME.2001.1237684.
- López-Linares Román, K., La Bruere, I. de, Onieva, J., Andresen, L., Qvortrup Holsting, J., Rahaghi, F. N., Macía, I., González Ballester, M. A., and San José Estepar, R. (2018). "3D Pulmonary Artery Segmentation from CTA Scans Using Deep Learning with Realistic Data Augmentation". In: *Image Analysis for Moving Organ, Breast, and Thoracic Images*. Springer International Publishing, pages 225–237.
- Lynch, D., Godwin, J., Safrin, S., Starko, K., Hormel, P., Brown, K., Raghu, G., King, T., Bradford, W., Schwartz, D., Webb, W., and Stud, I. P. F. (Aug. 2005). "High-resolution computed tomography in idiopathic pulmonary fibrosis: Diagnosis and prognosis". In: *American Journal of Respiratory and Critical Care Medicine* 172.4, pages 488–493. DOI: 10.1164/rccm.200412-17560C.
- Madan, R., Matalon, S., and Vivero, M. (Mar. 2016). "Spectrum of Smoking-related Lung Diseases Imaging Review and Update". In: *Journal of thoracic imaging* 31.2, pages 78–91. DOI: 10.1097/RTI.000000000000185.
- Madani, A., Zanen, J., Maertelaer, V. de, and Gevenois, P. A. (Mar. 2006). "Pulmonary emphysema: Objective quantification at mufti-detector row CT - Comparison with macroscopic

- and microscopic morphometry". In: *Radiology* 238.3, pages 1036–1043. DOI: 10.1148/radiol.2382042196.
- Mai, C., Verleden, S. E., McDonough, J. E., Willems, S., De Wever, W., Coolen, J., Dubbeldam, A., Van Raemdonck, D. E., Verbeken, E. K., Verleden, G. M., Hogg, J. C., Vanaudenaerde, B. M., Wuyts, W. A., and Verschakelen, J. A. (2017). "Thin-Section CT Features of Idiopathic Pulmonary Fibrosis Correlated with Micro-CT and Histologic Analysis". In: *Radiology* 283.1, pages 252–263. DOI: 10.1148/radiol.2016152362.
- Malone, J., Rossiter, J. M., Prabhu, S., and Goddard, P. (2004). "Identification of disease in CT of the lung using texture-based image analysis". In: *Conference Record of the Thirty-Eighth Asilomar Conference on Signals, Systems and Computers, 2004.* 2, 1620–1624 Vol.2. DOI: 10.1109/ACSSC.2004.1399431.
- Mascalchi, M., Diciotti, S., Sverzellati, N., Camiciottoli, G., Ciccotosto, C., Falaschi, F., and Zompatori, M. (Feb. 2012). "Low agreement of visual rating for detailed quantification of pulmonary emphysema in whole-lung CT". In: *Acta radiologica* 53.1, pages 53–60. DOI: 10.1258/ar.2011.110419.
- Matsuoka, S., Yamashiro, T., Matsushita, S., Kotoku, A., Fujikawa, A., Yagihashi, K., and Nakajima, Y. (2015a). "Quantitative CT evaluation in patients with combined pulmonary fibrosis and emphysema: Correlation with pulmonary function". In: *Acad Radiol* 22.5, pages 626–631. DOI: 10.1016/j.acra.2015.01.008.
- Matsuoka, S., Yamashiro, T., Matsushita, S., Kotoku, A., Fujikawa, A., Yagihashi, K., and Nakajima, Y. (May 2015b). "Quantitative CT Evaluation in Patients with Combined Pulmonary Fibrosis and Emphysema: Correlation with Pulmonary Function". In: *Academic Radiology* 22.5, pages 626–631.
- Mejia, M., Carrillo, G., Rojas-Serrano, J., Estrada, A., Suarez, T., Alonso, D., Barrientos, E., Gaxiola, M., Navarro, C., and Selman, M. (July 2009). "Idiopathic Pulmonary Fibrosis and Emphysema Decreased Survival Associated With Severe Pulmonary Arterial Hypertension". In: *Chest* 136.1, pages 10–15. DOI: 10.1378/chest.08-2306.
- Mendoza, C. S., Washko, G. R., Ross, J. C., Diaz, A. A., Lynch, D. A., Crapo, J. D., Silverman, E. K., Acha, B., Serrano, C., and Estepar, R. S. J. (2012). "Emphysema Quantification in a

- Multi-Scanner HRCT Cohort using Local Intensity Distributions". In: *IEEE ISBI*, pages 474–477.
- Mitani, Y., Hirayama, H., Yasuda, H., Kido, S., Hamamoto, Y., Ueda, K., and Matsunaga, N. (2000 2000). *A Gabor filter-based classification for diffuse lung opacities in thin-section computed tomography images*, page 783. ISBN: 0-7803-6400-7.
- Muller, N. L., Staples, C. A., Miller, R. R., and Abboud, R. T. (Oct. 1988). "Density Mask - an Objective Method to Quantitate Emphysema using Computed-Tomography". In: *Chest* 94.4, pages 782–787. DOI: 10.1378/chest.94.4.782.
- Nardelli, P., Jimenez-Carretero, D., Bermejo-Pelaez, D., Washko, G. R., Rahaghi, F. N., Ledesma-Carbayo, M. J., and Estepar, R. S. J. (Nov. 2018). "Pulmonary Artery-Vein Classification in CT Images Using Deep Learning". In: *IEEE Trans Med Imaging* 37.11, pages 2428–2440. DOI: 10.1109/TMI.2018.2833385.
- Ninaber, M. K., Stolk, J., Smit, J., Roy, E. J. L., Kroft, L. J. M., Bakker, M. E., Vries Bouwstra, J. K. de, Schouffoer, A. A., Staring, M., and Stoel, B. C. (May 2015). "Lung structure and function relation in systemic sclerosis: Application of lung densitometry". In: *European Journal of Radiology* 84.5, pages 975–979. DOI: 10.1016/j.ejrad.2015.01.012.
- Park, S. O., Seo, J. B., Kim, N., Lee, Y. K., Lee, J., and Kim, D. S. (2011). "Comparison of usual interstitial pneumonia and nonspecific interstitial pneumonia: Quantification of disease severity and discrimination between two diseases on HRCT using a texture-based automated system". In: *Korean J Radiol* 12.3, pages 297–307. DOI: 10.3348/kjr.2011.12.3.297.
- Park, S. C., Tan, J., Wang, X., Lederman, D., Leader, J. K., Kim, S. H., and Zheng, B. (Feb. 2011). "Computer-aided detection of early interstitial lung diseases using low-dose CT images". In: *Phys Med Biol* 56.4, pages 1139–1153.
- Park, Y. S., Seo, J. B., Kim, N., Chae, E. J., Oh, Y. M., Lee, S. D., Lee, Y., and Kang, S.-H. (2008). "Texture-based quantification of pulmonary emphysema on high-resolution computed tomography: Comparison with density-based quantification and correlation with pulmonary function test". In: *Invest Radiol* 43.6, pages 395–402.
- Paszke, A., Chaurasia, A., Kim, S., and Culurciello, E. (2016). "ENet: A Deep Neural Network Architecture for Real-Time Semantic Segmentation". In: *CoRR* abs/1606.02147.

- Pei, X. (2015 2015). "Emphysema Classification Using Convolutional Neural Networks". In: *Intelligent Robotics and Applications, Icira 2015, Pt i 9244*, pages 455–461. DOI: 10.1007/978-3-319-22879-2_42.
- Peng, L., Lin, L., Hu, H., Li, H., Ling, X., Wang, D., Han, X., Iwamoto, Y., and Chen, Y.-W. (2018 2018). "Classification of Pulmonary Emphysema in CT Images Based on Multi-Scale Deep Convolutional Neural Networks". In: *2018 25th IEEE International Conference on Image Processing (ICIP)*.
- Putman, R. K., Hatabu, H., Araki, T., Gudmundsson, G., Gao, W., Nishino, M., Okajima, Y., Dupuis, J., Latourelle, J. C., Cho, M. H., El-Chemaly, S., Coxson, H. O., Celli, B. R., Fernandez, I. E., Zazueta, O. E., Ross, J. C., Harmouche, R., Estépar, R. S. J., Diaz, A. A., Sigurdsson, S., Gudmundsson, E. F., Eiríksdóttir, G., Aspelund, T., Budoff, M. J., Kinney, G. L., Hokanson, J. E., Williams, M. C., Murchison, J. T., MacNee, W., Hoffmann, U., O'Donnell, C. J., Launer, L. J., Harris, T. B., Gudnason, V., Silverman, E. K., O'Connor, G. T., R. Washko, G., Rosas, I. O., and Hunninghake, G. M. (2016). "Association between interstitial lung abnormalities and all-cause mortality". In: *JAMA-J Am Med Assoc* 315.7, pages 672–681. DOI: 10.1001/jama.2016.0518.
- Regan, E. A., Hokanson, J. E., Murphy, J. R., Make, B., Lynch, D. A., Beaty, T. H., Curran-Everett, D., Silverman, E. K., and Crapo, J. D. (2010). "Genetic epidemiology of COPD (COPDGene) study design". In: *COPD* 7.1, pages 32–43. DOI: 10.3109/15412550903499522.
- Richeldi, L., Bois, R. M. du, and Raghu, G. (2014). "Efficacy and Safety of Nintedanib in Idiopathic Pulmonary Fibrosis". In: *N Engl J Med* 370, pages 2071–2082.
- Rokach, L. (2010). "Ensemble-based classifiers". In: *Artif Intell Rev* 33.1-2, pages 1–39. DOI: 10.1007/s10462-009-9124-7.
- Ronneberger, O., Fischer, P., and Brox, T. (2015). "U-Net: Convolutional Networks for Biomedical Image Segmentation". In: *Medical Image Computing and Computer-Assisted Intervention, (MICCAI 2015)* 9351, pages 234–241.
- Roth, H. R., Farag, A., Lu, L., Turkbey, E. B., and Summers, R. M. (2015 2015). "Deep convolutional networks for pancreas segmentation in CT imaging". In: *Medical Imaging 2015: Image Processing* 9413, 94131G. DOI: 10.1117/12.2081420.

- Roth, H. R., Lu, L., Liu, J., Yao, J., Seff, A., Cherry, K., Kim, L., and Summers, R. M. (2016). "Improving Computer-Aided Detection Using Convolutional Neural Networks and Random View Aggregation". In: *IEEE Trans Med Imaging* 35.5, pages 1170–1181.
- Ryu, J. H., Colby, T. V., Hartman, T. E., and Vassallo, R. (Jan. 2001). "Smoking-related interstitial lung diseases: a concise review". In: *European Respiratory Journal* 17.1, pages 122–132. DOI: 10.1183/09031936.01.17101220.
- Ryu, Colby, T., Hartman, T., and Vassallo, R. (2001). "Smoking-related interstitial lung diseases: a concise review". In: *European Respiratory Journal* 17.1, pages 122–132. ISSN: 0903-1936.
- Seibold, M. A., Wise, A. L., Speer, M. C., Steele, M. P., Brown, K. K., Loyd, J. E., Fingerlin, T. E., Zhang, W., Gudmundsson, G., Groshong, S. D., Evans, C. M., Garantziotis, S., Adler, K. B., Dickey, B. F., Bois, R. M. du, Yang, I. V., Herron, A., Kervitsky, D., Talbert, J. L., Markin, C., Park, J., Crews, A. L., Slifer, S. H., Auerbach, S., Roy, M. G., Lin, J., Hennessy, C. E., Schwarz, M. I., and Schwartz, D. A. (Apr. 2011). "A Common MUC5B Promoter Polymorphism and Pulmonary Fibrosis". In: *N Engl J Med* 364.16, pages 1503–1512. DOI: 10.1056/NEJMoa1013660.
- Selvaraju, R. R., Cogswell, M., Das, A., Vedantam, R., Parikh, D., and Batra, D. (2017). "Grad-CAM: Visual Explanations from Deep Networks via Gradient-based Localization". In: *2017 Proc IEEE Int Conf Comput Vis (ICCV 2017)*, pages 618–626. DOI: 10.1109/ICCV.2017.74.
- Setio, A. A. A., Ciompi, F., Litjens, G., Gerke, P., Jacobs, C., Riel, S. J. van, Wille, M. M. W., Naqibullah, M., Sánchez, C. I., and Ginneken, B. van (2016). "Pulmonary Nodule Detection in CT Images: False Positive Reduction Using Multi-View Convolutional Networks". In: *IEEE Trans Med Imaging* 35.5, pages 1160–1169. ISSN: 0278-0062.
- Shakeri, M., Tsogkas, S., Ferrante, E., Lippe, S., Kadoury, S., Paragios, N., and Kokkinos, I. (2016). "Sub-Cortical Brain Structure Segmentation using F-CNN's". In: *2016 IEEE 13th International Symposium on Biomedical Imaging (ISBI)*, pages 269–272.
- Shin, H.-C., Roth, H. R., Gao, M., Lu, L., Xu, Z., Nogues, I., Yao, J., Mollura, D., and Summers, R. M. (May 2016). "Deep Convolutional Neural Networks for Computer-Aided Detection: CNN Architectures, Dataset Characteristics and Transfer Learning". In: *IEEE Transactions on Medical Imaging* 35.5, pages 1285–1298. ISSN: 0278-0062, 1558-254X. DOI: 10.1109/TMI.2016.2528162. (Visited on 02/06/2020).

- Shin, K. E., Chung, M. J., Jung, M. P., Choe, B. K., and Lee, K. S. (2011). "Quantitative computed tomographic indexes in diffuse interstitial lung disease: Correlation with physiologic tests and computed tomography visual scores". In: *J Comput Assist Tomogr* 35.2, pages 266–271. DOI: 10.1097/RCT.0b013e31820ccf18.
- Simonyan, K. and Zisserman, A. (2014). "Very deep convolutional networks for large-scale image recognition". In: *ICLR*.
- Sluimer, I. C., Waes, P. F. van, Viergever, M. A., and Ginneken, B. van (Dec. 2003). "Computer-aided diagnosis in high resolution CT of the lungs". In: *Medical physics* 30.12, pages 3081–3090. DOI: 10.1118/1.1624771.
- Smith, B. M., Austin, J. H. M., Jr., J. D. N., D'Souza, B. M., Rozenshtein, A., Hoffman, E. A., Ahmed, F., and Barr, R. G. (Jan. 2014). "Pulmonary Emphysema Subtypes on Computed Tomography: The MESA COPD Study". In: *American Journal of Medicine* 127.1, 94.e7. DOI: 10.1016/j.amjmed.2013.09.020.
- Snider, G. L., Kleinerman, J. L., Thurlbeck, W. M., and Bengali, Z. H. (1985). "The Definition of Emphysema - Report of a National-Heart-Lung-And-Blood-Institute, Division of Lung-Diseases Workshop". In: *American Review of Respiratory Disease* 132.1, pages 182–185.
- Sørensen, L., Shaker, S. B., and Bruijne, M. D. (2010). "Quantitative analysis of pulmonary emphysema using local binary patterns". In: *IEEE Trans Med Imaging* 29.2, pages 559–569. DOI: 10.1109/TMI.2009.2038575.
- Sorensen, L., Shaker, S. B., and Bruijne, M. de (2010a). "Quantitative Analysis of Pulmonary Emphysema Using Local Binary Patterns". In: *IEEE Transactions on Medical Imaging* 29.2, pages 559–569. DOI: 10.1109/TMI.2009.2038575.
- Sorensen, L., Shaker, S. B., and Bruijne, M. de (2010b). "Quantitative Analysis of Pulmonary Emphysema Using Local Binary Patterns". In: *IEEE Trans Med Imaging* 29.2, pages 559–569.
- Srivastava, N., Hinton, G., Krizhevsky, A., Sutskever, I., and Salakhutdinov, R. (2014). "Dropout: A simple way to prevent neural networks from overfitting". In: *J Mach Learn Res* 15, pages 1929–1958.

- Srivastava, V. and Purwar, R. K. (2019-Jun-26 2019). "Classification of CT Scan Images of Lungs Using Deep Convolutional Neural Network with External Shape-Based Features." In: *Journal of Digital Imaging*. DOI: 10.1007/s10278-019-00245-9.
- Stoel, B. C., Putter, H., Bakker, M. E., Dirksen, A., Stockley, R. A., Piitulainen, E., Russi, E. W., Parr, D., Shaker, S. B., and Reiber, J. H. C. e. a. (2008). "Volume Correction in Computed Tomography Densitometry for Follow-up Studies on Pulmonary Emphysema". In: *Proc Am Thorac Soc* 5.9, pages 919–924. DOI: 10.1513/pats.200804-040qc.
- Sugino, K., Ishida, F., Kikuchi, N., Hirota, N., Sano, G., Sato, K., Isobe, K., Sakamoto, S., Takai, Y., and Homma, S. (Feb. 2014). "Comparison of clinical characteristics and prognostic factors of combined pulmonary fibrosis and emphysema versus idiopathic pulmonary fibrosis alone". In: *Respirology* 19.2, pages 239–245. DOI: 10.1111/resp.12207.
- Szegedy, C., Liu, W., Jia, Y., Sermanet, P., Reed, S., Anguelov, D., Erhan, D., Vanhoucke, V., and Rabinovich, A. (2015). "Going deeper with convolutions". In: *Proc IEEE Comput Soc Conf Comput Vis Pattern Recognit*. Volume 07-12-June-2015, pages 1–9. DOI: 10.1109/CVPR.2015.7298594.
- Talmadge, Bradford, W. Z., Castro-Bernardini, S., Fagan, E. A., Glaspole, I., Glassberg, M. K., Gorina, E., Hopkins, P. M., Kardatzke, D., Lancaster, L., Lederer, D. J., Nathan, S. D., Pereira, C. A., Sahn, S. A., Sussman, R., Swigris, J. J., Noble, P. W., and Grp, A. S. (May 2014). "A Phase 3 Trial of Pirfenidone in Patients with Idiopathic Pulmonary Fibrosis". In: *New England Journal of Medicine* 370.22, pages 2083–2092. DOI: 10.1056/NEJMoa1402582.
- Tanizawa, K., Handa, T., Nagai, S., Hirai, T., Kubo, T., Oguma, T., Ito, I., Ito, Y., Watanabe, K., Aihara, K., Ikezoe, K., Oga, T., Chin, K., Izumi, T., and Mishima, M. (July 2015). "Clinical impact of high-attenuation and cystic areas on computed tomography in fibrotic idiopathic interstitial pneumonias". In: *Bmc Pulmonary Medicine* 15, page 74. DOI: 10.1186/s12890-015-0069-0.
- Thielen, A., Klus, H., and Mueller, L. (June 2008). "Tobacco smoke: Unraveling a controversial subject". In: *Experimental and Toxicologic Pathology* 60.2-3, pages 141–156.
- Tzilas, V. and Bouros, D. (2016). "Combined Pulmonary Fibrosis and Emphysema, a clinical review". In: *COPD Research and Practice* 2.2.

- Uchiyama, Y., Katsuragawa, S., Abe, H., Shiraishi, J., Li, F., Li, Q., Zhang, C., Suzuki, K., and Doi, K. (Sept. 2003). "Quantitative computerized analysis of diffuse lung disease in high-resolution computed tomography". In: *Medical physics* 30.9, pages 2440–2454. DOI: 10.1118/1.1597431.
- Umakoshi, H., Iwano, S., Inoue, T., Li, Y., and Naganawa, S. (Sept. 2017). "Quantitative evaluation of interstitial pneumonia using 3D-curved high-resolution CT imaging parallel to the chest wall: A pilot study". In: *Plos One* 12.9, e0185532. DOI: 10.1371/journal.pone.0185532.
- Uppaluri, R., Hoffman, E. A., Sonka, M., Hunninghake, G. W., and McLennan, G. (Feb. 1999). "Interstitial lung disease - A quantitative study using the adaptive multiple feature method". In: *American Journal of Respiratory and Critical Care Medicine* 159.2, pages 519–525. DOI: 10.1164/ajrccm.159.2.9707145.
- Uppaluri, R., Mitsa, T., Sonka, M., Hoffman, E. A., and McLennan, G. (July 1997). "Quantification of pulmonary emphysema from lung computed tomography images". In: *American Journal of Respiratory and Critical Care Medicine* 156.1, pages 248–254. DOI: 10.1164/ajrccm.156.1.9606093.
- Uppaluri, R., MITSA, T., SONKA, M., HOFFMAN, E. A., and McLENNAN, G. (1997). "Quantification of Pulmonary Emphysema from Lung Computed Tomography Images". In: *Am J Resp and Crit Care* 156.1, pages 248–254. DOI: 10.1164/ajrccm.156.1.9606093.
- Walsh, S. L. F., Calandriello, L., Silva, M., and Sverzellati, N. (Nov. 2018). "Deep learning for classifying fibrotic lung disease on high-resolution computed tomography: a case-cohort study". In: *Lancet Resp Med* 6.11, pages 837–845. DOI: 10.1016/S2213-2600(18)30286-8.
- Wang, C., Moriya, T., Hayashi, Y., Roth, H., Lu, L., Oda, M., Ohkubo, H., and Mori, K. (2019). "Weakly-supervised deep learning of interstitial lung disease types on CT images". In: *Proc. SPIE* 10950, 109501H (7 pp.)–109501H (7 pp.) DOI: 10.1117/12.2512746.
- Wang, J., Li, F., Doi, K., and Li, Q. (2009a). "Computerized detection of diffuse lung disease in MDCT: The usefulness of statistical texture features". In: *Phys Med Biol* 54.22, pages 6881–6899. DOI: 10.1088/0031-9155/54/22/009.

- Wang, J., Li, F., Doi, K., and Li, Q. (Nov. 2009b). "Computerized detection of diffuse lung disease in MDCT: the usefulness of statistical texture features". In: *Physics in Medicine and Biology* 54.22, pages 6881–6899. DOI: 10.1088/0031-9155/54/22/009.
- Washko, G. R., Hunninghake, G. M., Fernandez, I. E., Nishino, M., Okajima, Y., Yamashiro, T., Ross, J. C., Estépar, R. S. J., Lynch, D. A., Brehm, J. M., Andriole, K. P., Diaz, A. A., Khorasani, R., D'aco, K., Sciurba, F. C., Silverman, E. K., Hatabu, H., and Rosas, I. O. (2011). "Lung volumes and emphysema in smokers with interstitial lung abnormalities". In: *N Engl J Med* 364.10, pages 897–906. DOI: 10.1056/NEJMoA1007285.
- Washko, G. R., Hunninghake, G. M., Fernandez, I. E., Nishino, M., Okajima, Y., Yamashiro, T., Ross, J. C., Estépar, R. S. J., Lynch, D. A., Brehm, J. M., Andriole, K. P., Diaz, A. A., Khorasani, R., D'Aco, K., Sciurba, F. C., Silverman, E. K., Hatabu, H., Rosas, I. O., and Investigators, C. (Mar. 2011). "Lung Volumes and Emphysema in Smokers with Interstitial Lung Abnormalities". In: *New Eng J Med* 364.10, pages 897–906. DOI: 10.1056/NEJMoA1007285.
- Washko, G. R., Lynch, D. A., Matsuoka, S., Ross, J. C., Umeoka, S., Diaz, A., Sciurba, F. C., Hunninghake, G. M., Estépar, R. S. J., Silverman, E. K., Rosas, I. O., and Hatabu, H. (2010). "Identification of Early Interstitial Lung Disease in Smokers from the COPD Gene Study". In: *Acad Radiol* 17.1, pages 48–53. DOI: 10.1016/j.acra.2009.07.016.
- Washko, G. R., Lynch, D. A., Matsuoka, S., Ross, J. C., Umeoka, S., Diaz, A., Sciurba, F. C., Hunninghake, G. M., Estépar, R. S. J., Silverman, E. K., Rosas, I. O., and Hatabu, H. (2010). "Identification of Early Interstitial Lung Disease in Smokers from the COPD Gene Study". In: *Academic Radiology* 17.1, pages 48–53.
- Webb, W. R. (2006). "Thin-section CT of the secondary pulmonary lobule: Anatomy and the image - The 2004 Fleischner lecture". In: *Radiology* 239.2, pages 322–338.
- Xu, Y., Beek, E. J. R. van, Yu, H., Guo, J., McLennan, G., and Hoffman, E. A. (Aug. 2006). "Computer-aided classification of interstitial lung diseases via MDCT: 3D adaptive multiple feature method (3D AMFM)". In: *Academic Radiology* 13.8, pages 969–978. DOI: 10.1016/j.acra.2006.04.017.
- Yamashiro, T., Matsuoka, S., Estépar, R. S. J., Bartholmai, B. J., Diaz, A., Ross, J. C., Murayama, S., Silverman, E. K., Hatabu, H., and Washko, G. R. (Feb. 2011). "Kurtosis and Skewness of Density Histograms on Inspiratory and Expiratory CT Scans in Smokers". In: *COPD-Journal of*

Chronic Obstructive Pulmonary Disease 8.1, pages 13–20. DOI: 10.3109/15412555.2010.541537.

Zhao, W., Xu, R., Hirano, Y., Tachibana, R., and Kido, S. (2015). “A sparse representation based method to classify pulmonary patterns of diffuse lung diseases”. In: *Comput Math Methods Med* 2015. DOI: 10.1155/2015/567932.