

Lagrangian Skeleton on the Gulf of Mexico Oil Spill

C. Mendoza^{1,*}, A.M. Mancho²⁾

1) Universidad Politécnica de Madrid. ETSI Navales.

Av. Arco de la Victoria s/n. 28040 Madrid, Spain

2) Instituto de Ciencias Matemáticas. CSIC-UAM-UC3M-UCM.

Serrano 121. 28006 Madrid, Spain

* e-mail: carolina.mendoza@upm.es

Abstract

The understanding of the circulation of ocean currents, the exchange of CO_2 between atmosphere and oceans, and the influence of the oceans on the distribution of heat on a global scale is key to our ability to predict and assess the future evolution of climate [1, 2]. Global climate change is affecting sea breathing through mechanisms not yet understood [3]. The ocean is important in the regulation of heat and moisture fluxes, and oceanic physical and bio-geochemical processes are major regulators of natural greenhouse gases. Understanding how oceans mix their waters is key to provide sound forecasts on the climate [1]. Global change also affects marine biodiversity and threatens the survival of ecosystems and exploitable resources. To predict not only the effects of global change on the oceans, but also the response time of climate feedback requires to improve detection systems and to open new lines of research.

We use a novel Lagrangian descriptor (function M , introduced in [4, 5]). It is based on the measure of the arclength of particle trajectories on the ocean surface at a given time. In [6, 7, 8] this technique has been proven to be successful for characterizing the Kuroshio current.

We employ this tool on velocity data sets on the Gulf of Mexico obtained from HYCOM project. The tool identifies the underlying Lagrangian structures in the oceanic currents. In particular invariant manifolds, hyperbolic and non-hyperbolic flow regions are detected. We study the influence of these structures on particle motions on the oil spill.

We acknowledge to CESGA for support with the supercomputer FINIS TERRAE. Thanks for support by grants: CSIC-OCEANTECH, I-Math C3-0104, MICINN-MTM2008-03754 and MTM2008-03840-E, and C. Madrid-SIMUMAT.

References

- [1] S. Bowen, M.S. Lozier, S. F. Gary, and C.W. Bning. *Nature* 459, 243-247, 2009.
- [2] K. Katija, and J.O. Dabiri. *Nature* 460, 624-626, 2009.
- [3] Q. Schiermeier. *Nature* 447, 522-524, 2007.
- [4] J. A. Jiménez Madrid, A.M. Mancho. Distinguished trajectories in time dependent vector fields. *Chaos* 19 (2009), 013111-1-013111-18.
- [5] C. Mendoza, A.M. Mancho. The hidden geometry of ocean flows. *Physical Review Letters* 105 (2010), 3, 038501-1-038501-4.
- [6] A.M. Mancho, S. Wiggins, A. Turiel, E. Hernández-Garca, C. López, E. García-Ladona. *Nonlinear Processes in Oceanic and Atmospheric Flows. Nonlinear Proc. Geoph* 17 (2010), 3, 283-285.
- [7] C. Mendoza, A.M. Mancho, Marie-Hélène Rio. The turnstile mechanism across the Kuroshio current: analysis of dynamics in altimeter velocity fields. *Nonlinear Proc. Geoph* 17 (2010), 2, 103-111.
- [8] C. Mendoza, A.M. Mancho. The Lagrangian description of aperiodic flows: a case study of the Kuroshio Current. preprint (2011).