



POSTER PRESENTATION

Open Access

Component resolved diagnosis in baker's asthma

Cristina Gómez-Casado^{1*}, María Garrido-Arandia², Catia Pereira³, Ana Aranda⁴, Manuela Catarino³, Alicia Armentia⁵, Santiago Quirce⁶, Paloma Campo⁴, Araceli Díaz-Perales²

From 5th International Symposium on Molecular Allergology (ISMA 2013)
Vienna, Austria. 6-7 December 2013

Background

Baker's asthma is one of the most common types of occupational asthma and its prevalence is increasing in the last years. Diagnosis of occupational asthma is complex. The poor specificity of current diagnostic approaches may be associated with insufficient purity of wheat extracts or lack of inclusion of major allergens in them. In this work, we use microarray technology to characterize the allergenic profiles of baker's asthma patients from three regions in Spain and to analyze the influence of other environmental allergens on the sensitization pattern.

Methods

A panel of wheat allergens was purified from natural sources and printed on a protein microarray by standard methods. Additionally, representative aeroallergens were included in the protein microarray. Individual sera from three groups of patients (baker's asthma-BA-, seasonal rhinitis-SR- and wheat food allergy-WFA-) and three regions in Spain (Madrid, Malaga and Valladolid) were incubated with the protein microarray and compared.

Results

WTAI-CM16 and Tri a 14 were the most prevalent allergens in the BA group (54 and 45%, respectively), covering a total of 64% of the BA population. Patients with seasonal rhinitis (SR) were not sensitized to wheat allergens. Tri a 14, the wheat LTP, was only recognized by BA patients. Although geographically different environmental allergenic load existed in the three regions, BA sensitization profiles were comparable.

Conclusion

Baker's asthma is a disease associated with the daily handling of wheat flour, regardless of local aeroallergens.

Authors' details

¹CBGP (UPM-INIA) Technical University Madrid, CBGP (UPM-INIA) Campus Montegancedo, Autovía M-40 Salida 38N 36S, Pozuelo de Alarcón (Madrid), 28223, Spain. ²Technical University CBGP (UPM-INIA), Pozuelo de Alarcón (Madrid), Spain. ³University of Lisbon, Faculty of Pharmacy, Lisbon, Portugal. ⁴Civil Hospital of Malaga, IMABIS Foundation, Malaga, Spain. ⁵Río Ortega University Hospital, Allergy Service, Valladolid, Spain. ⁶La Paz University Hospital, Allergy Service, Madrid, Spain.

Published: 17 March 2014

doi:10.1186/2045-7022-4-S2-P35

Cite this article as: Gómez-Casado et al.: Component resolved diagnosis in baker's asthma. *Clinical and Translational Allergy* 2014 **4**(Suppl 2):P35.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹CBGP (UPM-INIA) Technical University Madrid, CBGP (UPM-INIA) Campus Montegancedo, Autovía M-40 Salida 38N 36S, Pozuelo de Alarcón (Madrid), 28223, Spain

Full list of author information is available at the end of the article