

Article

Evaluating the Water Footprint of the Mediterranean and American Diets

Alejandro Blas ^{1,2,*}, Alberto Garrido ^{1,2} and Bárbara A. Willaarts ^{1,2}

¹ Water Observatory of the Botin Foundation, Madrid 28001, Spain; alberto.garrido@upm.es (A.G.); bwillaarts@gmail.com (B.A.W.)

² Research Center for the Management of Environmental and Agricultural Risks (CEIGRAM), Universidad Politécnica de Madrid, Madrid 28040, Spain

* Correspondence: a.blas@upm.es; Tel.: +34-91-452-4815

Academic Editors: Arjen Y. Hoekstra, Ashok K. Chapagain and Pieter R. van Oel

Received: 29 July 2016; Accepted: 8 October 2016; Published: 13 October 2016

Abstract: Global food demand is increasing rapidly as a result of multiple drivers including population growth, dietary shifts and economic development. Meeting the rising global food demand will require expanding agricultural production and promoting healthier and more sustainable diets. The goal of this paper is to assess and compare the water footprint (WF) of two recommended diets (Mediterranean and American), and evaluate the water savings of possible dietary shifts in two countries: Spain and the United States (US). Our results show that the American diet has a 29% higher WF in comparison with the Mediterranean, regardless of products' origin. In the US, a shift to a Mediterranean diet would decrease the WF by 1629 L/person/day. Meanwhile, a shift towards an American diet in Spain will increase the WF by 1504 L/person/day. The largest share of the WF of both diets is always linked to green water (62%–75%). Grey water in the US is 67% higher in comparison with Spain. Only five products account for 36%–46% of the total WF of the two dietary options in both countries, being meat, oil and dairy products the food items with the largest WFs. Our study demonstrates that adopting diets based on a greater consumption of vegetables, fruits and fish, like the Mediterranean one, leads to major water savings.

Keywords: dietary shifts; sustainability; consumption patterns; water consumption; green water; blue water

1. Introduction

World population will reach 9 billion by 2050 and global food production will have to increase between 70% and 100% [1]. Meeting the future food demand is a major challenge for the current food production systems, and even more doing it without compromising the environmental integrity [2]. Converting more land into cultivation and increasing crop yields have been largely promoted as key solutions to increase food production globally [3,4]. These production-oriented policies explain why global agricultural area has expanded over 11% since 1960 [5], and cereal crop production has almost doubled [4].

However, the increasing competition for land, water and energy [6], as well the multiple impacts agriculture has on the environment, require rethinking possible approaches to face the daunting challenge of satisfying increasing food demand within a resource-constrained planet. Many of the current efforts are placed in increasing the sustainability of food production through so-called “sustainable intensification”, which will require improving water use or nitrogen-phosphorus efficiency, implementing ecological-based management practices, judicious use of pesticides, and also reconverting much of the livestock production practices [4].

But beyond any productivity improvements, enhancement of consumption patterns and the promotion of healthier and sustainable diets will be of major importance to achieve positive

environmental effects [7,8]. Recently several studies have analyzed and recognized the importance of diets in future food security and sustainability [9–11]. The literature shows that larger environmental impacts originate from animal products-based diets in comparison to less meat-based and vegetarian diets [8,9,11–14]. Jalava et al. [9] compared the current global consumption patterns with diets containing low contents of animal products, and found that lower intake of animal products lead to important water savings, i.e., lower water footprint.

In the course of the last decades, several methodologies, including the water footprint assessment (WFA) [15], and the life cycle analysis (LCA) [16], have been developed to assess the impacts on water resources linked to food production and consumption patterns. Both approaches involve several-step process and the suitability of one over the other very much depends on the project goal. WFA is a suitable approach particularly when the overall purpose of the assessment is to identify options for water savings, reallocation and better management, and also in order to raise awareness about water issues [16]. A WFA involves four steps: (1) definition of the scope and goals of the assessment; (2) water accounting; (3) sustainability assessment; and (4) response formulation. There is ample literature exemplifying the usefulness of WFAs [17–21]. As opposed to the WFA, the LCA is more suitable when the goal is to evaluate the environmental impacts linked to different human activities, where water use is one among many different impacts that can be assessed [16]. LCA also involves four steps: (1) definition of the goal and scope; (2) inventory; (3) impact assessment and (4) interpretation. Organizations like the FAO have adopted the LCA methodology and the associated ISO 14046 [22], within its environmental sustainability program.

Both the WFA and the LCA rely in the use of quantitative indicators (e.g., the water footprint), although in different phases of the assessment [16]. Several authors have compared both methodologies [16,23], and applied them to different food products, such as tomato sauce [24], tea and margarine [25], biomass production from energy crops [26], or broccoli [27] to assess the advantages and disadvantages of each one.

A large number of the studies addressing the issue of diets and water impacts relied in the methodology proposed by the WFA. Vanham et al. [12], compared the water footprint (WF) of current consumption patterns with healthier and vegetarian diets in Europe, and found that improving diets might result in reductions of the diet's WF between 974 and 1611 L per person and day, equivalent to savings of 23%–38%. These authors also concluded that the consumption of animal products accounts for the largest share (46%) of the WF linked to the prevailing diets in Europe. Other studies comparing three European diets (current, healthy and vegetarian) across four European areas (west, north, south and east) showed that in all zones adopting healthy and vegetarian diets could lead to substantial WF reductions (up to 41%) [13]. Similar results have been obtained in other studies comparing different diets and dietary patterns at the country [11,14,28] and city level [29,30].

In much of the developed world, dietary shifts are causing important health problems [8]. This circumstance is driving national health and/or food agencies to campaign in favor of investing public funds to raise awareness among citizens about the importance of adopting healthier food habits. Countries like Spain are placing large efforts to reverse the growing obesity problems and involving different public institutions in the promotion of the Mediterranean diet [31,32]. In fact the Mediterranean diet has been recognized in many countries as a key strategy to improve a population's health with local, traditional and seasonal products [31,33]. Also, it has been recognized by UNESCO as a cultural World Heritage [34], and was selected by the UN Food Agricultural Organization (FAO) to develop a methodological approach to assess sustainability across different agro-ecological zones [35]. In fact, the Mediterranean diet is appreciated for its lower environmental impacts in relation with other meat-based diets [32,36].

Other countries facing serious obesity problems, like the United States of America (US), are also investing large efforts to reverse this trend [37]. In fact, the United States Department of Agriculture (USDA) in an attempt to raise awareness among consumers has elaborated several national dietary guidelines and recommended diets [37,38].

Yet, most of the studies published addressing the composition of diets and associated water savings are focused on making comparisons between different dietary patterns obtained from current consumption patterns of annual statistics, like the food balance sheets of FAO [39]. Little work has been done to compare real, local or seasonal recommended diets. Comparing recommended diets, elaborated with national and traditional recipes, dishes and products, can provide insightful results about the relationship between real consumption patterns and environmental effects, and the ingredients that generate the largest water savings.

Accordingly, the main objective of this paper is to assess the WF of two contrasting and recommended food diets, i.e., the Mediterranean and American, and to evaluate the water savings of possible dietary shifts in the two countries (Spain and US). Also, we attempt to deepen in the understanding of the relationship between products origin and consumption patterns, and their influence on the total WF and the diet's sustainability.

2. Materials and Methods

2.1. Menus and Diets Configuration

To characterize the composition and product quantities of the Mediterranean diet, two seasonal weeklong menus were defined (i.e., winter and summer) using the food guidelines elaborated by the Mediterranean Diet Foundation [40]. Each week menu included four daily meals: breakfast, morning snack, lunch and dinner. Dishes and product composition of each meal, as well as recommended daily intake quantities of each product were defined using traditional Mediterranean recipes (see Tables S1 and S2). The winter menu (Menu 1) was configured with winter seasonal products, whereas the summer menu (Menu 2) in the Mediterranean diet was configured with seasonal summer products. In total, over 116 products have been included in the elaboration of the two menus.

The American recommended diet (USDAR) was configured following the guidelines of the Center for Nutrition Policy and Promotion (CNPP) of the US Department of Agriculture [38] and the study by Haven et al. [37]. As with the Mediterranean diet, two weeklong menus (Menu 1 and Menu 2) were configured, each one containing four daily meals (breakfast, morning snack, lunch and dinner). No differences between menus in terms of seasonal products could be made with the USDAR. Recipes, product composition and weights for the two American menus were also defined using traditional recipes from the US (Tables S3 and S4). Overall, 103 products were included in the elaboration of the USDAR diet.

2.2. Water Footprint Calculation

2.2.1. Recommended Diets

We used the WFA approach, and particularly we focused on phase two (i.e., water accounting), to assess the water embedded in each menu and diet. Specifically we calculated the water footprint (WF) per person per day of the Mediterranean and USDAR diets as defined in the Global Water Footprint Standard [15].

The WF of a product is understood as the direct and indirect appropriation of freshwater resources required to produce the good, this being the final result the sum of three components: green, blue and grey [41]. Green WF refers to the rainwater stored in the soils and directly evapotranspired by crop products. The blue WF refers to the volume of surface and groundwater embedded in the production of a good. In agriculture, the blue WF refers to the total volume of irrigated water that is evapotranspired by a crop, and embedded in the production of livestock products. Finally, grey WF is an indicator of water quality degradation, and refers to the volume of freshwater required to assimilate the load of pollutants generated along the production chain of a product, in order to reach the quality standards established in the environmental regulations [15]. The WF of a product is the sum of all the water consumed along its different production steps [41]. When calculating the WF of a diet, we have only

considered the amount of water consumed in the production of each food item, without considering additional water requirements for cooking purposes (e.g., boiling, washing, etc.). This assumption was made due to the lack of detailed data regarding the amount of household water used for cooking and the fact that the largest fraction of the WF of food products lies at the field level, i.e., agricultural production [20].

To estimate the WF of the different weeklong menus we relied in the global WF database of crops [41], and livestock products [42]. Both databases provide average values of green, blue and grey WF of each product (in m³/t or L/kg) for the time series 1996–2005.

Considering the impossibility of determining the origin of all consumed products, we used national values to estimate the WF of the different menus. We estimated the WF of the Mediterranean diet assuming that all products consumed are produced in Spain, whereas for the USDAr we assumed that all products have been produced in the US. Products like coffee, cocoa, pineapple, pepper, cinnamon and mustard are not produced in either of the two countries and therefore they need to be imported. The WF of each imported product was calculated at the weighted average of the WF this product has in the main producer countries from which Spain and the US import them. The weighting factor is calculated based on the ratio of imports by origin of the product in each of the two countries (i.e., Spain and US). For instance the WF of coffee consumed in Spain was estimated as a weighted average of the WF coffee has in the major production centers from which Spain imports this product (i.e., Vietnam (40%), Brazil (50%) and Colombia (10%)). To define the origin of each imported product we used trade matrix from FAOSTAT [5], for the time series 1993–2013.

For each imported product in Spain, we used the following countries' WF data: Coffee, data from Vietnam (40%), Brazil (50%) and Colombia (10%); Cocoa, data from Ivory Coast (50%), Ghana (30%) and Indonesia (20%); Pineapple, data from Costa Rica (100%); Pepper, data from Vietnam (60%), Indonesia (30%) and India (20%); Cinnamon, data from Indonesia (50%) and China (50%); and Mustard, data from Canada (100%).

Regarding products imported into the US, we used the following countries' WF data: pepper, cinnamon and cocoa as in Spain (same WF values); vanilla, data from Indonesia (60%) and Madagascar (40%); cilantro/parsley/sesame, data from the UK (50%) and Russia (50%); clove, data from Indonesia (100%); and nutmeg, data from Guatemala (50%) and Indonesia (50%).

Lastly, saltwater fish and seafood (i.e., hake, tuna, sea bass, megrim, squid, mussels, cod, octopus, salmon, trout, shrimps and sardines) were assigned a WF equivalent to zero, since this study only evaluates the WF of raw products and does not include the water requirements for cooking. Other studies have calculated the fish WF but most of them refer to aquaculture production [43].

Hence, the daily WF of a menu k ($WF_{\text{menu } k}$, in liters per person and day) was calculated as:

$$WF_{\text{menu } k} = \sum_{j=1}^n (\text{Green } WF_j + \text{Blue } WF_j + \text{Grey } WF_j) \times W_j / 7 \quad (1)$$

where $(\text{Green } WF_j + \text{Blue } WF_j + \text{Grey } WF_j)$ is the total WF of a product j (in L/kg) and W_j is the average intake per person and week (in kg per person). Accordingly, the average WF of a diet i ($WF_{\text{diet } i}$, in liters per person and day) was estimated as:

$$WF_{\text{diet } i} = \sum_{K=1}^2 (WF_{\text{menu } k}) / 2 \quad (2)$$

To discern the relative weight that the different products have in the WF_{diet} (the 116 products identified in the Mediterranean menus and the 103 of the two American menus), we grouped them into 11 different food groups: (1) meat, fish and animal fats; (2) dairy products; (3) oil and vegetable fats; (4) legumes and nuts; (5) cereals and potatoes; (6) eggs; (7) vegetables; (8) sugar; (9) cocoa, chocolate and vanilla; (10) fruits and (11) drink and others.

2.2.2. Impacts of Shifting Diets

To assess the water impacts of changing diets we also estimated the WF_{diet} associated to a potential shift in the diets in the two countries—i.e., Spain adopting an American diet and the US adopting a Mediterranean diet. We assumed that all consumed products are also produced nationally, except for those products that are not produced in either of the two countries and need to be imported. Again, we relied on the WF databases of agricultural [41] and livestock products [42] to estimate the WF_{diet} .

3. Results

The WF_{menu} and WF_{diet} of the recommended Mediterranean and USAr diets, as well as the potential shift in dietary habits, are shown in Figure 1 for Spain (a) and the US (b). The WF_{menu} and WF_{diet} of USAr diet are higher than the Mediterranean diet, irrespectively of the products' origin (Spain or US).




Figure 1. Green, blue and grey water footprint (WF) for WF_{menu1} , WF_{menu2} and the total WF diet of Mediterranean diet (WF_{diet}), and WF_{menu1} , WF_{menu2} and the total WF diet of USAr diet (WF_{diet}), for: (a) Spain and (b) the US.

In Spain, the WF_{diet} of the Mediterranean diet is 5276 L per person and day, but adopting a USAr diet will increase the WF_{diet} nearly 29%, up to 6780 L per person and day. The majority of this increase results from the rise in green water, followed by blue water, and to a lesser extend grey water. In the US, the WF_{diet} of the USAr is 5632 L per person and day. Shifting towards a Mediterranean diet (4003 L per person and day), will decrease the WF_{diet} by 29%. Larger savings will be achieved in terms of green water (−1392 L/person and day) and grey water. However, in this diet-shift scenario blue WF will increase by 24%.

Green WF accounted for the largest share of both WF_{menu} and WF_{diet} in the two countries. In Spain, green WF contributes to 75% of WF_{diet} for the Mediterranean diet and 71% for the USAr diet. In the US, green WF accounted for 62% of WF_{diet} in the Mediterranean and 69% in the USAr diet. Blue WF was the second largest fraction in WF_{diet} in Spain for both Mediterranean (16%) and USAr diet (19%). On the other hand, grey WF accounted for the second largest fraction of WF_{diet} in the US for Mediterranean (20%) and USAr diets (21%).

If we took into account only the blue and green WF components of WF_{menu} and WF_{diet} , in order to discern the impacts of water resources quantity, there would be water savings equivalent to 1277 L per person and day in Spain by consuming a Mediterranean diet instead of a USAr diet. Similar values

were obtained in the US for green and blue WFs, where changing from a USDAR to a Mediterranean diet will imply a net reduction of 1252 L of water per person and day.

Comparing the WF_{diet} in the two different countries for the same diet revealed that, while the Mediterranean diet is more efficient in terms of water consumption in all cases, adopting a Mediterranean diet in the US would deliver greater water savings (up to 24% or 1273 L per person and day in comparison with the same diet consumed in Spain). This is due to the greater water productivity of the US's agriculture. On the other hand, USDAR diet in Spain will increase the WF_{diet} 20% (1148 L/person and day) in comparison with the consumption of this diet in the US.

Figure 2 shows the contribution of the different groups of products to the WF_{diet} in Spain and US. Dairy products, oil and vegetable fats, and meat, fish and animal fats accounted for the 68% of the green component of the Mediterranean diet in Spain (up to 2662 L per person and day) (Figure 2a). A shift towards an American diet of Spanish consumers would increase the consumption of dairy and meat, fish and animal products groups by 30% (+547 L per person and day) and therefore enlarge the green WF_{diet} up to 4808 L/person and day. Legumes, cereals and potatoes, and eggs also represent a significant share of the green fraction of the WF_{diet} under a Mediterranean and USDAR diet in Spain (22% and 19% respectively).

Changing the consumption patterns in the US and adopting a Mediterranean diet would deliver significant green water savings (Figure 2a). Many of these water savings are related to lower green WF values associated to oil and vegetable fats (70% lower, 407 L/person and day less), meat and fish products (22% lower, 211 L/person and day less) and cocoa, chocolate and vanilla (93% lower, 257 L/person and day less) in the Mediterranean diet.

The share of blue WF among product groups is more evenly distributed in the case of the Mediterranean diet in Spain (Figure 2b). On the other hand, legumes and nuts, and cereals and potatoes groups account for the 45% of the blue component of the USDAR diet (572 L/person and day). So the largest differences in terms of blue water WF among diets in Spain are due to the higher water consumption of legumes and nuts (almost six times higher, +273 L/person and day) as well as cereals and potatoes (almost five times higher, +195 L/person and day) in the USDAR diet in comparison with the Mediterranean one.

In the case of US, the Mediterranean diet has higher blue WF than USDAR diet. Legumes and nuts, oil and vegetable fats, and vegetables account for 62% of the blue component of the Mediterranean diet in the US (up to 455 L per person and day). A shift towards a Mediterranean diet of American consumers would increase the blue WF of these products groups: legumes and nuts (68% higher, +42 L/person and day), oil and vegetable fats (nearly five times higher, +166 L/person and day) and vegetables (nearly three times higher, +89 L/person and day).

Concerning grey WF (Figure 2c) in both countries, the USDAR diet has higher values than the Mediterranean one, especially in US. Meat, fish and animal fats and dairy products account for 55% of the grey WF of the Mediterranean diet in Spain (260 L per person and day). Consuming a USDAR diet would lead to an increase of dairy products (46% higher, 70 L/person and day) and oil and vegetable fats (35 times higher, 67 L per person and day). Also, very significantly, the WF of cocoa, chocolate and vanilla group is higher and increases up to 480 times in changing to a USDAR diet (53 L/person and day).

In the case of the US, the legumes and nuts group alone accounts for almost 50% of the grey component of the USDAR diet (576 L/person and day). A change to a Mediterranean diet would afford a 35% reduction (202 L/person and day) in this product group.

The analysis of the individual products' WF reveals that a limited number of products contribute the most to the green, blue, grey and total WF_{diet} account for up to 36%–46% of the total in both countries and dietary options. Tables 1 and 2 show the five products that contribute the most to the green, blue, grey and total WF_{diet} for both diets and countries. Olive oil is the product which accounts the most to the WF_{diet} of the Mediterranean diet, both in Spain and the US, as shown in Tables 1 and 2 respectively. On the other hand, in the USDAR diet, semi-skimmed milk is the product that accrues the

largest share of the WF_{diet} (16%, equivalent to 1085 L per person day) in Spain (Table 1). While in the US (Table 2) the product that claims the largest WF_{diet} is beef meat (14%, 789 L per person and day).

Most of the products that influence the most for green, blue, grey (and thus in WF_{diet}) for both dietary options and countries originate from only three products groups: (a) meat, fish and animal fats; (b) dairy products; and (c) oils and vegetable fats. Moreover, products from the group of legumes and nuts account for the major part of the grey WF for both diets in US.

Table 1. The five products that contribute the most in L/person and day (lpd) and % to the green, blue, grey and total WF_{diet} for the Mediterranean and US diets in Spain.

Mediterranean Diet											
Green WF			Blue WF			Grey WF			WF_{diet}		
Product	lpd	%	Product	lpd	%	Product	lpd	%	Product	lpd	%
Olive oil	867	22%	Olive oil	207	24%	Milk	61	13%	Olive oil	1055	20%
Milk	355	9%	Milk	52	6%	Eggs	43	9%	Milk	475	9%
Beef meat	276	7%	Sugar	52	6%	Cheese	33	7%	Eggs	316	6%
Eggs	236	6%	Asparagus	43	5%	Beef meat	28	6%	Beef meat	316	6%
Cheese	197	5%	Eggs	34	4%	Chicken	28	6%	Cheese	264	5%
Rest	2010	51%	Rest	474	55%	Rest	279	59%	Rest	2847	54%
USDAr Diet											
Green WF			Blue WF			Grey WF			WF_{diet}		
Product	lpd	%	Product	lpd	%	Product	lpd	%	Product	lpd	%
SK milk ¹	817	17%	Soy milk	267	21%	SK milk ¹	140	20%	SK milk ¹	1084	16%
Beef meat	529	11%	Oats	165	13%	Margarine	70	10%	Beef meat	610	9%
Margarine	385	8%	SK milk ¹	115	9%	Beef meat	56	8%	Margarine	542	8%
Chicken	192	4%	Sesame	76	6%	Vanilla	49	7%	Oats	407	6%
Oats	192	4%	Margarine	51	4%	Oats	42	6%	Soy milk	339	5%
Rest	2692	56%	Rest	598	47%	Rest	343	51%	Rest	3796	56%

Note: ¹ SK milk = Semi-skimmed milk.

Table 2. The five products that contribute the most in L/person and day (lpd) and % to the green, blue, grey and total WF_{diet} for the Mediterranean and US diets in the US.

Mediterranean Diet											
Green WF			Blue WF			Grey WF			WF_{diet}		
Product	lpd	%	Product	lpd	%	Product	lpd	%	Product	lpd	%
Milk	323	13%	Olive oil	212	29%	Chickpeas	126	16%	Olive oil	400	10%
Beef meat	298	12%	Almonds	51	7%	Almonds	79	10%	Milk	400	10%
Olive oil	174	7%	Asparagus	44	6%	Lentils	63	8%	Beef meat	320	8%
Bread	124	5%	Hazelnuts	37	5%	Hazelnuts	55	7%	Chickpeas	160	4%
Flour	124	5%	Milk	29	4%	Milk	47	6%	Pork Meat	160	4%
Rest	1439	58%	Rest	358	49%	Rest	419	47%	Rest	2562	64%
USDAr Diet											
Green WF			Blue WF			Grey WF			WF_{diet}		
Product	lpd	%	Product	lpd	%	Product	lpd	%	Product	lpd	%
Beef meat	736	19%	Oats	77	13%	Beans	268	23%	Beef meat	788	14%
SK milk ¹	387	10%	Vanilla	53	9%	Chickpeas	152	13%	SK milk ¹	451	8%
Vegetable oil	232	6%	Apple juice	35	6%	Coffee	128	11%	Beans	338	6%
Margarine	194	5%	SK milk ¹	35	6%	Lentils	105	9%	Oats	282	5%
Oats	194	5%	Rice	35	6%	SK milk ¹	58	5%	Coffee	225	4%
Rest	2131	55%	Rest	355	60%	Rest	455	39%	Rest	3548	63%

Note: ¹ SK milk = Semi-skimmed milk.




Figure 2. Characterization of the green (a), blue (b) and grey (c) components of the WF_{diet} of the Mediterranean and USDAr diets in Spain and the US for the different product groups.

4. Discussion

USDAr diet has a greater average intake quantity of dairy products (756 g/person and day) than the Mediterranean diet (405 g/person and day). Also, the USDAr diet has a greater share of legumes and nuts daily intake (246 g/person and day, in comparison with 89 g/person and day in the Mediterranean diet). On the other hand, in the Mediterranean diet the intake quantity of vegetables is higher (1021 g/person and day versus 613 in USDAr diet), as well as fruits (624 g/person in the Mediterranean diet in comparison with the 545 g/person and day of the USDAr diet). Regarding meat, fish and animal fats food group, they have similar consumption rates of meat and animal fats: 187 g/person and day in the Mediterranean diet and 173 g/person and day in the USDAr one. However, in the Mediterranean diet the consumption of fish is much higher (146 versus 48 g/person and day), which significantly reduces the WF_{diet} linked to the consumption of animal proteins. In short, the Mediterranean diet relies on a larger intake of vegetables, fruits and fish, while in the USDAr diet the fraction of dairy, legumes and nuts products is larger. Tables S1–S4 contain a full description of the products and quantities included of each menu for both diets.

4.1. Consumption Patterns: Water Footprint of Mediterranean and USDAr Diets

In both countries, adopting a Mediterranean diet would lead to major water savings per person. In Spain, maintaining a Mediterranean diet will save up to 1504 L per person per day ($\approx 29\%$); while in the US, shifting into a Mediterranean diet would deliver even greater benefits (net water savings of 1629 L per person per day, equivalent to 29%). The Mediterranean summer menu (menu 2) has a WF 7% smaller than the winter menu (menu 1) in Spain and 1.5% in the US. These differences can be attributed to the larger calorie content of the winter menu, since the proportion of food groups is similar in the two menus. Our results stand within the ranges of Vanham et al. [13], who found that healthy diets lead to water savings ranging between 3% and 30%. Moreover, our results are consistent with those studies where diets based on the consumption of vegetables and fruits [8,9,11–14,28] or totally vegetarian [44,45] have a smaller WF compared to animal products-based or non-vegetarian diets.

Some authors have shown similar water savings (up to 33%) when western dietary patterns are replaced by the adoption of a Mediterranean diet [32]. Other comparisons between the Mediterranean recommended diet and other western meat-based dietary patterns have been reported [8,28,32], concluding also that there are major water savings using a Mediterranean diet. But little is known about comparisons with other real and official recommended diets.

4.2. Products' Origins Matter: Water Footprints in Spain and US

This study also shows that production factors also explain part of the changes in WF_{diet} beyond the shifts in diets. As shown in Figure 1 showed, a shift towards a Mediterranean diet in the US will decrease the WF_{diet} up to 4003 L per person and per day, a value below the WF_{diet} calculated for a Mediterranean diet in Spain (5276 L per person and per day). Also, looking only at the consumptive fraction of the WF_{diet} (green WF + blue WF), the lowest values appear for Mediterranean diet in the US. This means that, not only consumption patterns, but also products' origins (prevailing climate and production conditions) are very significant factors to be taken into account for minimizing WFs. It also implies that the mixture of dietary patterns based on the consumption of vegetables, fish and fruits, followed by improved production conditions (i.e., lower WF), can contribute to achieve major water savings.

In countries like Spain, high values for green and blue WF can be explained because of the prevailing semi-arid climate conditions. Several studies have shown this close relationship between drier climate conditions and higher values of green and blue WFs [12,13,46]. High temperatures and low rainfall contribute to higher WF in L/kg of products, because of the lower yields under large production areas [47]. Further, in semi-arid conditions there is a greater need for irrigation, which increases the blue WF [13,48].

Our results demonstrate that the green and blue components of the WF_{diet} are higher in Spain compared to the US's regardless of the type of diet. These results are mainly due to the differences of green WF values in the US in comparison with Spain of oil and vegetable fats in the Mediterranean diet (80% lower in the US, -688 L/person and day), and because the dairy products in the US diet (51% lower in the US, -661 L/person and day). Despite the differences in water use efficiency across countries, green water is the most important component of the WF_{diet} in both countries. These results are in accordance with other studies that concluded the dominance of green water in food production [10,12,14,17,19,28].

The US is also more efficient in the use of blue water in food production. Our results showed that shifting towards a Mediterranean diet in the US is 15% lower than the blue WF of the Mediterranean diet when consumed in Spain. Also, adopting a US diet in the US would afford a reduction of blue WF of 54% in comparison with the adoption of the same diet in Spain. These differences in blue WF are due to the higher efficiency of the US in the production of dairy products and legumes and nuts.

Despite the comparative advantages of the US in terms of blue and green water use in the production of different food items, it is less competitive in terms of grey water use. For both diets, the grey component of the WF_{diet} is up to 67% higher in the US in comparison with Spain. Legumes and nuts are the food products that show the largest differences in terms of grey water among the two countries. These results are in accordance with those authors that reported high levels of nitrate concentrations in the US farming areas under irrigation, or where the use of nitrogen fertilizers have increased significantly in the last years [49,50].

4.3. Data Limitation and Further Research: Water Footprint of Every Product within the Diet

The calculations of the WF_{diet} in this research are based on global WF datasets for crop and livestock products [41,42]. Previous research has found that WF estimations for the same products and countries can differ significantly [17,19,20]. The specific local climate and agricultural practices have a large effect on the WF of products.

Results of this research showed that olive oil is the product accounting for a large share of the Mediterranean WF_{diet} in both countries. According to Mekonnen and Hoekstra [41], the WF of olive oil produced in Spain accounts for 12.1 m³/L and 4.7 m³/L in the US. Salmoral et al. [19], conducted a detailed study on the WF of olive oil across the five most important production centers in Spain and found large variations, ranging between 13.1 m³/L and 22.4 m³/L [19].

Olive has been cultivated for millennia in the Mediterranean region, and has been considered by many authors a symbol of environmental sustainable systems [51–53]. These studies also described olive cultivation as a low-intensity production system, and usually associated with old trees, small yields, and receiving low inputs for both labor and materials [53]. These factors (mainly low yields) probably explain the large WF (particularly green water) of olives and olive oil. In view of the above, further research is needed to evaluate the high green WFs.

Other products that have a large influence in the WF_{diet} are dairy and meat. De Miguel et al. [17], studied the water footprint of the Spanish pork industry and obtained lower values than those estimated by Mekonnen and Hoekstra [42] (6094 L/kg versus 7184.3 L/kg respectively). The wide range of WF estimations for the same products and countries requires a detailed comparison of the values reported by Mekonnen and Hoekstra [41,42]. Also, more national and local studies are needed. This also requires further research on the calculation of products' WFs at national and more local scales.

5. Conclusions

Few studies have evaluated the WF of real recommended and daily menus, using traditional and national-local recipes and dishes with individual products analysis. As this study has shown, changing consumption patterns towards recommended diets based on a high intake of vegetables, fruits and fish would deliver significant water savings, in some cases larger than those associated to increasing

efficient production. But this message is less likely to be embraced by the general public because of the lack of knowledge about the environmental impacts of current consumption patterns and in particular linked to the diets [54,55]. This study demonstrates how important diets are for consumers and the environment, and supports other studies which argue that diets do matter when referring to sustainability [45,56]. Also, this paper highlights the benefits linked to embracing the Mediterranean diet not just because of its potential health benefits, but also because it is a less water intensive diet. As shown, adopting a Mediterranean diet would lead to major water savings in Spain and also in the US. Therefore, our findings support the conclusion that diets based on low meat consumption could also be more environmentally sustainable in terms of water savings, contributing to address the health-environment problem [8].

A further conclusion of this research is that the origin of the products also matter. More water savings can be achieved when efficient production systems coexist with sustainable consumption patterns. Further research is required to assess the sustainability of diets, e.g., through LCA impact [57] or by conducting a sustainability WFA [15], since the WF of diets only provides insight on the amount of water embedded in food production, regardless of the impacts such water consumption generates in the production regions.

The largest share of the WF of the two diets analyzed in this paper is always linked to green and blue water. Nevertheless, grey WF is considerably larger in the US, mainly because of larger nitrogen pollution in water resources caused in the production of legumes and nuts.

A few products have a large influence and account for the major part of the WF of a diet in both countries and dietary options. Meat, oils, vegetable fats and dairy products are the most influential and important ones. In the USDAR diet the product which accounts the most for the final WFs values is semi-skimmed milk in Spain, and beef meat in the US. Olive oil is the product which contributes the largest percentage of water footprint in the Mediterranean diet in Spain and the US. Despite olives being millennial, local and landscape-adapted trees, their high green WF values make olive oil one of the major water consuming products, even more than meat and dairy. Further research is thus required to assess the WF of the food components that influence the WF of diets the most, in order to obtain more accurate estimates.

Supplementary Materials: The following are available online at www.mdpi.com/2073-4441/8/10/448/s1, Table S1: Recipes, dishes and products' quantity of winter menu (menu 1) of Mediterranean diet, Table S2: Recipes, dishes and products' quantity of summer menu (menu 2) of Mediterranean diet, Table S3: Recipes, dishes and products' quantity of menu 1 of USDAR diet, Table S4: Recipes, dishes and products' quantity of menu 2 of USDAR diet.

Acknowledgments: This research has been funded by the the Water Observatory of the Botin Foundation. The authors are grateful to Michelaina Jhonson for her support in the collection of information and to Maite M. Aldaya for providing valuable comments during the design phase of this research.

Author Contributions: Alejandro Blas, Alberto Garrido and Bárbara A. Willaarts conceived and designed the experiments; Alejandro Blas performed the experiments and analyzed the data; Alejandro Blas and Bárbara A. Willaarts wrote the paper; Alejandro Blas, Alberto Garrido and Bárbara A. Willaarts contributed to the analysis tools; Alejandro Blas, Alberto Garrido and Bárbara Willaarts contributed to the discussion; Alberto Garrido and Bárbara A. Willaarts supervised the experimental design.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Godfray, H.C.J.; Beddington, J.R.; Crute, I.R.; Haddad, L.; Lawrence, D.; Muir, J.F.; Pretty, J.; Robinson, S.; Thomas, S.M.; Toulmin, C. Food Security: The Challenge of Feeding 9 Billion People. *Science* **2010**, *327*, 812–818. [[CrossRef](#)] [[PubMed](#)]
2. Flachsbarth, I.; Willaarts, B.; Xie, H.; Pitois, G.; Mueller, N.D.; Ringler, C.; Garrido, A. The Role of Latin America's Land and Water Resources for Global Food Security: Environmental Trade-Offs of Future Food Production Pathways. *PLoS ONE* **2015**, *10*, e0116733. [[CrossRef](#)] [[PubMed](#)]
3. Matson, P.A.; Parton, W.J.; Power, A.G.; Swift, M.J. Agricultural Intensification and Ecosystem Properties. *Science* **1997**, *277*, 504–509. [[CrossRef](#)] [[PubMed](#)]

4. Tilman, D.; Cassman, K.G.; Matson, P.A.; Naylor, R.; Polasky, S. Agricultural Sustainability and Intensive Production Practices. *Nature* **2002**, *418*, 671–677. [[CrossRef](#)] [[PubMed](#)]
5. Food and Agriculture Organization of the United Nations. Statistical Database. Available online: <http://faostat3.fao.org/home/E> (accessed on 1 January 2015).
6. Tilman, D.; Fargione, J.; Wolff, B.; Antonio, C.D.; Dobson, A.; Howarth, R.; Schindler, D.; Schlesinger, W.H.; Simberloff, D.; Swackhamer, D. Forecasting Agriculturally Driven Environmental Change. *Am. Assoc. Adv. Sci.* **2001**, *292*, 281–284. [[CrossRef](#)] [[PubMed](#)]
7. Carlsson-Kanyama, A.; González, A.D. Potential Contributions of Food Consumption Patterns to Climate Change. *Am. J. Clin. Nutr.* **2009**, *89*, 1704S–1709S. [[CrossRef](#)] [[PubMed](#)]
8. Tilman, D.; Clark, M. Global Diets Link Environmental Sustainability and Human Health. *Nature* **2014**, *515*, 518–522. [[CrossRef](#)] [[PubMed](#)]
9. Jalava, M.; Kumm, M.; Porkka, M.; Siebert, S.; Varis, O. Diet Change—A Solution to Reduce Water Use? *Environ. Res. Lett.* **2014**, *9*. [[CrossRef](#)]
10. Rockström, J.; Falkenmark, M.; Karlberg, L.; Hoff, H.; Rost, S.; Gerten, D. Future Water Availability for Global Food Production: The Potential of Green Water for Increasing Resilience to Global Change. *Water Resour. Res.* **2009**, *45*, 1–16. [[CrossRef](#)]
11. Pimentel, D.; Pimentel, M. Sustainability of Meat-Based and Plant-Based Diets and the Environment. *Am. J. Clin. Nutr.* **2003**, *78*, 660–663.
12. Vanham, D.; Mekonnen, M.M.; Hoekstra, A.Y. The Water Footprint of the EU for Different Diets. *Ecol. Indic.* **2013**, *32*, 1–8. [[CrossRef](#)]
13. Vanham, D.; Hoekstra, A.Y.; Bidoglio, G. Potential Water Saving through Changes in European Diets. *Environ. Int.* **2013**, *61*, 45–56. [[CrossRef](#)] [[PubMed](#)]
14. Vanham, D. The Water Footprint of Austria for Different Diets. *Water Sci. Technol.* **2013**, *67*, 824–830. [[CrossRef](#)] [[PubMed](#)]
15. Hoekstra, A.Y.; Chapagain, A.K.; Aldaya, M.M.; Mekonnen, M.M. *The Water Footprint Assessment Manual*; Earthscan: London, UK, 2011.
16. Boulay, A.-M.M.; Hoekstra, A.Y.; Vionnet, S. Complementarities of Water-Focused Life Cycle Assessment and Water Footprint Assessment. *Environ. Sci. Technol.* **2013**, *47*, 11926–11927. [[CrossRef](#)] [[PubMed](#)]
17. De Miguel, Á.; Hoekstra, A.Y.; García-Calvo, E. Sustainability of the Water Footprint of the Spanish Pork Industry. *Ecol. Indic.* **2015**, *57*, 465–474. [[CrossRef](#)]
18. Chico, D.; Aldaya, M.M.; Garrido, A. A Water Footprint Assessment of a Pair of Jeans: The Influence of Agricultural Policies on the Sustainability of Consumer Products. *J. Clean. Prod.* **2013**, *57*, 238–248. [[CrossRef](#)]
19. Salmoral, G.; Aldaya, M.M.; Chico, D.; Garrido, A.; Llamas, R. The Water Footprint of Olives and Olive Oil in Spain. *Span. J. Agric. Res.* **2011**, *9*, 1089–1104. [[CrossRef](#)]
20. Chico, D.; Salmoral, G.; Llamas, M.R.; Garrido, A.; Aldaya, M.M. *The Water Footprint and Virtual Water Exports of Spanish Tomatoes*; Fundación Botín: Madrid, Spain, 2010; Volume 8.
21. Dumont, A.; Salmoral, G.; Llamas, M.R. The Water Footprint of a River Basin with a Special Focus on Groundwater: The Case of Guadalquivir Basin (Spain). *Water Resour. Ind.* **2013**, *2*, 60–76. [[CrossRef](#)]
22. International Organization for Standardization (ISO). *ISO/DIS 14046.2: Environmental Management—Water Footprint—Principles, Requirements and Guidelines*; ISO: Geneva, Switzerland, 2013.
23. Berger, M.; Finkbeiner, M. Water Footprinting: How to Address Water Use in Life Cycle Assessment? *Sustainability* **2010**, *2*, 919–944. [[CrossRef](#)]
24. Manzardo, A.; Mazzi, A.; Loss, A.; Butler, M.; Williamson, A.; Scipioni, A. Lessons Learned from the Application of Different Water Footprint Approaches to Compare Different Food Packaging Alternatives. *J. Clean. Prod.* **2016**, *112*, 4657–4666. [[CrossRef](#)]
25. Jefferies, D.; Muñoz, I.; Hodges, J.; King, V.J.; Aldaya, M.; Erwin, A.E.; Milà i Canals, L.; Hoekstra, A.Y. Water Footprint and Life Cycle Assessment as Approaches to Assess Potential Impacts of Products on Water Consumption. Key Learning Points from Pilot Studies on Tea and Margarine. *J. Clean. Prod.* **2012**, *33*, 155–166. [[CrossRef](#)]
26. Pacetti, T.; Lombardi, L.; Federici, G. Water-Energy Nexus: A Case of Biogas Production from Energy Crops Evaluated by Water Footprint and LCA Methods. *J. Clean. Prod.* **2015**, *101*, 278–291. [[CrossRef](#)]

27. Milà, L.; Chapagain, A.; Orr, S.; Chenoweth, J.; Anton, A.; Clift, R. Assessing Freshwater Use Impacts in LCA, Part 2: Case Study of Broccoli Production in the UK and Spain. *Int. J. Life Cycle Assess.* **2010**, *5*, 598–607. [[CrossRef](#)]
28. López-Gunn, E.; Willaarts, B.; Dumont, A.; Niemeyer, I.; Martínez-Santos, P. The concept of water and food security in Spain. In *Water, Agriculture and the Environment in Spain: Can We Square the Circle?*; CRC Press: Leiden, The Netherlands, 2012; pp. 21–34.
29. Vanham, D.; Pozo, S.; Pekcan, A.G.; Keinan-boker, L.; Trichopoulou, A.; Gawlik, B.M. Water Consumption Related to Different Diets in Mediterranean Cities. *Sci. Total Environ.* **2016**, *573*, 96–105. [[CrossRef](#)] [[PubMed](#)]
30. Vanham, D.; Mak, T.N.; Gawlik, B.M. Urban Food Consumption and Associated Water Resources: The Example of Dutch Cities. *Sci. Total Environ.* **2016**, *565*, 232–239. [[CrossRef](#)] [[PubMed](#)]
31. Bach-faig, A.; Berry, E.M.; Lairon, D.; Reguant, J.; Trichopoulou, A.; Dernini, S.; Medina, F.X.; Battino, M.; Belahsen, R. Mediterranean Diet Pyramid Today. Science and Cultural Updates. *Public Health Nutr.* **2011**, *14*, 2274–2284. [[CrossRef](#)] [[PubMed](#)]
32. Sáez-Almendros, S.; Obrador, B.; Bach-Faig, A.; Serra-Majem, L. Environmental Footprints of Mediterranean versus Western Dietary Patterns: Beyond the Health Benefits of the Mediterranean Diet. *Environ. Health* **2013**, *2*, 1–8. [[CrossRef](#)] [[PubMed](#)]
33. Bonaccio, M.; Iacoviello, L.; De Gaetano, G. The Mediterranean Diet: The Reasons for a Success. *Thromb. Res.* **2012**, *129*, 401–404. [[PubMed](#)]
34. The United Nations Educational, Scientific and Cultural Organization. Representative List of the Intangible Cultural Heritage of Humanity. Available online: <http://www.unesco.org/culture/ich/en/RL/mediterranean-diet-00884> (accessed on 1 February 2015).
35. Smith, C.; McDonald, G. Assessing the Sustainability of Agriculture at the Planning Stage. *J. Environ. Manag.* **1998**, *52*, 15–37. [[CrossRef](#)]
36. Capone, R.; El Bilali, H.; Debs, P.; Cardone, G.; Driouech, N. Mediterranean Food Consumption Patterns Sustainability: Setting Up a Common Ground for Future Research and Action. *Am. J. Nutr. Food Sci.* **2014**, *1*, 37–52. [[CrossRef](#)]
37. Haven, J.; Rihane, C.; Britten, P.; Johnson-Bailey, D.; Lino, M.; Rahavi, E.; Ciampo, M. Healthy Eating on a Budget at ChooseMyPlate.gov. *J. Acad. Nutr. Diet.* **2015**, *114*, 834–837. [[CrossRef](#)] [[PubMed](#)]
38. United States Department of Agriculture. Choose My Plate. Available online: <http://www.choosemyplate.gov> (accessed on 1 March 2016).
39. The Food and Agriculture Organization Corporate Statistical Database. Food Balance/Food Balance Sheets. Available online: <http://faostat3.fao.org> (accessed on 1 May 2016).
40. Fundación Dieta Mediterránea. Available online: <http://dietamediterranea.com/> (accessed on 1 January 2016).
41. Mekonnen, M.M.; Hoekstra, A.Y. The Green, Blue and Grey Water Footprint of Crops and Derived Crop Products. *Hydrol. Earth Syst. Sci.* **2011**, *15*, 1577–1600. [[CrossRef](#)]
42. Mekonnen, M.M.; Hoekstra, A.Y. A Global Assessment of the Water Footprint of Farm Animal Products. *Ecosystems* **2012**, *15*, 401–415. [[CrossRef](#)]
43. Pahlow, M.; Van Oel, P.R.; Mekonnen, M.M.; Hoekstra, A.Y. Increasing Pressure on Freshwater Resources due to Terrestrial Feed Ingredients for Aquaculture Production. *Sci. Total Environ.* **2015**, *536*, 847–857. [[CrossRef](#)] [[PubMed](#)]
44. Marlow, H.J.; Hayes, W.K.; Soret, S.; Carter, R.L.; Schwab, E.R.; Sabate, J. Diet and the Environment: Does What You Eat Matter? *Am. J. Clin. Nutr.* **2009**, *89*, 1699–1703. [[CrossRef](#)] [[PubMed](#)]
45. Reijnders, L.; Soret, S. Quantification of the Environmental Impact of Different Dietary. *Am. J. Clin. Nutr.* **2003**, *78*, 664–668.
46. Vanham, D.; Bidoglio, G. A Review on the Indicator Water Footprint for the EU28. *Ecol. Indic.* **2015**, *26*, 61–75. [[CrossRef](#)]
47. Olesen, J.E.; Trnka, M.; Kersebaum, K.C.; Skjelvåg, A.O.; Seguin, B.; Peltonen-sainio, P.; Rossi, F.; Kozyra, J.; Micale, F. Impacts and Adaptation of European Crop Production Systems to Climate Change. *Eur. J. Agron. J.* **2011**, *34*, 96–112. [[CrossRef](#)]
48. Wriedt, G.; Van Der Velde, M.; Aloe, A.; Bouraoui, F. Estimating Irrigation Water Requirements in Europe. *J. Hydrol.* **2009**, *373*, 527–544. [[CrossRef](#)]
49. Power, J.F.; Schepers, J.S. Nitrate Contamination of Groundwater in North America. *Agric. Ecosyst. Environ.* **1989**, *26*, 165–187. [[CrossRef](#)]

50. Hallberg, G.R. Pesticides Pollution of Groundwater in the Humid United States. *Agric. Ecosyst. Environ.* **1989**, *26*, 299–367. [[CrossRef](#)]
51. Duarte, F.; Jones, N.; Fleskens, L. Traditional Olive Orchards on Sloping Land: Sustainability or Abandonment? *J. Environ. Manag.* **2008**, *89*, 86–98. [[CrossRef](#)] [[PubMed](#)]
52. Loumou, A.; Giourga, C. Olive groves: “The Life and Identity of the Mediterranean”. *Agric. Hum. Values* **2003**, *20*, 87–95. [[CrossRef](#)]
53. Blondel, J. The “Design” of Mediterranean Landscapes: A Millennial Story of Humans and Ecological Systems during the Historic Period. *Hum. Ecol.* **2006**, *34*, 713–729. [[CrossRef](#)]
54. Gephart, J.A.; Davis, K.F.; Emery, K.A.; Leach, A.M.; Galloway, J.N.; Pace, M.L. The Environmental Cost of Subsistence: Optimizing Diets to Minimize Footprints. *Sci. Total Environ.* **2016**, *553*, 120–127. [[CrossRef](#)] [[PubMed](#)]
55. Tobler, C.; Visschers, V.H.M.; Siegrist, M. Eating Green. Consumers’ Willingness to Adopt Ecological Food Consumption. *Appetite* **2011**, *57*, 674–682. [[CrossRef](#)] [[PubMed](#)]
56. Goodland, R. Environmental Sustainability in Agriculture: Diet Matters. *Ecol. Econ.* **1997**, *23*, 189–200. [[CrossRef](#)]
57. Kounina, A.; Margni, M.; Bayart, J.-B.; Boulay, A.-M.; Berger, M.; Bulle, C.; Frischknecht, R.; Koehler, A.; Milà i Canals, L.; Motoshita, M.; et al. Review of Methods Addressing Freshwater Use in Life Cycle Inventory and Impact Assessment. *Int. J. Life Cycle Assess.* **2013**, *18*, 701–721. [[CrossRef](#)]



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).