

Title: Creating of a xylarium and herbarium with a preventive conservation profile: review of procedures and materials

Author(s): Silva, JL (Silva, Jose Luis); Bordalo, R (Bordalo, Rui); Pissarra, J (Pissarra, Jose); de Palacios, P (de Palacios, Paloma); Vieira, C (Vieira, Cristiana); Vieira, E (Vieira, Eduarda)

Source: MUSEUM MANAGEMENT AND **DOI:** 10.1080/09647775.2024.2375989

Creating xylarium and herbarium with a preventive conservation profile: review of procedures and materials

José Luís Silva¹, Rui Bordalo¹, José Pissarra², Paloma de Palacios³, Cristiana Vieira⁴, Eduarda Vieira¹

[1] Universidade Católica Portuguesa, School of Arts, Research Center for the Science and Technology of the Arts, Portugal

[2] Green UPorto and Department of Biology - Faculty of Sciences, University of Porto, Portugal

[3] School of Forestry and Natural Environment Engineering, Universidad Politécnica de Madrid, Spain

[4] Museu de História Natural e da Ciência da Universidade do Porto (MHNC-UP/UPorto/PRISC), Praça Gomes Teixeira, 4099-002 Porto, Portugal

Abstract

The significance of plant material sample deposits is increasingly apparent beyond the scientific community. Their frequent usage by various industries, history-related researchers, and law enforcement agencies, for example, justifies its growing importance and necessity. Therefore, maintaining the existing xylarium and herbarium worldwide offers the opportunity to access centuries of data, species, and samples no longer present in their original locations or, in some cases, extinct altogether. However, while the preservation of these plant material archives is important, it is equally significant to ensure the proper collection, structure, organization, and classification of the xylarium and herbarium currently being built.

If these archives are established with a preventive conservation perspective from the outset, which avoids degradative factors beforehand, and plans all actions, materials, and uses so as not to endanger future research or the application of analytical techniques that may yet to exist, this will ensure future accessibility to these samples and, consequently, all the knowledge and possibilities they hold. With the aim of developing a xylarium and an herbarium, this paper examines crucial factors including sample drying and processing, primary pests, preventative and curative measures, as well as archival materials, facilities, and handling procedures.

Conteúdo

Abstract 1

33	Introduction	3
34	1. A vouchered xylarium for the 21 st century	3
35	2. Xylarium reference collections.....	5
36	3. Herbarium reference Collections	5
37	4. Photographic documentation	6
38	5. Sample labelling	7
39	6. Colour charting.....	8
40	7. Drying and pressing process for xylarium and herbarium	8
41	8. Mounting specimens.....	10
42	9. Label template and archival material.....	11
43	10. Main pests of xylariums and herbarium	12
44	11. Conservation methods for xylarium and herbarium.....	13
45	11.1. Fumigation of the collection	13
46	11.2. Modified Atmosphere Treatment (Anoxia).....	14
47	11.3. Microwaves and ultraviolet radiation.....	14
48	11.4. Temperature control.....	15
49	11.5. Freezing samples.....	15
50	11.6. Vacuum chambers.....	16
51	11.7. Regular inspections	16
52	11.8. Continuous monitoring devices	16
53	12. Storage, Archiving and Handling of xylarium and herbarium Samples.....	17
54	12.1. Storage facilities	18
55	12.2. Archival System	19
56	12.3. Handling procedures	19
57	13. Conclusion	20
58	Acknowledgements.....	21
59	References.....	22
60		
61		
62		

Introduction

The notion of wood collections initially emerged in central Europe during the late 18th century. This practice involved the creation of wooden libraries, which frequently consisted of boxes crafted from the wood of relevant tree species. These containers also held other components of the selected species, such as dried flowers and leaves (Feuchter-Schawelka & Freitag, 2001). Wood collections, intended for anatomical studies, may have been originated in the mid-19th century, as for example, the Royal Botanic Gardens, Kew, which began collecting wood specimens with a scientific methodology since 1847. Subsequently, numerous xylaria (plural of xylarium) were established around the world, with their peak of development occurring during the 1940s and 1950s (Wiedenhoeft, 2014).

In many instances, these collections were little more than a accumulation of wood and herbarium samples. Due to the absence of preventive measures, they soon displayed a subpar state of conservation, ultimately leading to vigorous and, in certain cases, irreversible measures that restricted their scientific utility and usage.

Nowadays, preventive conservation is no longer optional, but a mandatory approach. Although curative conservation and restoration, and their before and after conditions, capture the public's attention, it is the invisible preventive approach that yields the most satisfactory long-term outcomes for cultural heritage. Preventive conservation aims to prevent damage from occurring by conducting research, planning, and conserving. To maintain, instead of restoring, is the pivotal statement here and is being implemented across various sectors including society, economy, environment, and cultural heritage, as previously mentioned.

The Michalski et al. (2016) ABC method is widely accepted and implemented as a guideline for holistic consideration of collections housed within buildings. This method takes a macroscopic approach, considering the building's integral relationship with its environment, as well as microscopic details such as the building's implementations and impacts. Cultural heritage cannot be viewed as a distinct and self-contained aspect of human knowledge. Nowadays, it is increasingly regarded as a dynamic platform for sharing knowledge and expertise. As such, it absorbs and disseminates valuable experience and know-how.

Focusing on the environment and heritage sectors, we evaluate the construction measures for xylarium and its herbarium. We adopt a preventive conservation approach and establish foundations to incorporate future investigations, techniques, and methods. The collection is maintained in a climate-controlled and environmentally friendly situation to prevent damage. This approach ensures that stored samples remain available and useful in the future, enabling the preservation or enhancement of their physical and chemical characteristics and profiles over time.

1. A vouchered xylarium for the 21st century

A collection of wood samples offers a consistent and manageable resource for analysing, comparing and studying wood that enables investigators to explore more research areas besides wood identification.

The standout feature of a xylarium created in the 21st century is its potential to be an invaluable resource for future investigations, resources, and technologies that are yet to be discovered. In order for any future xylarium to be scientifically confirmed and retested at any time, it is paramount that vouchered samples are associated with an herbarium for the scientific

verifiability of the sample preserved. This statement extends beyond mere scientific accreditation but guarantees the potential for future confirmation and re-examination of the current samples, facilitating continuous adaptation to future identity checks, taxonomic changes, and discoveries (Barker & Flinders, 2005; Ross, 1973 cited in Greve et al., 2016).

Recent research (Bebber et al., 2010; Schmitz et al., 2019) has identified previously unclassified specimens, utilizing innovative genetic technologies (Beck & Semple, 2015; Jiao et al., 2018; Yu et al., 2017), and mapping biodiversity changes over time in response to global alterations (IPCC, 2014). It is important to maintain objectivity, clarity, and precision of language, avoiding subjectivity and biased language. It is also essential to adhere to conventional academic writing standards, including regular formatting, clear structure, and precise word choice, free from grammatical errors and spelling mistakes.

The correlation of wood samples to other parts of the specimen, including leaves, flowers, fruits, and bark, holds significant importance. Therefore, the concurrent establishment of an herbarium is vital (D. Bridson & L. Forman, 2010; Schmitz et al., 2019). These herbaria form a vast and intricate source of information that can have applications beyond their original purpose (British Columbia Ministry of Forests, 1996; Drobniak, 2008; Greve et al., 2016; Pyke & Ehrlich, 2010). The scientific value attributed to herbarium records is increasingly significant as a source of information on global biodiversity, as highlighted by organizations like the IUCN Red List of Threatened Species (IUCN, 2022) and IPBES - Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES, n.d.) (Greve et al., 2016). As evidence of its increasing relevance, online access to herbarium is becoming more accessible through shared portals such as the Global Biodiversity Information Facility (GBIF, 2021).

Even if not affiliated with an xylarium, the range of research areas covered by an herbarium is extensive. These include confirming the identity of specimens or discovering new variants (taxonomy); providing material for morphological measurements (systematics); verifying scientific names (nomenclature); supplying material for DNA analysis (genetics) (Edinburgh, 2022); and for geographic information system studies (ecology) (Staudinger et al., 2012), or contemporary specimens for comparison with fossils in paleobotany. It can also provide access to pollen for taxonomic, systematic, and pollination studies, as well as for allergy research in insect ecology, pollination ecology, and medical studies. It also documents temporal changes in the phenological traits of species (Staudinger et al., 2012). It enables microscopic examination for anatomy and morphology, along with chemical analysis for lead-uptake, pollution documentation, bio-prospecting, and chemical concentration. Additionally, information on common names and local uses of plants was gathered for anthropology, linguistics, ethnobotany, and economic botany purposes, while the geographic distribution of species was determined according to Pyke and Ehrlich (2010). Temporal changes in the environment were monitored (Greve et al., 2016) and documentation on minor climate cycles (paleoecology) or providing carbon isotope ratios (for example, Lewis and Clark specimens from 200 years ago have increased C12) showed evidence of climate change. The institution serves as a repository for voucher specimens globally, for use in fields such as ecology, ethnobotany, and environmental impact studies (Funk, 2007).

Continuous support for xylarium and herbarium is of utmost importance to ensure a consistent supply of specimens that can be used for various scientific purposes (Greve et al., 2016). The inclusion of historical wood and herbarium samples, sourced from art objects, historical structures, and archaeological sites, can provide valuable information to a range of fields, including Dendrochronology, Paleobotany, conservation & restoration, the history of

commerce, environmental biology, and potential areas of research not yet identified. All these factors will facilitate the establishment of a xylarium, which is capable of meeting the increasingly frequent and specific requirements of the industry, judiciary, art history, archaeology, and others (Beeckman, 2003; Carlquist, 1982; Cartwright, 2015; Collins & Cruickshank, 2012; Schmitz et al., 2019).

2. Xylarium reference collections

Constructing a well-grounded xylarium and collecting its samples is a complex undertaking. This process requires significant time, physical effort, and financial investment and should not be expected to be flawless, but rather aimed at achieving the highest attainable standards (Schmitz et al., 2019). When striving for a scientifically rigorous xylarium, Table 1 provides relevant factors to consider.

Table 1 – Protocol for the construction of a xylarium. (British Columbia Ministry of Forests, 1996; D. Bridson & L. Forman, 2010; Perkins, 2022; RHS - Royal Horticultural Society, 2013; Schmitz et al., 2019, 2020; Wiedenhoef, 2014).

I.	It is advisable to take note of the field trip season, particularly when collecting deciduous species, as this may impede the collection of leaves, flowers, and fruits intended for the herbarium.
II.	Ideally, a botanist ought to be present to identify the specimen before collection.
III.	Use a GPS to obtain and record latitude and longitude coordinates (in decimal degrees with at least four decimals) when collecting samples. Ensure the location data is in the WGS84 datum to accurately pinpoint the plant collection location. Alternatively, utilise a GPS Logger app on a smartphone.
IV.	When feasible, collect a sample that encompasses the bark, sapwood, and heartwood.
V.	Ideally, procure a sample (a disc) from the entire trunk. When unattainable, an wedge-shaped piece that stretches from the centre of the trunk to the bark is ideal. However, if not possible either, cut a sample that measures more than 5 cm wide, about 20 cm tall, and if possible, up to the centre of the specimen.
VI.	Ensure to record the location of sample collection within the specimen at ground height, chest height, or branch.
VII.	Collect multiple samples from the same specimen, if possible, for analysis of internal variations and as a backup for lost or destroyed material or future trade with other xylarium.
VIII.	Ideally, choose trees without cracks or any visible signs of infestation or infection.
IX.	Avoid using chemicals or pencil marks on the sample during processing as this may affect future stable isotopes, DART, and NIRS analyses.
X.	When possible, participate or be present during sample collection. If not possible, work only with samples that have attached information such as species, variety, and geographic origin.
XI.	Preferably, use xylarium linked to herbarium vouchers when working with donated samples from other collections. If not possible, have the samples identified by an experienced wood anatomist to maintain reliability.

A worldwide index of xilaria is available in the Index Xylariorum 4.1 that was compiled by Anna H. Lynch and Peter E. Gasson (version March 2010) and updated by IAWA under supervision of Frederic Lens (March 2016) - (Lynch et al., 2016).

3. Herbarium reference Collections

Collecting plant specimens to create an herbarium is a precise task that requires certainty that the collected and processed specimen matches the specified species. To ensure scientific credibility and value, confirmation from a taxonomist is necessary. It is essential to consider factors such as the sample's original location and environment typology when conducting research (British Columbia Ministry of Forests, 1996).

Although the procedure for collecting xylarium and herbarium specimens are related, they are not the same. Therefore, Table 2 provides additional relevant points to the previous information.

Table 2 - Protocol for constructing an herbarium (British Columbia Ministry of Forests, 1996; Perkins, 2022; RHS - Royal Horticultural Society, 2013; Schmitz et al., 2019, 2020).

I.	Note the field trip season, as this may impede the collection of leaves, flowers, and fruits, particularly when targeting deciduous species.
II.	Ideally, a botanist should be present for specimen identification before collection.
III.	Use a GPS to acquire latitude and longitude coordinates (in decimal degrees with at least four decimals) and employ WGS84 as the datum to locate the plant collection spot. Alternatively, use a GPS Logger app if in possession of a smartphone.
IV.	Handpick specimens and samples with good condition and without any signs of animal/insect damage, deformities or diseases.
V.	Whenever possible, include all plant parts that may help with identification. Choose different plant parts that enable identification and not just the biggest leaf or flower.
VI.	It is advisable to collect several samples from the same specimen to allow for the analysis of internal variations and the use of other techniques, as well as providing a backup for lost or destroyed material or future trade with other herbarium.
VII.	A polythene bag should be used to hold the specimen, along with an identifier. Ensure that only one bag is used for each specimen. Any excess air in the bag should be expelled before sealing it.
VIII.	To maintain the specimens inside the bag during collection, it is advised to avoid direct sunlight and hot places, and where possible, store them in a cooler. If this is not feasible, sprinkle water inside the bag or add a damp paper towel.
IX.	Avoid collecting fallen leaves from the ground, young trees or sprouts as their morphological characteristics may vary from those of mature trees.
X.	It is recommended to avoid using chemicals or pencil marks on the samples during processing as it may affect stable isotopes, DART, and NIRS analyses.
XI.	In cases where it is not feasible to be present during collection, it is advisable to work only with samples that have attached information, such as species, variety, and geographic origin.
XII.	If it is not possible to have samples with attached information, it is suggested to get the samples identified by an experienced botanist.

A worldwide index of herbaria is available in the Index Herbarium, which was originally published by the International Association for Plant Taxonomy, and subsequently maintained by the New York Botanical Garden - (New Your Botanical Garden, n.d.).

4. Photographic documentation

Wherever feasible, photographic documentation of the sample, surroundings, process, and materials collected should be taken for future reference. This is a vital and valuable source of information. In contrast to previous decades, it is now an essential, cost-effective, and speedy procedure due to digital photography. The protocol specified in Table 3 applies to both xylarium and herbarium specimens.

Table 3 - Procedure for photographic documentation of xylarium and herbarium samples. (Schmitz et al., 2019).

I.	When feasible, employ a standard-sized object (e.g., ruler, coin, or scale bar).
II.	Capture an image of the overall shape and the surroundings of the specimen.
III.	Obtain general images of the bark, leaves, flowers, fruits, thorns, and any other relevant details.
IV.	Take close-up photographs of any notable characteristics such as diseases, malformations, or fissures.

5. Sample labelling

There are several different methods for labelling samples, especially in the field. In this case, we used several authors as references, with particular attention to Schmitz et al. (2019). Table 4 provides information applicable to both xylarium and herbarium collections.

Table 4 - Protocol for xylarium and herbarium field description and specimen labelling. (British Columbia Ministry of Forests, 1996; D. Bridson & L. Forman, 2010; Perkins, 2022; RHS - Royal Horticultural Society, 2013; Schmitz et al., 2019, 2020).

Wherever possible, the identifier should register:	
I.	Taxonomic classification (confirmed/probable).
II.	The number of samples collected.
III.	The full date of collection.
IV.	Season of collection.
V.	Name(s) of collector(s).
VI.	Name(s) of region/city/place of collection and GPS location, if possible.
VII.	Type of material collected (wood, leaves, flowers, fruits, cones, bark or other).
VIII.	The location of collection of the wood sample within the specimen (at ground level, at breast height, or on a branch), in case of a xylarium.
IX.	The circumference or diameter (in centimetres) of the xylarium specimens at breast height. The specimen should be standing or recently felled.
X.	The height of the specimen (in metres) should be estimated using a clinometer and laser meter/tape measure or a hypsometer.
XI.	Ensure that all samples taken (bark, cambium, wood, leaf, flower, and fruit) from the same specimen are labelled with a distinct and consistent sample code.
XII.	Do not repeat the same number, even if the collection site is the same, the plant was collected at a later date, or it was grown in a greenhouse.
XIII.	Canopy shape, growth shape, and height (spreading, scraggly, columnar) must be described.
XIV.	The habitat description, including slope, riparian area, associated vegetation, and soil type, is also necessary.
XV.	The concise description of the specimen including physical attributes, dimensions, and any other pertinent details.

XVI. The coordinates using a GPS mobile phone application where feasible to accurately determine the plant collection site.

6. Colour charting

The sample colour charting can be a valuable identification factor to consider in certain situations. In this order, some relevant factors are underlined in Table 5.

Table 5 - Indicators for colour charting. Adapted from (British Columbia Ministry of Forests, 1996; RHS - Royal Horticultural Society, 2013).

I.	Compare the colour of the fresh material with a plant indicator colour chart (e.g., Royal Horticultural Society colour chart).
II.	When the colour of the sample is as close as possible to the colour on the chart, add the colour names or numbers to the identifier in the bag.
III.	Colour change or loss is inevitable. However, some samples may return to their original colour as the sample ages or dries.
IV.	Avoid using the chart in direct sunlight or artificial light, the light from a north facing window is ideal.
V.	Be aware that you may need to record more than one colour in different parts of a specimen.
VI.	Many cultivars only show differences in colour characteristics.

7. Drying and pressing process for xylarium and herbarium

The drying time depends, among other factors, on the thickness of the sample, the species density, and the drying conditions. For a piece of wood to reach a stable level of dryness during air drying (with normal air circulation) in a stable (without drastic fluctuations in relative humidity) and protected environment (e.g., without exposure to sunlight) usually takes from a few months to several years. This process can be accelerated by kiln drying. Specific kiln schedules have been developed to control temperature and relative humidity in accordance with the moisture content and stress situation within the wood, thus minimizing shrinkage-caused defects (Boone and others 1988). Moisture content below 20% when introduced into a xylarium, shows minimal susceptibility to biological attacks, few chemical changes over time, and acceptable dimensional changes (Wiedenhoef, 2014).

In the context of herbarium specimen handling, the main objective is to avoid curling, wrinkling, and warping of specimen leaves, flowers, and thin twigs during drying. Although there is some tolerance (even though not advisable) regarding the dryness of xylarium specimens during storage, the complete dryness of herbarium specimens is the most important factor in storage, which is why the following steps are of paramount importance. The materials listed in Table 6 are only indicative and may be replaced by products with similar characteristics or which provide comparable results.

Table 6 – Materials for herbarium samples pressing process.

I.	Pressing boards - consist of a pair of 6mm thick plywood or hardboard sections cut to around 300mm wide x 450 long.
II.	Blotting paper - acid-free 300 gsm blotting paper is the most recommended material for drying out specimens quickly, which also gives the best results in what preservation of colour and other characteristics is concerned.

III.	Corrugated card - Lengths of 450mm minimum is required.
IV.	Foam sheet - sponge/foam sheets cut to format size.
V.	Straps - two 25mm wide x 1.5m long canvas luggage straps with sliding plastic buckles.
VI.	Tensioning bars - wood bars 30mm x 12mm x 460mm long, bevelled at one end to facilitate pushing under the straps.

The protocol for pressing herbarium samples follows a series of steps that, although not rigid, must be followed as closely as possible to obtain similar results. To that end, the process is exposed in Table 7.

Table 7 - Protocol for pressing herbarium samples (British Columbia Ministry of Forests, 1996; Heath et al., 2017; RHS - Royal Horticultural Society, 2013; Vieira, 2023).

I.	Place two pieces of corrugated cardboard (corrugation facing corrugation) over a press board.
II.	Place a double layer of blotting paper on top.
III.	Place your specimen, trying to maintain the natural position of the plant, but making sure that at least one leaf and one flower are visible underneath. If it is not possible to turn the specimen while it is still attached to the plant, remove it, and fix it separately.
IV.	Specimens with long leaves or stems can be folded to fit the given format.
V.	Once the layout has been decided and the appropriate taxonomic classification has been determined, place another double layer of blotting paper over it.
VI.	Place two more pieces of corrugated cardboard (corrugation facing corrugation) on top of the ensemble, followed by another pressing board. If another specimen is to be made, the same procedure is followed, with the press board placed as the last layer before pressing.
VII.	If the specimen is bulky, a sheet of foam should be placed on top of the second set of blotting papers and corrugated board.
VIII.	When all the layers are in place, the second pressing board is placed on top.
IX.	Pressure can be applied by wrapping the straps around the entire assembly, placing them 1/3 from the end of each press board and tying a knot as tightly as possible. Once this is done, place the tensioning rods under the strap at right angles to it and pull them as far as possible towards the side edge of the boards to increase the pressure.

The aim is to dry the specimen as quickly as possible to retain maximum colour and avoid the formation of mould. To this end, some form of low heat source is possible but not essential (Drobnik, 2008), although it is advisable never to exceed 50 °C (120 °F) and never to use an oven or microwave for this process (Perkins, 2022; Wilken, 2009). Table 8 shows the steps to be taken once the drying process has begun.

Table 8 – Drying process protocol (British Columbia Ministry of Forests, 1996; Perkins, 2022; RHS - Royal Horticultural Society, 2013; Vieira, 2023).

I.	An inspection should be made after 24 hours. Care should be taken when handling the blotting paper to avoid tearing the specimen, which is likely to be in a limp state.
II.	If any of the leaves or petals are accidentally folded, they can be placed in the correct position and temporarily fixed with a post-it note.
III.	Ideally, if the blotting paper and corrugated cardboard are found to be damp, they should be replaced with dry ones, but if they are not damp or show no signs of mould,

	it's more common to keep the same materials until the whole ensemble is completely dry.
IV.	Make sure that the corrugated cardboard is not crushed and that there is room for air to circulate.
V.	Tighten the strap daily and allow the whole ensemble to dry for at least a week.
VI.	It can be difficult to conclude that a specimen is completely dry, but if the specimen lifts off the paper easily, is stiff and light to the touch, it should be ready for storage. If in doubt, continue the drying process for a longer period.

8. Mounting specimens

Methods of preparing, mounting, and storing herbarium specimens vary widely between herbarium, depending primarily on the type of plant specimen being processed (Perkins, 2022; Wilken, 2009). While most plant parts can be easily stored and pressed, bulbs, fruits, cones, bark or even large diameter woody stems may need to be cut or dried in boxes or paper bags and stored in acid-free boxes, depending on space requirements (D. Bridson & L. Forman, 2010; Perkins, 2022; Wilken, 2009). Despite the different options in the bibliography consulted, Table 9 summarises the most frequently mentioned materials used.

Table 9 - Materials used for mounting herbarium specimens.

I.	Acid-free or PH-neutral mounting card. Colour and size are optional, but usually white and around A3 size (297 x 420 mm).
II.	Acid-free adhesive, ideally non-water soluble (Drobnik, 2008). Wilken (2009) suggests the reversible adhesive grade A methylcellulose with a molecular weight of 4000, and Cabral (2018) and Heath (2017) suggest a white water-soluble adhesive with a neutral pH. The use of poly(vinyl acetate) emulsion adhesives is discouraged as these tend to become acidic as they age (Down et al, 1996 cited in Wilken, 2009).
III.	Gummed linen hinging tape.
IV.	Transparent polypropylene covers (Cabral & Sales, 2018).

The protocol for mounting specimens also varies between authors and is summarised in Table 10.

Table 10 - Protocol for mounting herbarium specimens.

I.	Place your cardboard label template on the right-hand corner of the mounting sheet.
II.	Fragments such as seeds, petals, leaves or others, should be preserved. For that, they should be placed in a fragment pouch. Usually, it is placed in the bottom left-hand corner.
III.	Fragment pouches can be made of the same acid-free mounting card by folding an A4 sheet into an A6 or A7 size (D. Bridson & L. Forman, 2010; Heath et al., 2017; Perkins, 2022; RHS - Royal Horticultural Society, 2013).
IV.	The specimen may be fixed to the mounting card resorting to small dabs of glue on its underside, which can also be done with a cocktail stick or dissecting needle dipped in glue for finer work (RHS - Royal Horticultural Society, 2013). On the other hand, small strips of gummed linen hinging tape can be glued to the card while trapping the specimen underneath. In this way, the specimen is always free of glue and because of that, manageable. Bigger or more abundant strips of gummed tape should be used on the stalk and flower stems.
V.	To guarantee the fixation of the glue in the desired place, use small sandbags to press it down.

VI.	As extra protection, the entire ensemble may be placed inside a polypropylene transparent cover (Cabral & Sales, 2018).
VII.	In addition to the procedure referred, a more dated and less frequently used method is stitching. In this case, parts of the specimen, usually the bigger and bulbous ones can be stitched to the paper sheet instead of glued. It is mainly used in stems, bulbs, or fruits (Cabral & Sales, 2018; Heath et al., 2017; Perkins, 2022).

9. Label template and archival material

The label on xylarium and herbarium specimens should contain all the essential information and be as durable as possible to avoid unidentified specimens. Loss of this element is one of the most damaging factors for the scientific validity of old xylarium and herbarium. The longevity of labels, particularly handwritten but also printed, is limited, mainly due to the chemical nature of these materials, which results in a relatively rapid degradation that can occur as little as 3 years after the information was written.

The degradation of the material is mainly visible as a change in the original colour (black ballpoint pens and inks become brownish-grey, while blue ones tend to become reddish, pinkish or very pale). The disappearance of the colour in the paper is also a common occurrence, resulting in the appearance of an increasingly wider band around all characters, often leading to the complete disappearance of the written message (Drobnik, 2008). The same applies to low quality inks for computer inkjet printers, as most of them are not water resistant. Of course, paper, as a highly absorbent material, favours the reaction of these materials with water and accelerates their rapid degradation (Drobnik, 2008).

Another factor to be taken into account is the chemical composition of the ink, since, despite the colourfastness of some brands, many of them contain acidic components and microelements such as iron (Fe) and chromium (Cr), which allow the acidification of the paper with the decomposition of the cellulose present in it, but also promote the development of bacteria and microscopic fungi with the transfer of these ions to these microorganisms (Szostak-Kotowa, 2001 cited in Drobnik, 2008; Perkins, 2022).

But the material limitations do not end there. The quality of the ink can be improved by purchasing special archival inks or heritage grade pen inks with a special composition, but it is also important to be aware that the adhesives used have a direct impact on the material they are used to affix. In particular, dense liquid adhesives can gradually dissolve some of the ink's constituents, resulting, for example, in the purpling of the white bars on barcode adhesive labels, which reduces contrast and makes the label unrecognisable to barcode scanners.

A proven material is graphite or carbon. These materials have low reactivity, are not water soluble and don't oxidise, resulting in a permanent black material that has been proven to be still present and legible in the Linnaeus samples. The only limiting factor is the ability to be deliberately erased with a rubber (Drobnik, 2008).

A relatively new technique is that of laser printing and xerographic copies (printed or photocopied labels), which, although considered to be a durable material, have yet to be tested. They are made of a solid material that is melted (not dissolved) and consists mainly of black polymers, which reduces the possibility of diffusion in the printing paper as well as in an adhesive layer (Drobnik, 2007).

Since most of the paper produced until the 1990s is acidic, which causes embrittlement and yellowing of the material due to oxidation of the lignin and changes in the structure of the cellulose chains (Zyska, 1995 cited in Drobnik, 2008), it becomes clear that acid-free (pH neutral or alkaline) paper and adhesives are absolutely necessary for the construction of a herbarium (Drobnik, 2008).

A very common adhesive used to fix specimens to the paper sheet is commonly referred to as 'white glue' and is based on PVA - polyvinyl acetate. This widely used water soluble adhesive has as its main disadvantage the fact that it tends to become acidic with age (Down et al., 1996 cited in Wilken, 2009), but more importantly it is susceptible to degradation by filamentous fungi, and more rarely by algae, yeasts, lichens and bacteria (Cappitelli & Sorlini, 2008), especially when the relative humidity in the herbarium room increases, making its use highly controversial among specialists, as the long term effects of this potential substrate for bacteria are still unknown. Nevertheless, the two main methods of mounting herbarium specimens, the Archer method (Archer, 1950) (in which the specimen is fixed to the herbarium sheet by narrow strips of plastic or paper) and the pasting method (in which the entire specimen is glued to the sheet) both use this type of adhesive.

The alternative methods of stitching specimens to the sheet (Cabral & Sales, 2018; Heath et al., 2017; Perkins, 2022) or pinning are more expensive and labour-intensive methods but are still chosen by some herbarium (Drobnik, 2008). Table 11 lists the materials and information that should be present.

Table 11 – Materials and mandatory information items.

Materials	
I.	Archival quality, acid-free labels.
II.	Acid-free adhesive. Wilken (2009) suggests the reversible adhesive Grade A methylcellulose with a molecular weight of 1500.
Mandatory information items	
I.	Item number.
II.	Origin. Ideally with GPS coordinates.
III.	Wild collection vs. breeding.
IV.	Date of collection.
V.	Collector.
VI.	Taxonomic classification of specimen. Should include scientific classification and family name.
VII.	Always indicate if the sheet is part of a larger specimen that had to be mounted on more than one sheet (e.g., 1 of 2; 2 of 3, etc.). If this is not the case, always mark the sheet as 1 of 1.
VIII.	Additional space for relevant information specific to the item.
IX.	The definition and inclusion of a barcode can be a valuable addition to save time and avoid errors (Cabral & Sales, 2018).

10. Main pests of xylaria and herbaria

Pests in a xylarium and their effects vary according to the geographical location of the storage. In addition to the well-known xylophagous insects, fungi and bacteria are also important microorganisms. All these species have a destructive effect on this type of biological material.

The most common xylarium pests found in Southern Europe are *Anobium punctatum* (furniture beetle); *Hylotrupes bajulus* (house longhorn beetle); *Nacerdes malamura* (wharf borer); *Lyctus brunneus* (brown powder post beetle); *Cryptotermes brevis* (powder post termite); *Reticulitermes lucifugus* (subterranean termite); *Coptotermes acinaciformis* (subterranean termite) (Lieutier et al., 2004; Toriti et al., 2021).

Although the same pests can also attack herbarium specimens, there are some other protagonists that have a specific impact on herbarium deposits. Again, depending on the geographical location of the repository, the nature of the pests and their effects vary considerably. For that reason, it is important to a concise knowledge of the main insect species present in your geographic location (D. Bridson & L. Forman, 2010). In addition to the fungi and bacteria already mentioned, beetles, spiders and other insects can also have a destructive effect on this type of biological material. The most common herbarium pests are: *Lepisma saccharinum* (Silverfish); *Psocoptera* order (Booklice); *Lasioderma serricorne* (Cigarette or tobacco beetles); *Trogoderma angustum* (Herbarium Beetle); *Dermestidae* family (Cockchafer) and *Stegobium paniceum* (Drugstore beetles) (British Columbia Ministry of Forests, 1996; Child, 1998; D. Bridson & L. Forman, 2010; Edinburgh, 2022; Wilken, 2009).

11. Conservation methods for xylarium and herbarium

Occasional infestations are almost inevitable in any museum or repository, even when preventive measures are taken. However, when they do occur, curative measures must be taken as soon as possible. Actions such as:

- I. Immediately isolate any objects suspected or found to be infested to prevent the spread of the infestation.
- II. If possible, seal items immediately in polyethylene zip-lock bags.
- III. Thoroughly clean the area where the infested items were located.
- IV. It is important to identify the pest in order to understand its life cycle and behaviour.

With this information, decisions can be made on the best treatment for the object(s), collection or storage type (D. Bridson & L. Forman, 2010; Pinniger, 1998).

Most of the methods, materials and intervention parameters described below apply to both xylarium and herbarium with minor differences. However, it should be borne in mind that the conservation of herbarium is a highly complex area, as the material to be protected is much more sensitive to degradation than, for example, the wood samples that make up a xylarium. Furthermore, herbarium is more sensitive to certain treatments due to the very nature of the material they preserve, whereas xylarium, as a collection of solid wood samples, can withstand more aggressive and simpler treatments.

The most common threats to the permanence of a herbarium are factors such as past infestations, the natural ageing of the paper used for labels, sheets, covers or envelopes, together with the degradation of inks, adhesives and printed texts (Drobnik, 2008), but it is important to stress the high susceptibility of these collections to factors such as fluctuations in environmental humidity and, consequently, uncontrolled drying or fungal attack.

11.1. Fumigation of the collection

One of the most common methods of curing and preventing infestation and infection was the fumigation of collections. Despite its relative success, the main limitation of this method is that the specimens and the paper sheets (in the case of herbarium) often remained toxic to

human handlers for long periods of time. From the mid-18th century to the end of the 20th century, highly toxic chemicals were used in xylarium and herbarium, as shown in Table 12.

Table 12 - Chemicals most commonly used for fumigation of collections (Briggs et al., 1983; D. Bridson & L. Forman, 2010; Hawks, 2001; Hawks et al., 2004; Kurmanow, 2019; U.C., 2008; Whitmore, 1965; Whitten et al., 1999).

• Mercuric chloride - HgCl₂ • Potassium cyanide – KCN • Carbon tetrachloride - CCl₄ • Dichlorodiphenyltrichloroethane (DDT) - C₁₄H₉Cl₅ • Lauryl Pentachlorophenol - C₆HCl₅O	• Arsenic trioxide - As₂O₃ • Carbon disulfide - CS₂ • Naphthalene - C₁₀H₈ • Lindane - C₆H₆Cl₆ • Organofosforado 2,2 diclorovinil dimetilfosfato (DDVP) - C₄H₇Cl₂O₄P
--	---

The absorption of these toxic compounds through the skin and by inhalation is still a real threat today, mainly because their remnants are still present in all parts of old plant sample deposits (Drobnik, 2008). The use of moth balls (naphthalene or paradichlorobenzene) has been discontinued because of their toxicity (Pereira & Hammond, 2001; Pereira & Wolf, 2001), as the use of commercial insecticides such as Raid and Vapona has been described as more dangerous to the user than to the pest (RHS - Royal Horticultural Society, 2013). A 'newer' problem that many of the old plant specimens have, is the fact that most toxic compounds have a deleterious effect on the DNA of the specimens, making their use for genetic research unviable (Drobnik, 2008).

This method is particularly harmful to herbarium users because the materials stored are generally more absorbent than most woods, adding to the absorbent properties of the mounting paper sheets and folders, which remain toxic to human handlers for long periods of time.

Because of the high risks and extreme hazards associated with the use of highly toxic chemicals, other approaches have been developed. This chemical-free perspective, commonly referred to as Integrated Pest Management (IPM), involves the use of multiple techniques to achieve the same result, which is the destruction of the infestation or vector (Edinburgh, 2022).

11.2. Modified Atmosphere Treatment (Anoxia)

This method has been used to replace fumigation methods, which are much more dangerous for humans. Anoxia takes place in an airtight space (metal chambers or portable special plastic bags) where the oxygen level is extremely low (less than 0.1%). The duration of treatment depends on factors such as temperature (at 25 °C or above it can take 2 to 3 weeks, while at 20 °C or below it can take 4 to 5 weeks) or between 30-38° C as mentioned in Bridson & L. Forman (2010) and the size of the objects (nitrogen can be introduced for large objects), but other chemicals such as carbon dioxide and argon are increasingly being used. This method is now widely available commercially and is particularly applicable to most collection typologies because of its portability (Burke, 1999; Pinniger, 1998).

11.3. Microwaves and ultraviolet radiation

Ultraviolet (UV) radiation and microwave heating are much more damaging to herbarium specimens than to xylarium specimens because of the rapid ageing of paper due to cellulose and lignin degradation, which also increases the loss of paper flexibility and the intense drying of

specimens that are often very fragile by nature. In addition, UV light and microwaves damage the DNA of the specimens, making their use for genetic research unviable (Drobnik, 2008).

At present, the most effective, ecological and beneficial measures to control degradation are temperature and humidity control, regular or even permanent freezing of samples and the use of vacuum chambers.

11.4. Temperature and relative humidity control

Temperatures below 13 °C prevent egg hatching and full metamorphosis of all herbarium insects and reduce their motility (Rumball & Pinniger, 2003), while relative humidity (RH) below 55% combined with temperatures below 18 °C stops the growth of moulds (British Columbia Ministry of Forests, 1996; Rumball & Pinniger, 2003). The Royal Botanic Garden Edinburgh (2022) maintains a stable temperature of 20 °C and 50% relative humidity.

11.5. Freezing samples

Sample freezing temperatures and time vary widely between authors. Table 13 summarises these values and time periods.

Table 13 - Freezing temperatures and time periods.

Authors	Degrees (Celsius, °C)	Freezing time
Strang (1992)	set values that range from -18° C to -32 °C	minimum of 72 hours
Rumball & Pinniger (2003)		
Raphael (1994)	-20 °C	7 to 10 days
British Columbia Ministry of Forests (1996)	-20 °C	48 hours
Bridson & L. Forman (2010)	Minimum -18° C, ideally -30 °C	17 hours
Royal Horticultural Society (2013)	No temperature mentioned	36 hours
Royal Botanical Gardens Kew (2017)	-40 °C	4 days
The Royal Botanical Garden of Edinburgh (2022)	-29 °C	5 days

Although less damaging to herbarium than to xylarium, it is important to note that water vapour tends to condense on the surface of the specimen and must evaporate before the specimens are returned to the shelf (Drobnik, 2007). To avoid this, the Royal Horticultural Society (2013) recommends sealing the specimen(s) in a sealed polythene bag prior to freezing, emphasising the importance of keeping the bag sealed during the defrosting period to prevent moisture absorption by the specimen.

It should be noted that the freezing equipment used must be able to reduce the temperature inside the object in less than 24 hours, since, especially in cold climates, if the freezing process is too slow, some insect species may acclimatise and survive the freezing process (Pinniger, 1998).

The parameters for temperatures and exposure times are the same as above, but it should be noted that tightly rolled tapestries and skins (Pinniger, 1998), large volume boxes or large fascicles require a longer freezing time due to their natural insulation against cold (D. Bridson & L. Forman, 2010; Drobnik, 2008). If possible, the temperature in the centre of these large items should be recorded (Pinniger, 1998). Freezing should be a routine annual procedure, or more frequently if pests or the result of pest metabolic activity are detected (RHS - Royal Horticultural Society, 2013).

11.6. Vacuum chambers

Vacuum chambering is considered the most effective method of pest control. This method destroys the protoplasts within the cell wall, thereby terminating any animal, fungal or bacterial life (Drobnik, 2008).

Particularly for newly acquired or loaned specimens, these procedures are mandatory to ensure the continued stability of herbarium specimens. Irrespective of the protection measures taken, all specimens should be regularly inspected for moisture, mould or insect infestation.

11.7. Regular inspections

Regular inspection and cleaning of xylarium and herbarium cabinets, shelves, work areas, floors and files is the most effective and economical way to prevent pest introduction. Particularly in the case of herbarium, the use of a vacuum cleaner is preferred to the use of wet cloths, mops, or similar systems, as these methods introduce moisture into the environment. Detailed information on this point can be found in the National Park Service Museum Handbook, Part I, Chapter 5, Biological Infestations and Chapter 13, Museum Housekeeping (NPS National Park Services., 2000) (British Columbia Ministry of Forests, 1996; Edinburgh, 2022; KSU Herbarium, 2002; Wilken, 2009).

Due to the fragile and less resistant nature of herbarium specimens, regular inspection and routine cleaning of cabinets, shelves, work areas, floors and files is by far the most important, effective, ecological, and economical measure to prevent pest infestation.

11.8. Continuous monitoring devices

As mentioned above, regular inspections are an effective method of control, but constant surveillance is not possible. Therefore, devices such as sticky traps provide a 24/7 defence against these constant enemies of museum objects and buildings.

This method can be used to trap a wide range of pest species, both crawling and flying, from their mobile larval stages to their adult stages. In addition to the capture itself, data can be obtained on their numbers, seasonal cycles and hiding places. However, this tool should not be used as a pest control device, but only as a pest monitor (Child, 1998).

The forms, shapes, sizes and brands are numerous and readily available on the market, being the two main types of traps, suspended tent traps for catching flying insects and sticky cardboard traps for the ground. The latter may sometimes contain a general food attractant. Typically, funnel traps and UV electrocution types don't have a regular presence in museums.

Another mechanism is the use of semiochemicals, which is essentially the use of chemicals produced by a particular species to influence the behaviour of another individual of the same or a different species. In this specific case, the use of female pheromones to attract the male (Guarino et al., 2020). As shown in Table 14, the use of the synthesised version of these pheromones has been developed for specific insect species that affect both xylarium and herbarium.

Table 14 - Insect species for which synthetic pheromones have been developed.

• <i>Anobium punctatum</i> - Furniture beetle	• <i>Tineola bisselliella</i> - Webbing clothes moth
---	--

• <i>Stegobium paniceum</i> - Drug store beetle	• <i>Lasioderma serricorne</i> - Cigarette beetle
---	---

This strategy, particularly when combined with a food attractant (Guarino et al., 2020) and sticky traps, is highly effective, especially when a particular insect species is suspected of becoming a potential hazard. In particular, in enclosed areas, this option is very successful in attracting large numbers of male insects and is therefore a valuable indicator of likely total insect numbers (Child, 1998; Edinburgh, 2022). The placement of these traps should follow the protocol shown in Table 15.

Table 15 - Trap placement protocol.

I.	Traps should be marked with the date of installation.
	Traps should be placed:
-	On the walls, as most insects do not cross open areas.
-	At wall/floor junctions, corners, near doors, ventilation systems or other entry points.
-	In any dark, hidden area that is difficult to access and inspect.
-	In display cases (if possible).
II.	Hanging tent traps should preferably be placed near windows.
III.	Place on windowsills to catch insects that may be attracted by light.
IV.	The Royal Botanic Garden Edinburgh (2022) also uses UV lights on windowsills and under workstations.

The purpose of these traps is to catch insects, but also to identify them. Traps should be inspected regularly, and the insects caught (adults or larvae) should be identified, counted and, if possible, the direction in which they entered the trap should be noted, as this may indicate a source of entry. Basic room hygiene is essential, otherwise the sticky part of the traps will be filled with dirt and fluff instead of insects (Child, 1998; Edinburgh, 2022), but traps should still be checked weekly and more frequently in the summer months. These traps should be replaced over time as the captured insect may become a food source for other insects and, despite the longevity of the glue, it will eventually dry out. New traps should be set after a disinfestation treatment to monitor the success of the operation (Child, 1998) and all this information should be recorded.

12. Storage, Archiving and Handling of xylarium and herbarium samples

The methods of storing and archiving xylarium specimens are limited. Most xylarium organise their specimens in open racks, shelves, drawers, or closed cabinets with specimens arranged side by side. This furniture typology is mostly dependent on the size and scope of the collection, the physical and spatial conditions, and the economic resources available. There are, however, parameters that are not, such as the paramount importance of the specimens being completely dried and free of pests before storage.

It is important to emphasise that the presence and storage of histological slides associated with the xylarium, and herbarium will facilitate and speed up their use by researchers, avoiding the need for constant recourse to the wood or herbarium specimens for sectioning (Beeckman, 2003; Lamb & Curtis, 2005). Table 16 shows the materials and protocol for storing xylarium.

Table 16 - Materials and protocol for xylarium storage.

Materials	
I.	Open or preferably closed metal racks, shelves, drawers, or cabinets.
II.	Archival quality acid free labels.
III.	Acid-free glue. Wilken (2009) suggests the reversible adhesive grade A methylcellulose with a molecular weight of 1500.
Protocol	
I.	The height, width and thickness of the specimen will depend largely on what the original specimen allows and other issues such as availability, level of protection, transport, market value etc.
II.	Xylarium specimens are usually placed side by side.
III.	The identification label should face outwards if possible.
IV.	No coatings of any kind (varnish, wax or other) should be applied to the specimens as this may interfere with future application of analytical techniques.
V.	Samples should not be exposed to direct sunlight and should ideally be stored in a light-free environment in order to preserve, as far as possible, the natural colours of the different specimens.
VI.	If possible, a sheet of blotting paper should be used as a divider between each wood sample.

500

501 There are various methods to archive herbarium specimens, highly depending on the
502 collection's size, scope, physical, space, and economic conditions. The language should be
503 objective, balanced, and formal, devoid of ornamental, biased phrases. Additionally, the text's
504 grammar, spelling, and punctuation must be correct and consistent. Subjective evaluations will
505 be avoided, unless clearly marked as such. It is crucial always to ensure specimens are entirely
506 dried and free of pests before being stored. It is important to note that xylarium samples do not
507 require any specific material for proper storage. In contrast, herbarium necessitate the use of
508 particularly high-grade materials. Table 17 presents the materials and protocol for herbarium
509 storage.

510 Table 17 - Materials and protocol for herbarium storing.

Materials	
I.	Protective bags are constructed from flimsy acid-free paper with a weight of 60 gsm.
II.	Genus covers - archival natural history herbarium folders.
III.	Large format sealable (zipper) polythene bags.
Protocol	
I.	Attach the specimen to a mounting card using a thin adhesive and label the bottom-right corner accordingly.
II.	Place the document in a folder with a clear label, ensuring the label is also placed in the bottom-right corner.
III.	The specimen is ready for filing (RHS - Royal Horticultural Society, 2013).
IV.	Herbarium specimens are typically stacked atop one another in a standard format.

511 12.1. Storage facilities

512 Wooden shelves and cabinets are no longer used in storage facilities due to their
513 susceptibility to infestation (British Columbia Ministry of Forests, 1996). As a result, metal
514 furniture cabinets are the prevailing storage mechanism for xylarium and herbarium collections
515 (British Columbia Ministry of Forests, 1996). The most common herbarium storage system for
516 small and medium-sized samples involves metal cabinets with movable shelves and drawers.

Although not strictly necessary, closed metallic cabinets can effectively reduce damage caused by dust, insects, rodents, air pollutants, and light and provide a stable microenvironment with controlled relative humidity and temperature fluctuations (Donald R. Cumberland, 1993; Vieira, 2023; Wilken, 2009).

One significant aspect of these cabinets is the gasket lining the door jamb, which creates a positive seal against air infiltration and biological infestations. Older gaskets oxidised rapidly, losing flexibility and structural integrity in around two to five years primarily due to exposure to ultraviolet radiation. Furthermore, these materials were made from felt impregnated with arsenic or mercuric chloride, intended to kill any insects seeking access to the interior of the cabinet. Consequently, these materials pose a serious hazard for herbarium staff (Donald R. Cumberland, 1993). Long-lasting, chemically inert, and stable silicone polymer is a more suitable material for this purpose (Donald R. Cumberland, 1993; Edinburgh, 2022).

12.2. Archival System

The most prevalent approach for storing xylarium and herbarium samples is to file them in alphabetical order, with the genus and species being alphabetically arranged. Family classifications can also be maintained in alphabetical order or organized according to a phylogenetic sequence (British Columbia Ministry of Forests, 1996; Wilken, 2009).

12.3. Handling procedures

Handling procedures for xylarium samples differ from those for herbarium specimens. Generally, no specific requirements exist for handling xylarium samples. However, when the xylarium houses highly degraded, old, or fragile specimens, such as palm trees, centenarians, or archaeological samples, precautions should be taken to avoid accidents.

Wearing surgical gloves prevents the deposition of hand grease, which could result in oxidation in the future. Additionally, it is necessary to wear gloves when handling known toxic species to protect the handler. Even if the wood is free of dangerous chemical compounds, it is important to consider potential allergic reactions some individuals may have to specific types of wood (Meier, 2022; Osborn, 2022).

Handling specimens, irrespective of their nature, should always be considered a challenging operation that necessitates thoughtful planning and focus. Especially with herbarium specimens, the careless handling of excessively dehydrated, delicate, and occasionally harmed vegetative samples can be exceptionally taxing and destructive. Consequently, Table 18 outlines the procedure that must be pursued.

Table 18 – Herbarium handling protocol (British Columbia Ministry of Forests, 1996; KSU Herbarium, 2002; RHS - Royal Horticultural Society, 2013; Wilken, 2009).

I.	Whenever feasible, please ensure the specimen remains affixed to the sheet and prevent any potential bending.
II.	To transport the specimen's folder from the storage cabinet to the examination table, use a rolling cart, a sturdy surface such as a plywood board or stiff cardboard, or a box.
III.	Always examine the specimen on a flat, sturdy, and horizontal surface, avoiding standing.
IV.	When stacking folders, be mindful of their weight to prevent damage to the specimens at the bottom.

V.	When carrying out microscopic observations, it is preferable to use a microscope equipped with task lights that can be attached to an adjustable arm (similar to a boom stand).
VI.	It is recommended to avoid flipping through sheets inside a folder. Instead, take out all the sheets together and analyse each one individually.
VII.	Ensure that mounted specimens are placed specimen-side up in the sheets and avoid turning them over.
VIII.	Finally, replace the specimen sheets in the same order as previously organized.
IX.	Before refiling, it is advisable to carry out a preventive conservation revision to identify pests, signs of pests, broken fragments, or incorrect labels.
X.	It is not recommended to leave specimens outside of the cabinets in the herbarium room overnight because most herbarium beetles are more active during the night.

13. Xylaria and Herbaria trends and future prospects

As far back as 1973, William L. Stern lamented the ongoing decline of institutional wood collections (Stern, 1973). The future trajectory of xylaria and herbaria collections presents however an intriguing and evolving narrative, since, historically, the vitality of most collections has been closely tethered to the commitment of one or more dedicated researchers within a particular institution. With the departure of such researchers, activities and, at times, the very existence of the xylarium and herbarium have been imperilled. The reason for this, lays frequently in the undeniable difficulty of articulating the value of biocultural collections to the general public. Demonstrating the tangible benefits that stem from the resources and dedication invested in establishing, curating, and expanding such collections remains an ongoing hurdle for most scientists who advocate for them. However, recent societal developments have transformed this landscape for the better.

In the particular case of xylaria, many venerable collections have witness closures or amalgamations. For instance, in the past 55 years, the Forest Products Laboratory (FPL) in Madison, WI, has absorbed the Yale University xylarium, the Chicago Field Museum xylarium, the Jessup wood collection, and the Houghton, Michigan xylarium (Wiedenhoeft, 2014). It is worth noting, however, that in certain regions of the world, particularly in Brazil, new xylaria are sprouting, suggesting that rather than an overall decline, we may be witnessing a shift in the epicenters of active xylaria (Wiedenhoeft, 2014).

With a growing emphasis on combatting illegal logging, species extinction and climate change, xylaria and herbaria has witnessed a growing in inquiries from botanists, historians and federal law enforcement agents, seeking scientific expertise for all types of identifications. The strength of these identifications fundamentally hinges on the robustness of the reference collection underpinning them; hence, collections are central to ensuring scientifically sound forensic identification. Through such undertakings, it has become possible to enhance the recognition and appreciation of the scientific and societal value of a xylaria and herbaria. Demonstrating that a xylaria, with its seemingly unassuming assemblage of wooden blocks, and herbaria with its collection of old dried plants possesses tangible worth may serve as a means to elevate public awareness regarding all biocultural collections, thereby encouraging their meticulous curation, expansion, and preservation for posterity.

14. Conclusion

In conclusion, implementing measures to prevent degradation, decrease contamination risk, and limit the exposure of collected objects to degradation factors would lead to

584 maintenance that is less demanding, restoration-focused, and consumes fewer time and
585 economic resources for any collection, storeroom, or museum. Preventing or mitigating
586 collection degradation directly affects the scientific, cultural, and economic value of a museum
587 institution, making it the ultimate goal. This is particularly crucial when managing storage areas
588 with highly degradable and consumable materials, such as wood samples and dried plants.

589 Although producing a sturdy, scientifically-sound and future-proof xylarium and herbarium
590 requires significant amounts of time, physical effort, and economic and material resources, it is
591 crucial that the construction and structuring of this research tool conform to the necessary steps
592 in order to achieve high-quality results.

593 **Acknowledgements**

594 We would like to acknowledge the Portuguese Fundação para a Ciência e Tecnologia (FCT)
595 for the PhD grant (UI/BD/151009/2021) for José Luis Amorim Silva and for funding the Strategic
596 Projects UID/EAT/0622/2016 of CITAR - Research Center for the Science and Technology of the
597 Arts, Porto, Portugal

References

- Archer, W. A. (1950). NEW PLASTIC AID IN MOUNTING HERBARIUM SPECIMENS. *Rhodora*, 52(624), 298–299. <http://www.jstor.org/stable/23306129>
- Barker, J. A., & Flinders, B. A. H. (2005). *Key to a Selection of Arid Australian Hardwoods & Softwoods*. https://keys.lucidcentral.org/keys/v3/arid/default_wip.htm
- Bebber, D. P., Carine, M. A., Wood, J. R. I., Wortley, A. H., Harris, D. J., Prance, G. T., Davidse, G., Paige, J., Pennington, T. D., Robson, N. K. B., & Scotland, R. W. (2010). Herbaria are a major frontier for species discovery. *Proceedings of the National Academy of Sciences*, 107(51), 22169–22171. <https://doi.org/10.1073/pnas.1011841108>
- Beck, J. B., & Semple, J. C. (2015). Next-Generation Sampling: Pairing Genomics with Herbarium Specimens Provides Species-Level Signal in *Solidago* (Asteraceae). *Applications in Plant Sciences*, 3(6), 1500014. <https://doi.org/10.3732/apps.1500014>
- Beeckman, H. (2003). A xylarium for the sustainable management of biodiversity: the wood collection of the Royal Museum for Central Africa, Tervuren, Belgium. *Bulletin de l'APAD*, 26. <https://doi.org/10.4000/apad.3613>
- Briggs, D., Sell, P. D., Block, M., & I'ons, R. D. (1983). MERCURY VAPOUR: A HEALTH HAZARD IN HERBARIA. *New Phytologist*, 94(3), 453–457. <https://doi.org/10.1111/j.1469-8137.1983.tb03458.x>
- British Columbia Ministry of Forests. (1996). *Techniques and Procedures for Collecting, Preserving, Processing, and Storing Botanical Specimens*. [file:///C:/Users/jl_6/Desktop/Doutoramento geral/PhD aulas - horários/Disciplinas/Preventive Conservation of Built Heritage, Movable and Integrated Assets/wp18.pdf](file:///C:/Users/jl_6/Desktop/Doutoramento%20geral/PhD%20aulas%20-%20hor%C3%A1rios/ Disciplinas/ Preventive%20Conservation%20of%20Built%20Heritage,%20Movable%20and%20Integrated%20Assets/wp18.pdf)
- Burke, J. (1999). *Anoxic Microenvironments: A Treatment For Pest Control Conserve O Gram 3/9*. [file:///C:/Users/jl_6/Desktop/Doutoramento geral/PhD aulas - horários/Disciplinas/Preventive Conservation of Built Heritage, Movable and Integrated Assets/Conserv-O-Gram-Author Guidelines.pdf](file:///C:/Users/jl_6/Desktop/Doutoramento%20geral/PhD%20aulas%20-%20hor%C3%A1rios/ Disciplinas/ Preventive%20Conservation%20of%20Built%20Heritage,%20Movable%20and%20Integrated%20Assets/Conserv-O-Gram-Author%20Guidelines.pdf)
- Cabral, C., & Sales, F. (2018). *Restoration of the Willkomm Herbarium: curating an old collection at COI. Universidade de Coimbra*. https://www.uc.pt/en/herbario_digital/willkomm_herbarium/restauro
- Cappitelli, F., & Sorlini, C. (2008). Microorganisms Attack Synthetic Polymers in Items Representing Our Cultural Heritage. *Applied and Environmental Microbiology*, 74(3), 564–569. <https://doi.org/10.1128/AEM.01768-07>
- Carlquist, S. (1982). The Use of Ethylenediamine in Softening Hard Plant Structures for Paraffin Sectioning. *Stain Technology*, 57(5), 311–317. <https://journals.scholarsportal.info/browse/00389153/v57i0005>
- Cartwright, C. R. (2015). The principles, procedures and pitfalls in identifying archaeological and historical wood samples. *Annals of Botany*, 116(1), 1–13. <https://doi.org/10.1093/aob/mcv056>
- Child, R. E. (1998). *Monitoring Insect Pests With Sticky Traps Conserve O Gram 3/7*. [file:///C:/Users/jl_6/Desktop/Doutoramento geral/PhD aulas - horários/Disciplinas/Preventive Conservation of Built Heritage, Movable and Integrated Assets/wp18.pdf](file:///C:/Users/jl_6/Desktop/Doutoramento%20geral/PhD%20aulas%20-%20hor%C3%A1rios/ Disciplinas/ Preventive%20Conservation%20of%20Built%20Heritage,%20Movable%20and%20Integrated%20Assets/wp18.pdf)

- 639 horários/Disciplinas/Preventive Conservation of Built Heritage, Movable and Integrated
640 Assets/Conserv-O-Gram-03-07.pdf
- 641 Collins, R. A., & Cruickshank, R. H. (2012). The seven deadly sins of DNA barcoding. *Molecular*
642 *Ecology Resources*, n/a-n/a. <https://doi.org/10.1111/1755-0998.12046>
- 643 D. Bridson, & L. Forman. (2010). *Herbarium Handbook* (Third Edition, 3rd). Royal Botanic
644 Gardens, Kew.
- 645 Donald R. Cumberland, Jr. (1993). *Installing The Retrofit Gasket Kit Conserve O Gram 4/3*.
646 <https://www.nps.gov/museum/publications/conservoogram/04-03.pdf>
- 647 Drobnik, J. (2007). *Zielnik i zielnikoznawstwo* (Warsaw). Wydawnictwo Naukowe PWN.
648 <https://ksiegarnia.pwn.pl/Zielnik-i-zielnikoznawstwo,68737776,p.html>
- 649 Drobnik, J. (2008). Modern techniques of herbarium protection. *Scripta Facultatis Rerum*
650 *Naturalium Universitatis Ostraviensis*, 186, 243–246.
- 651 Edinburgh, R. B. G. (2022). *CARE AND CONSERVATION OF SPECIMENS*.
652 [https://www.rbge.org.uk/science-and-conservation/herbarium/specimen-preparation-](https://www.rbge.org.uk/science-and-conservation/herbarium/specimen-preparation-care/care-and-conservation-of-herbarium-specimens/)
653 [care/care-and-conservation-of-herbarium-specimens/](https://www.rbge.org.uk/science-and-conservation/herbarium/specimen-preparation-care/care-and-conservation-of-herbarium-specimens/)
- 654 Feuchter-Schawelka, A., & Freitag, W. & G. D. (2001). *Alte Holzsammlungen. Die Ebersberger*
655 *Holzbibliothek: Vorgänger, Vorbilder und Nachfolger*. (Deutscher Sparkassen Verlag, Ed.).
- 656 Funk, V. (2007). *100 Uses For a Herbarium (well at least 72)*. Fairchild Tropical Botanic Garden.
657 <http://www.virtualherbarium.org/vh/100usesaspt.html>
- 658 GBIF. (2021). *GBIF | Global Biodiversity Information Facility*. <https://www.gbif.org/>
- 659 Greve, M., Lykke, A. M., Fagg, C. W., Gereau, R. E., Lewis, G. P., Marchant, R., Marshall, A. R.,
660 Ndayishimiye, J., Bogaert, J., & Svenning, J.-C. (2016). Realising the potential of herbarium
661 records for conservation biology. *South African Journal of Botany*, 105, 317–323.
662 <https://doi.org/10.1016/j.sajb.2016.03.017>
- 663 Guarino, S., Basile, S., Caimi, M., Carratello, A., Manachini, B., & Peri, E. (2020). Insect pests of
664 the Herbarium of the Palermo botanical garden and evaluation of semiochemicals for the
665 control of the key pest *Lasioderma serricorne* F. (Coleoptera: Anobiidae). *Journal of*
666 *Cultural Heritage*, 43, 37–44.
667 <https://doi.org/https://doi.org/10.1016/j.culher.2019.10.009>
- 668 Hawks, C. (2001). Historical survey of the sources of contamination of ethnographic materials in
669 museum collections. *Collection Forum*, 16(1–2), 2–11.
- 670 Hawks, C., Makos, K., Bell, D., Wambach, P. F., & Burroughs, G. E. (2004). An inexpensive method
671 to test for mercury vapor in herbarium cabinets. *TAXON*, 53(3), 783–790.
672 <https://doi.org/10.2307/4135451>
- 673 Heath, J., Jones, M., & Wallace, M. (2017). *Preserving plants for the future*. Royal Botanical
674 Gardens Kew. <https://www.kew.org/read-and-watch/preserving-plants-for-the-future>
- 675 Hoadley, R. B. (2000). *Understanding Wood: A Craftsman's Guide to Wood Technology*. The
676 Taunton Press.

677 IPBES. (n.d.). *IPBES - Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem*
678 *Services*. Retrieved January 13, 2022, from <https://www.ipbes.net/>

679 IPCC. (2014). *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III*
680 *to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*.
681 https://www.ipcc.ch/site/assets/uploads/2018/05/SYR_AR5_FINAL_full_wcover.pdf

682 IUCN. (2022). *IUCN Red List of Threatened Species*. <https://www.iucnredlist.org/>

683 Jiao, L., Yu, M., Wiedenhoef, A. C., He, T., Li, J., Liu, B., Jiang, X., & Yin, Y. (2018). DNA Barcode
684 Authentication and Library Development for the Wood of Six Commercial Pterocarpus
685 Species: the Critical Role of Xylarium Specimens. *Scientific Reports*, 8(1), 1945.
686 <https://doi.org/10.1038/s41598-018-20381-6>

687 KSU Herbarium. (2002). *Integrated Pest Management Plan*. Kansas State University Herbarium.
688 <https://www.k-state.edu/herbarium/>

689 Kurmanow, M. G. (2019). Review of biocides used as prevention and intervention measures for
690 historic artefacts, with special regard to herbaria collections. *Notes Konserwatorski*, 21,
691 121–161. <https://doi.org/10.36155/NK.21.00004>

692 Lamb, S. H., & Curtis, A. B. (2005). *A Guide for Developing a Wood Collection* (Forest Products
693 Society for the International Wood Collectors Society, Ed.).

694 Lieutier, F., Day, K. R., Evans, A., Claude, B., & F., G. (2004). *Bark and Wood Boring Insects in*
695 *Living Trees in Europe, a Synthesis* (F. Lieutier, K. R. Day, A. Battisti, J.-C. Grégoire, & H. F.
696 Evans, Eds.). Springer Netherlands. <https://doi.org/10.1007/978-1-4020-2241-8>

697 Lynch, A. H., Gasson, P. E., & Lens, F. (2016). *IAWA Index Xylariorum*.
698 <https://globaltimbertrackingnetwork.org/products/iawa-index-xylariorum/>

699 Meier, E. (2022). *WOOD ALLERGIES AND TOXICITY*. The Wood Database. [https://www.wood-](https://www.wood-database.com/wood-articles/wood-allergies-and-toxicity/)
700 [database.com/wood-articles/wood-allergies-and-toxicity/](https://www.wood-database.com/wood-articles/wood-allergies-and-toxicity/)

701 Michalski, S., Jr., J. L. P., CCI, & ICCROM. (2016). The ABC Method: a risk management approach
702 to the preservation of cultural heritage. In *The ABC Method: a risk management approach*
703 *to the preservation of cultural heritage*. Government of Canada, Canadian Conservation
704 Institute (CCI), International Centre for the Study of the Preservation and Restoration of
705 Cultural Property (ICCROM). [https://www.iccrom.org/sites/default/files/2017-](https://www.iccrom.org/sites/default/files/2017-12/risk_manual_2016-eng.pdf)
706 [12/risk_manual_2016-eng.pdf](https://www.iccrom.org/sites/default/files/2017-12/risk_manual_2016-eng.pdf)

707 New Your Botanical Garden, N. (n.d.). *INDEX HERBARIORUM*. Retrieved April 21, 2022, from
708 <http://sweetgum.nybg.org/science/ih/>

709 NPS National Park Services. (2000). The Museum Handbook Part I: Museum Collections. In *NPS*
710 *Museum handbook* (Revised ed, p. 1441). National Park Service, U.S. Dept. of the Interior.

711 Osborn, L. (2022). *TOXIC WOODS*. Woodcraft Magazine.
712 https://www.woodcraft.com/blog_entries/toxic-woods

713 Pereira, M., & Hammond, B. (2001). *Chronology Of Pesticides Used On National Park Service*
714 *Collections - " Conserve O Gram 2/16*. file:///C:/Users/jl_6/Desktop/Doutoramento
715 *geral/PhD aulas - horários/ Disciplinas/Preventive Conservation of Built Heritage, Movable*
716 *and Integrated Assets/Conserv-O-Gram-02-16.pdf*

- Pereira, M., & Wolf, S. J. (2001). *Physical Properties And Health Effects of Pesticides Used On National Park Service Collections Conserve O Gram* 2/17. file:///C:/Users/jl_6/Desktop/Doutoramento geral/PhD aulas - horários/ Disciplinas/ Preventive Conservation of Built Heritage, Movable and Integrated Assets/ Conserv-O-Gram-Author Guidelines.pdf
- Perkins, K. (2022). *Preparation of Plant Specimens for Deposit as Herbarium Vouchers*. University of Florida Herbarium (FLAS). <https://www.floridamuseum.ufl.edu/herbarium/voucher.htm>
- Pinniger, D. B. (1998). *Controlling Insect Pests: Alternatives To Pesticides Conserve O Gram* 3/8. file:///C:/Users/jl_6/Desktop/Doutoramento geral/PhD aulas - horários/ Disciplinas/ Preventive Conservation of Built Heritage, Movable and Integrated Assets/ Conserv-O-Gram-Author Guidelines.pdf
- Pyke, G. H., & Ehrlich, P. R. (2010). Biological collections and ecological/environmental research: a review, some observations and a look to the future. *Biological Reviews*, 85(2), 247–266. <https://doi.org/10.1111/j.1469-185X.2009.00098.x>
- Raphael, T. (1994). *An Insect Pest Control Procedure: The Freezing Process Conserve O Gram* 3/6.
- RHS - Royal Horticultural Society. (2013). *RHS Herbarium guide to making Herbarium Specimens*. <https://www.plantheritage.org.uk/media/1894/rhs-guide-to-making-herbarium-specimens.pdf>
- Rumball, N., & Pinniger, D. (2003). Use of temperature to control an infestation of biscuit or drugstore beetle *Stegobium Paniceum* (L.) (Coleoptera: anobidae) in a large economic botany collection. *Collection Forum*, 18(1–2), 50–58. <https://www.bcin.ca/bcin/detail.app?id=233947>
- Schmitz, N. (ed.) (ed.), Beeckman, H., Blanc-Jolivet, C., Boeschoten, L. E., Braga, J. J. W. B., Cabezas, J. A., Chaix, G., Crameri, S., Degen, B., Deklerck, V., Dormontt, E., Espinoza, E., Gasson, P., Haag, V., Helmling, S., Horacek, M., Koch, G., Lancaster, C., Lens, F., ... Zuidema, P. (2020). Overview of current practices in data analysis for wood identification. A guide for the different timber tracking methods. In *Global Timber Tracking Network* (Issue June, p. 142). GTTN - Global Timber Tracking Network. <https://doi.org/10.13140/RG.2.2.21518.79689>
- Schmitz, N., Haag, V., Blanc-Jolivet, C., Boner, M., Cervera, M. T., Chavesta, M., Cronn, R., Deklerck, V., Diaz-Sala, C., Dormontt, E., Gasson, P., Gehl, D., Hermanson, J. C., Honorio Coronado, E., Lancaster, C. A., Lens, F., Liendo Hoyos, E., Martínez-Jarquín, S., Montenegro, R. A., ... Wiemann, M. (2019). *General sampling guide for timber tracking. How to collect reference samples for timber identification*. <https://doi.org/10.13140/RG.2.2.26883.96806>
- Staudinger, M. D., Grimm, N. B., Staudt, A., Carter, S. L., Stuart III, F. S., Kareiva, P., Ruckelshaus, M., & Stein, B. A. (2012). *Impacts of climate change on biodiversity, ecosystems, and ecosystem services: technical input to the 2013 National Climate Assessment*. <http://pubs.er.usgs.gov/publication/70039460>
- Stern, W. L. (1973). The wood collection: what should be its future? *Arnoldia*, 33, 67–80.
- Strang, T. J. K. (1992). A review of published temperatures for the control of insect pests in museums. *Collection Forum*, 8(2), 41–67.

- Toriti, M., Durand, A., & Fohrer, F. (2021). *Traces of Common Xylophagous Insects in Wood* (S. N. Switzerland, Ed.). Springer International Publishing. <https://doi.org/10.1007/978-3-030-66391-9>
- U.C., U. de C. (2008). *COI - Herbarium of University of Coimbra*. https://www.uc.pt/en/herbario_digital/activities/pestcontrol
- Vieira, C. (2023). Hortus Siccus – o herbário como um testemunho botânico visual, científico e técnico. In *Modos de Editar - Arquivo em Aberto* (pp. 88–96). Edições i2ADS – Instituto de Investigação em Artes, Design e Sociedade Faculdade de Belas Artes da Universidade do Porto.
- Whitmore, T. C. (1965). THE HONIARA TECHNIQUE. *TAXON*, 14(5), 164–165. <https://doi.org/10.2307/1217552>
- Whitten, W. M., Williams, N. H., & Glover, K. V. (1999). Sulphuryl fluoride fumigation: effect on DNA extraction and amplification from herbarium specimens. *TAXON*, 48(3), 507–510. <https://doi.org/10.2307/1224563>
- Wiedenhoeft, A. C. (2014). Curating xylaria. In *Curating Biocultural collections A handbook* (Vol. 9). Forest Service U.S. DEPARTMENT OF AGRICULTURE.
- Wilken, D. H. (2009). *Preparing And Storing Herbarium Specimens Conserve O Gram 11/12*. [file:///C:/Users/jl_6/Desktop/Doutoramento geral/PhD aulas - horários/Disciplinas/Preventive Conservation of Built Heritage, Movable and Integrated Assets/Conserv-O-Gram-Author Guidelines.pdf](file:///C:/Users/jl_6/Desktop/Doutoramento%20geral/PhD%20aulas%20hor%C3%A1rios/ Disciplinas/Preventive%20Conservation%20of%20Built%20Heritage,%20Movable%20and%20Integrated%20Assets/Conserv-O-Gram-Author%20Guidelines.pdf)
- Yu, M., Jiao, L., Guo, J., Wiedenhoeft, A. C., He, T., Jiang, X., & Yin, Y. (2017). DNA barcoding of vouchered xylarium wood specimens of nine endangered Dalbergia species. *Planta*, 246(6), 1165–1176. <https://doi.org/10.1007/s00425-017-2758-9>