

Three-phase equilibria of hydrates from computer simulation. II. Finite-size effects in the carbon dioxide hydrate

J. Algaba,¹ S. Blazquez,² E. Feria,¹ J. M. Míguez,¹ M. M. Conde*,³ and F. J. Blas*¹

¹Laboratorio de Simulación Molecular y Química Computacional, CIQSO-Centro de Investigación en Química Sostenible and Departamento de Ciencias Integradas, Universidad de Huelva, 21007 Huelva, Spain

²Departamento de Química Física, Facultad de Ciencias Químicas, Universidad Complutense de Madrid, 28040 Madrid, Spain

³Departamento de Ingeniería Química Industrial y Medio Ambiente, Escuela Técnica Superior de Ingenieros Industriales, Universidad Politécnica de Madrid, 28006, Madrid, Spain

(*Electronic mail: felipe@uhu.es)

(Dated: 28 December 2025)

In this work, the effects of finite size on the determination of the three-phase coexistence temperature (T_3) of carbon dioxide (CO₂) hydrate have been studied by molecular dynamic simulations and using the direct coexistence technique. According to this technique, the three phases involved (hydrate-aqueous solution-liquid CO₂) are placed together in the same simulation box. By varying the number of molecules of each phase it is possible to analyze the effect of simulation size and stoichiometry on the T_3 determination. In this work, we have determined the T_3 value at 8 different pressures (from 100 to 6000 bar) and using 6 different simulation boxes with different numbers of molecules and sizes. In 2 of these configurations, the ratio of the number of water and CO₂ molecules in the aqueous solution and the liquid CO₂ phase is the same as in the hydrate (stoichiometric configuration). In both stoichiometric configurations, the formation of a liquid drop of CO₂ in the aqueous phase is observed. This drop, which has a cylindrical geometry, increases the amount of CO₂ available in the aqueous solution and can in some cases lead to the crystallization of the hydrate at temperatures above T_3 , overestimating the T_3 value obtained from direct coexistence simulations. The results obtained in this work highlight that the choice of a correct initial configuration is essential to accurately estimate the three-phase coexistence temperature of hydrates by direct coexistence simulations.

*Corresponding authors: felipe@uhu.es and maria.mconde@upm.es

I. INTRODUCTION

Gas hydrates, crystalline structures formed by the encapsulation of gas molecules within water cages, have garnered significant attention due to their implications in energy storage, gas separation, and environmental processes¹⁻⁴. Gas hydrates are a subject of considerable interest due to their diverse applications depending on the guest molecule they encapsulate. Hydrogen hydrates, for instance, have been proposed as an alternative method for storing hydrogen, presenting potential applications in energy storage⁵⁻⁷. On the other hand, methane hydrates represent a natural and alternative source of energy, particularly when found in seabed deposits^{1,8-11}. However, in the context of natural gas transport, hydrate formation within pipelines poses a significant challenge, prompting research into hydrate inhibition strategies. Ionic liquids, for instance, have garnered attention as inhibitors in mitigating hydrate formation during gas transport^{12,13}. In addition to their occurrence on Earth, clathrate hydrates play a crucial role in the context of icy satellites within our solar system, particularly under conditions involving salty water.¹⁴⁻²⁰

Particularly, carbon dioxide (CO₂) hydrates have become a pivotal area of research due to their profound implications in addressing global environmental challenges, particularly those associated with the increasing levels of carbon dioxide in the Earth's atmosphere^{21,22}. Thus, CO₂ hydrates are of particular interest in the context of carbon capture and sequestration (CCS) strategies, which aim to mitigate the adverse

impacts of anthropogenic CO₂ emissions.²³⁻²⁷

In 2015, some of us studied the three-phase equilibrium line of CO₂ hydrate employing the Direct Coexistence technique²⁸. We showed that to accurately replicate the experimental behavior of T_3 at various pressures, it was necessary to introduce positive deviations (specifically, $\chi=1.13$) to the energetic Lorentz-Berthelot (LB) rules. By incorporating these deviations, we successfully reproduced the three-phase line of CO₂ hydrate, as shown in Fig. 2 (black crosses). It is noteworthy that our focus was primarily on the coexistence phase when CO₂ was in its liquid state (i.e., above 4 MPa).

When delving into the realm of simulations concerning CO₂ hydrates, it is noteworthy to highlight several notable contributions in the field. In a pioneering work, Tung et al.²⁹ computed the three-phase line of the carbon dioxide hydrate using the TIP4P-Ew³⁰ model for water and the EPM2³¹ for CO₂. They obtained a T_3 of 285(2) K at 400 bar, in good agreement with our work of 2015²⁸. Another such significant work was conducted by Costandy et al. in 2015³². Similar to the work of Míguez et al.²⁸, they used the direct coexistence technique and utilized the same force fields (TIP4P/Ice for water and TraPPE for CO₂). However, they employed different cutoff values and altering water-CO₂ interactions due to a distinct deviation from the LB rules. Their findings, particularly at 400 bar, yielded a T_3 of 283.5(1.7) K, differing slightly from the results obtained by Míguez et al. of 287(2) K at the same pressure²⁸. In a separate endeavor, Waage and coworkers³³, employing a different technique, evaluated the dissociation temperature of CO₂ hydrate using interfacial energy calculations. Remarkably, they employed the same force fields as Costandy et al.³² and Míguez et al.²⁸. However, the dispersive interactions and cutoff values implemented by

Waage and the team were distinct from the other works. Their results showed a T_3 of 284.5 K at 400 bar.

There exists other significant works in which the T_3 of CO₂ hydrates is computed using different force fields for water and CO₂^{34,35} or using Monte Carlos simulations in the Grand Canonical ensemble³⁶. Even there are theoretical works in which the three-phase equilibrium of this hydrate is evaluated by using PC-SAFT and Peng-Robinson equation³⁷ or by using semiempirical equations³⁸. Theoretical calculations using the van der Waals-Platteeuw have also been employed to evaluate the occupation of the CO₂ hydrates³⁹. Also, the kinetics of these hydrates have been studied^{29,40}. Even the addition of salt to the hydrates (due to they can be found at the seabed) has been computationally studied for CO₂ hydrates⁴¹ (and recently for methane hydrates in a wide range of concentrations^{42,43}). Collectively, these works offer insights into the thermodynamics and stability of hydrates under diverse conditions.

Importantly, in 2022 we developed a new methodology to evaluate the T_3 of gas hydrates baptized as the *solubility method*^{44,45}. In short, by calculating on the one hand the solubility of methane in a gas-liquid system (which decreases with temperature) and on the other hand solubility of methane in a hydrate-liquid system (which increases with temperature). The two solubility curves intersect at a certain temperature that is the T_3 at a fixed pressure. We employed this approach for the methane hydrate⁴⁴ and for the carbon dioxide hydrate⁴⁵ with larger system sizes than in the direct coexistence simulations of previous works but using the same force fields (TIP4P/Ice + TraPPE) and the same cutoff. By using this methodology we estimated a T_3 of 290(2) K at 400 bar which slightly differs from previous calculations.

The discussed previous works suggest that system sizes or cutoff values can have an effect on the determination of the T_3 . This observation aligns with earlier research that has delved into finite-size effects across various systems. Previous studies, for Lennard-Jones potentials⁴⁶ or for the evaluation of surface tensions^{47,48}, have investigated the influence of finite-size effects in different contexts. Additionally, the impact of finite sizes has been explored in square well fluids⁴⁹. Notably, a recent study conducted a comprehensive analysis of how finite sizes affect the determination of the melting temperature of ice⁵⁰. This body of research collectively underscores the significance of considering finite-size effects in simulations across various systems and provides valuable insights into their implications on observed phenomena.

In this work, we have conducted a thorough examination of the finite-size effects influencing the determination of the three-phase equilibrium temperature for CO₂ hydrates. Similar to our parallel work on methane hydrate finite-size effects, we have employed the TIP4P/Ice force field for water and the TraPPE for CO₂, and we systematically have varied system sizes by manipulating the number of molecules in the hydrate, liquid, and gas phases.

The structure of our study is organized as follows: Section II outlines the methodology and simulation details, providing a comprehensive overview of our approach. In Section III, we present the results, encompassing the three-phase equilibrium

temperatures for different system sizes of carbon dioxide hydrate at different pressures. Finally, Section IV encapsulates the main conclusions drawn from this extensive study, summarizing key findings and insights into the impact of finite-size effects on the determination of carbon dioxide hydrate equilibrium temperatures.

II. METHODOLOGY AND SIMULATION DETAILS

In order to study finite-size effects on the three-phase coexistence line of the CO₂ hydrate, we have employed the same methodology introduced by Conde and Vega.⁵¹ Following this methodology, the three phases involved (CO₂ hydrate, aqueous, and pure CO₂ phases) are put together in the same simulation box. Due to the use of periodic boundary conditions, this arrangement ensures that the three phases are in contact, with one of the phases surrounded by the other two. In this work, all the simulation boxes have been generated using the same arrangement explained previously regardless of the size and number of molecules of the system.

The CO₂ hydrate exhibits a sI crystallographic structure. The unit cell of the sI hydrate structure is formed by six tetradecahedrons, 5¹²6², and two pentagonal dodecahedrons, 5¹² cages. Hence, one sI hydrate unit cell is formed by 46 molecules of water and 8 cages. The initial sI CO₂ hydrate unit cell has been built taking explicitly into account that hydrates are proton-disordered structures. To obtain this arrangement, hydrogen atoms are placed using the algorithm proposed by Buch *et al.*⁵² This allows the generation of solid configurations satisfying the Bernal-Fowler rules,⁵³ with zero (or at least negligible) dipole moment. Also, in this work, we have assumed single occupancy of all the CO₂ hydrate cages, which means that each cage of the hydrate is occupied by a molecule of CO₂. In this work, 6 different initial configurations have been built by varying the replication factor of the hydrate unit cell and the number of initial molecules of water and CO₂ in the aqueous and CO₂ phases respectively (see Table I),

Once the initial configuration is generated, *NPT* simulations are carried out by fixing the pressure at 8 different values (see Table II) and varying the temperature. If the selected temperature is above the temperature at which the three phases coexist, T_3 , the hydrate phase becomes unstable and melts. If the selected temperature is below the T_3 , the hydrate phase will grow until extinguishing the CO₂ and/or aqueous phases depending on the initial amount of molecules of CO₂ and water in both phases respectively. If the amount of water and CO₂ in the aqueous and CO₂ phases is the same as that in the hydrate phase (stoichiometric composition), at $T < T_3$, the hydrate will grow until extinguishing both phases, and the final configuration box will be formed by a unique bulk hydrate phase (configurations 1 and 3 from Table I). Contrary, if the amount of CO₂ molecules in the CO₂ phase is larger than the number of molecules of water in the aqueous phase (using the stoichiometric composition of the hydrate as a reference), the hydrate will grow until extinguishing the aqueous phase. In this case, the final configuration box will be formed

Configuration	Hydrate phase			H ₂ O-rich liquid phase	CO ₂ -rich liquid phase	Stoichiometric
	Unit Cell	Water	CO ₂	Water	CO ₂	
0	2x2x2	368	64	368	192	No
1	2x2x2	368	64	368	64	Yes
2	3x3x2	828	144	828	432	No
3	3x3x3	1242	216	1242	648	No
4	3x3x3	1242	216	1242	216	Yes
5	4x4x2	1472	256	1472	768	No
6	4x4x4	2944	512	2944	1536	No

TABLE I: Initial number of molecules for each phase (hydrate phase, liquid phase, and gas phase) in the different configurations studied in this work. For the hydrate phase, the replication factor of the unit cell is indicated in each case. Configuration 0 corresponds to the original system studied by Míguez *et al.*²⁸ but it is not studied in this paper. The last column indicates whether the system composition is stoichiometric.

by a CO₂ hydrate phase in equilibrium with a pure CO₂ phase (configurations 2, 4, 5, and 6 from Table I). The evolution of the system can be determined from the analysis of the potential energy as a function of the simulation time. If $T < T_3$, the potential energy will decrease as a consequence of the increment of hydrogen bonds formed when the hydrate phase grows. Contrary, if $T > T_3$, the potential energy will increase as a consequence of the destruction of the hydrogen bonds of the CO₂ hydrate phase. Finally, the dissociation temperature, T_3 , is estimated by analyzing if the hydrate phase grows or melts, at a fixed value of pressure, as a function of the temperature. The T_3 value is estimated as the intermediate value between the lowest temperature at which the CO₂ hydrate melts and the highest temperature at which the CO₂ hydrate grows at a given value of pressure. Some of us have previously used this methodology to study the three-phase coexistence line of the CO₂ hydrate.²⁸ However, in this previous work, only one size system was used and the finite size effects were not taken into account.

In this work, all the molecular dynamic simulations have been carried out using GROMACS (version 4.6, double precision).⁵⁴ CO₂ molecules are modeled through the TraPPE (Transferable Potentials for Phase Equilibria) force field⁵⁵ and water molecules are described using the widely-known TIP4P/Ice model.⁵⁶ As in our previous work,²⁸ Lorentz-Berthelot combining rules have been applied, but the Berthelot combining rule for the unlike dispersive interactions between water and CO₂ molecules has been modified by a $\xi = 1.13$ factor. This factor was used in order to match the results obtained in our previous work²⁸ with experimental results taken from the literature. However, must be taken into account that this factor is rectifying the discrepancies with the experimental results, not only for the weakness of the unlike interaction descriptions of the molecular models but also for the finite-size effects presented in our previous study. We have used a Verlet leapfrog algorithm⁵⁷ to numerically solve Newton's motion equations. The time step used is 2, fs since all the models employed in this work are rigid. We have also used the Nosé-Hoover thermostat⁵⁸ and the Parrinello-Rahman barostat,⁵⁹ to ensure that simulations are performed at constant temperature and pressure. The time constant used for both, thermostat and barostat, is 2ps. It is important to remark that the barostat has been applied anisotropically to

avoid any stress from the solid hydrate structure. We have used a cutoff distance of 1.0nm for dispersive and Coulombic interactions. No long-range corrections have been applied for the dispersive interactions. For the case of the coulombic interactions, long-range particle mesh Ewald (PME) corrections⁶⁰ have been applied with a width of the mesh is 0.1, nm and a relative tolerance of 10^{-5} .

III. RESULTS

In this section, we show the results obtained to understand the role of finite-size effects in determining the dissociation temperature, as a function of pressure, of the CO₂ hydrate. As in paper I⁶¹, we estimate the T_3 in different systems of varying sizes. Table I shows the initial number of molecules forming each phase and the unit cell replication factor for the 6 size-dependent configurations. We have also included, for comparison reasons, the configuration 0 corresponding to the system studied by Míguez *et al.*²⁸ In this work, we extend the study of paper I and consider 8 different pressures in each configuration, from 100 to 6000 bar. This allows to analyze how the T_3 depends on the size of the system simulated in a wide range of pressures.

As in our previous work, we consider two different scenarios: systems in which the H₂O- and CO₂-rich liquid phases are formed from stoichiometric configurations, and systems in which the liquid phases are formed from configurations with a higher number of CO₂ molecules in the CO₂-rich liquid phase. In the first case, when $T < T_3$, the simulations evolve into a single phase of CO₂ hydrate (configurations 1 and 4). In the second case, the systems evolve into two phases, CO₂ hydrate and CO₂-rich liquid phase (configurations 2, 3, 5, and 6). Note that in the first estimation of the three-phase coexistence temperature of the CO₂ hydrate, the configuration used was non-stoichiometric (configuration 0).

Note that the dissociation line of the CO₂ hydrate is not completely analogous to that of the CH₄ hydrate considered in paper I.⁶¹ In this case, the three-phase involves the corresponding solid phase and two liquid phases, a water-rich liquid phase and a CO₂-rich liquid phase; in the CH₄ case, one of the fluid phases is not liquid since the conditions of temper-

P (bar)	Configuration							Experiment
	0	1	2	3	4	5	6	
100	284(2)	301(2)	283(2)	287(2)	289(2)	282(2)	289(2)	283.6
400	287(2)	299(2)	286(2)	289(2)	291(2)	284(2)	289(2)	286.2
1000	289(2)	299(2)	287(2)	294(2)	299(2)	286(2)	294(2)	289.7
2000	292(2)	301(2)	291(2)	296(2)	303(2)	289(2)	296(2)	292.6
3000	287(2)	301(2)	291(2)	296(2)	301(2)	291(2)	296(2)	293.7
4000	284(2)	296(2)	289(2)	293(2)	299(2)	289(2)	293(2)	293.4
5000	–	289(2)	284(2)	289(2)	296(2)	281(2)	291(2)	292.1
6000	–	279(2)	276(2)	286(2)	291(2)	276(2)	289(2)	–

TABLE II: Three-phase coexistence temperatures (T_3), at different pressures, for the CO₂ hydrate obtained for the 6 size-dependent configurations as obtained from MD-*NPT* computer simulations. Simulations for configuration 0 correspond to the original simulation obtained by Míguez *et al.*²⁸ Experimental data taken from the literature is also included in the Table.³

ature and pressure are supercritical and the CH₄-rich phase behaves as a fluid of low density that can be considered as a gas phase.

A. Effect of the stoichiometric configuration

We first consider two stoichiometric configurations, i.e., configurations in which the number of molecules of CO₂ and H₂O in the hydrate phases is the same as those in both liquid phases. The first configuration we analyze (configuration 1) is a system formed from 368 and 64 water and CO₂ molecules in the hydrate phase (all the cages of the hydrates are fully occupied, i.e., 8 CO₂ molecules per 46 water molecules in each unit cell in a $2 \times 2 \times 2$ sI structure) and the same number of molecules of each species in the liquid phases. According to the nomenclature used previously in paper I,⁶¹ this is a stoichiometric configuration. As we have already seen, it is important to compare the results obtained from simulations using stoichiometric systems with simulation data obtained using non-stoichiometric configurations. In this work, we compare the results with simulation data obtained by Míguez *et al.*²⁸ in which some of us estimated the three-phase coexistence temperature of the CO₂ hydrate using the same solid configuration ($2 \times 2 \times 2$) surrounded by liquid water (with the same number of water molecules) and a second liquid phase formed from 192 CO₂ molecules, i.e., the triple number than that of the hydrate phase. This system is the configuration 0. According to the nomenclature used in this work, configuration 0 is not stoichiometric (see Table I).

We have determined the three-phase coexistence temperature of configuration 1, T_3 , in a wide range of pressures, from 100 to 6000 bar, following the methodology already explained in paper I⁶¹ and in Section II. We first concentrate at a given pressure, 2000 bar, and then analyze the behavior of the system at different pressures. Fig. 1a shows the evolution of the potential energy of the system, U , as a function of time, at 2000 bar and temperatures from 290 to 305 K. As can be seen, two clear behaviors are observed. At the highest temperatures, 302 and 305 K, the potential energy increases very quickly over time, indicating the melting of the CO₂ hydrate. How-

ever, at low temperatures, 300, 302, and 305 K, the potential energy shows a sharp decrease, which is more pronounced as the temperature is lower, indicating that the hydrate solid phase grows. Since at 302 K the potential energy increases and at 300 K, the potential energy decreases, the three-phase coexistence temperature at 2000 bar is estimated at $T_3 = 301(2)$. It is interesting to compare this prediction with the T_3 corresponding to the configuration 0 at the same pressure. According to Table II, the T_3 obtained by Míguez *et al.*²⁸ was 292(2) K. According to this, the dissociation temperature of the stoichiometric configuration is 9 K above that of the non-stoichiometric configuration of the liquid phases. This result is in agreement with our findings in paper I,⁶¹ indicating that the use of stoichiometric configurations produces an overestimation of the T_3 .

Note that characteristic times at which the CO₂ hydrate crystallizes are shorter than those compared with the CH₄ hydrate. This is clearly seen comparing Figs. 1a of this work and Fig. 2a of paper I.⁶¹ In both cases, the system has exactly the same number of molecules in each phase. As can be seen, in the case of the CO₂, crystallization of the hydrate occurs relatively soon, between 10 and 75 ns. However, the CH₄ hydrate needs even 200 ns to crystallize if temperature is close of the T_3 . This is an expected behavior since the solubility of CO₂ in water is 10 times higher than that of CH₄ in water, approximately. This difference in solubility explains the difference in the growth times of both hydrates.^{45,62}

As we have previously mentioned, stoichiometric configurations simulated at temperatures below the T_3 of the hydrate, at the corresponding pressure, evolve into singular phases of hydrate. This is confirmed by the sharp decrease in the potential energy already shown in Fig. 1a. As we will see later, this behavior indicates that the growing mechanism of the hydrate from these configurations is not only due to a layer-by-layer formation of the hydrate. In fact, as it happens with the CH₄ hydrate,⁶¹ this behavior confirms the presence, not of a bubble of CO₂ but a liquid drop of CO₂ within the liquid phase. Recall that CO₂ and water exhibit liquid-liquid immiscibility at the conditions at which the hydrate forms.³ This liquid drop collapses, at a certain time, producing a very quick formation of the hydrate phase corroborated by the abrupt decrease in the




FIG. 1: Evolution of the potential energy, as a function of time, for the 6 size-dependent configuration analyzed in the work. The results are obtained from MD-NPT simulations at 2000 bar.

potential energy at this time. We have checked that the liquid drop mechanism is observed in the whole range of pressures considered in configuration 1. It is also important to remark on the quantitative difference found between the decrease of the potential energy of the CH₄ and CO₂ hydrates. In the first case, as can be seen in Fig. 2a of paper I,⁶¹ the sharp decrease is more pronounced than that compared with the CO₂ hydrate, as shown in Fig. 1a of this work. This indicates a quicker dissolution of the CH₄ bubbles than the CO₂ liquid drop. A more detailed account of the formation of the liquid drops in the CO₂ hydrate is presented below.

In paper I, we have concentrated on the estimation of the T_3 at a fixed pressure, 400 bar. In this work, we extend the study and consider the dissociation temperature of the CO₂ hydrate in a wide range of pressures. We have used the same procedure to evaluate the T_3 of the hydrate in a wide range of pressures, from 100 to 5000 bar. At each pressure, we have simulated several temperatures, separated 5 K, from 280 up to 310 K, approximately. The results obtained are presented in Table II (predictions obtained for the configuration 0 by Míguez *et al.*²⁸ are also included for comparison reasons).

To have an overall vision of the results obtained along the whole range of pressures, we have represented all the T_3 estimated in this work in a PT or pressure-temperature projection, as shown in Fig. 2. Particularly, Fig. 2a shows the diagram corresponding to configuration 1. We have also included the simulation results of the configuration 0, studied by Míguez *et al.*²⁸ using a non-stoichiometric configuration (see Table I for further details). As can be

seen, the main effect of using a stoichiometric instead of a non-stoichiometric configuration is to displace the whole dissociation line towards high temperatures. Particularly, the dissociation line of configuration 1 is located 9 – 14 K, depending on the pressure, above the dissociation line of configuration 0 (non-stoichiometric). Note that at 100 bar, differences between both T_3 values are 14 K. See the Table II for further details. Differences between the T_3 values of stoichiometric and non-stoichiometric ($2 \times 2 \times 2$) are much larger than in the case of the CH₄ hydrate.⁶¹

We now study a second stoichiometric configuration but considering a larger system. In this case, we replicate the unit cell of the hydrate three times ($3 \times 3 \times 3$), instead of the two times of configuration 1 ($2 \times 2 \times 2$). According to this, the initial hydrate phase is now formed from 1242 and 216 water and CO₂ molecules, respectively. In order to have the same stoichiometry in the liquid phase, we surround the initial hydrate phase by a slab with 1242 water molecules and a second slab with 216 CO₂ molecules. This system corresponds to configuration 4 presented in Table I, which is similar to configuration 6 presented in paper I for the CH₄ hydrate.⁶¹ It is interesting to compare the results obtained from this new stoichiometric configuration and check if the CO₂ hydrate shows the same general behavior.

Fig. 1d shows the evolution of the potential energy of configuration 4, as a function of time, at 2000 bar and several temperatures, from 285 to 302 K. As can be seen, there is an increase in the time needed to observe the crystallization or




FIG. 2: Pressure–temperature or PT projection of the dissociation line of the CO₂ hydrate. The green squares (a), violet down triangles (b), maroon diamonds (c), green circles (d), violet left triangles (e), and maroon right triangles (f) represent the predictions obtained from simulation using the corresponding configuration. The red circles represent the experimental data taken from the literature³ and the blue curve is the experimental vapor pressure of the CO₂. Black crosses stand for the results obtained in the work of Míguez *et al.*²⁸

melting of the hydrate. It is worthy to mention that the times required to observe one of the behaviors are not so large as in the case of the CH₄ hydrate. This indicates again that CO₂ hydrates grow much quicker than CH₄ hydrates due to the high solubility of CO₂ in water. The dissociation temperature of the hydrate at 2000 bar ranges between 302 (the lowest temperature showing an increase in potential energy) and 300 K (the highest temperature showing a decrease in potential energy). According to this, $T_3 = 301(2)$ K. The corresponding dissociation temperature of configuration 0 (non-stoichiometric), at the same pressure (2000 bar), is 292(2) K. This is the same value obtained in the case of configuration 1. The potential energy curves, as functions of time for temperatures below the T_3 , exhibit a similar behavior to those observed in configuration 1 (see Figs. 1a and 1d), with two minor differences: (1) the slope of the sharp decrease is now lower; and (2) the time required to observe the decrease is higher. The reason is due to the large size of configuration 4 compared with that of configuration 1 (3.375 times in terms of number of molecules). As we will see later, this also confirms the quick formation of liquid drops before the layer-by-layer growth of the hydrate at temperatures below the T_3 .

The complete dissociation line of configuration 4 is also presented in Fig. 2d in a pressure-temperature projection of the phase diagram as in the previous case. We have also included the results corresponding to configuration 0 obtained by Míguez *et al.*²⁸. As it happens for configuration 1 (Fig. 2a), the dissociation line of configuration 4 is shifted towards higher temperatures at all pressures. However, the displace-

ment is pressure-dependent if we compare it with the displacement observed in the case of configuration 1. Now, the shift is smaller at low pressures, similar at intermediate pressures, and larger at high pressures. This plot, with the results discussed in the previous paragraph, confirms that the liquid drop formation during the CO₂ hydrate growth is caused by the stoichiometric composition of the system, regardless of its size and the guest.

As in the case of the CH₄ hydrate study in our paper I,⁶¹ it is possible to observe the complete growth sequence of a liquid drop for configurations 1 and 4. Here we only show the formation of the CO₂ liquid drop in system 4, which corresponds to the initial $3 \times 3 \times 3$ configuration of the hydrate. It is important to mention that we have observed the same mechanism of formation of liquid drops, not only at all the pressures considered in this work (100 – 6000 bar) for this configuration but also at all of the pressures for configuration 1 ($2 \times 2 \times 2$ system). The main difference between drops observed in configuration 1 is that they exhibit a smaller size compared with those formed in configuration 3. As an example, Figure 3 shows four snapshots that illustrate the evolution of the configuration 4, at 2000 bar and 285 K. In the first snapshot (top), the initial hydrate phase is located on the left, with the pure water phase in the middle, and the CO₂ pure phase on the right (0 ns). In the second snapshot (second panel from top), a liquid drop of CO₂ is formed within the water-rich liquid phase that can be clearly observed next to the water-CO₂ interface (3 ns). It is interesting to compare this drop with the CH₄ bubble presented in paper I.⁶¹ In this case, the drop




FIG. 3: Snapshots taken at different times during 100ns simulation run for configuration 4 at 285 K and 2000 bar. The formation of a liquid drop and the growth of carbon dioxide hydrate are shown. From top to bottom: $t = 0$ ns, three-phase initial configuration. $t = 3$ ns, formation of a carbon dioxide drop within the liquid phase. $t = 4$ ns, rupture of the carbon dioxide drop and the formation of an oversaturated solution. $t = 20$ ns, complete growth of the CO₂ hydrate. Water molecules are represented as white and red sticks and CO₂ molecules as cyan (carbon atom) and blue spheres (oxygen atoms).

is more difficult to distinguish than in the methane case due to the high solubility of CO₂ in the water-rich liquid phase. Then, the liquid drop ruptures and supersaturated aqueous solution of CO₂ appears in the central region of the simulation box, as can be seen in the third snapshot from the top shown in Fig. 1d (4ns). Finally, the last snapshot (bottom) shows the complete formation of the hydrate phase (20ns). Notice that this time corresponds to the end of the slope shown in Fig. 1d when the complete growth of the hydrate occurs. As we have already mentioned, this is the expected behavior since configuration 4 corresponds to a stoichiometric system and the final state corresponds to a single hydrate phase.

The mechanism of the formation of the bubble, in the case of the CH₄ hydrate, and the liquid drop, in the case of the CO₂ hydrate, is completely analogous. However, the time scales are very different. In the first case, the formation of the bubble corresponds to the exact moment when the potential energy starts to drop abruptly (see Fig. 2f of paper I). Contrary, in the second case (CO₂), the formation of the drop starts at 0.8 ns. Note that potential energy in this case starts to drop at the beginning of the simulation. In addition to this, the temporal sequence at which the bubble forms, it ruptures, and the solid phase occupies the whole simulation box in the case of the CH₄ hydrate is 415 ns → 440 ns → 500 ns, approximately. However, in the case of the CO₂ hydrate is really different 3 ns → 4 ns → 20 ns, approximately. The reason for these differences has to be found in the high difference between the solubility of CO₂ and CH₄ in water: in this case, the high concentration of CO₂ in water acts as a driving force that enhances the diffusion of CO₂ molecules in the water-rich liquid phase, the formation of the liquid drop, and finally, the formation of the complete hydrate phase.

As we have already mentioned, in paper I we have observed the formation of CH₄ bubbles for stoichiometric configurations of the hydrate.⁶¹ Particularly, we have discussed in detail not only the formation and stability of the CH₄ bubble, but also its shape. Simulation results of paper I demonstrate that the CH₄ bubbles exhibit a cylindrical shape instead of a spherical one. An obvious question arises in this context: are the CO₂ liquid drops observed in the current simulations spherical or cylindrical, as in the case of the CH₄ hydrate? To answer this question, we analyze the snapshot presented in Fig. 3b ($t = 3$ ns) corresponding to the simulation of configuration 4 at 285 K and 2000 bar. It is important to recall again that we have observed the formation of liquid drops of CO₂, not only at this particular pressure but in the whole range of pressures along the dissociation line of the hydrate. Particularly, we have also observed the drops in configuration 1 ($2 \times 2 \times 2$). However, since the solubility of CO₂ in water is high (10 times higher than that of CH₄ in water, approximately), it is more difficult to distinguish the drops. For this reason, we concentrate here in liquid drops of configuration 4.

Fig. 4 shows the projection of the xz plan (top), as well as the yz projection (bottom) of the same configuration. To help the visualization of the drop we have omitted the water molecules. As can be seen, the second projection (yz) clearly shows that the liquid drop extends along the whole y -axis of the simulation box. As it discussed in paper I,⁶¹ and accord-




FIG. 4: Snapshots of the xz (top) and yz (bottom) planes taken for configuration 4 at 285 K and 2000 bar, showcasing the emergence of a CO₂ liquid drop within the liquid phase at $t = 3$ ns. For clarity, water molecules are omitted, while CO₂ molecules are represented as cyan (carbon atom) and blue spheres (oxygen atoms). CO₂ molecules are visible within the hydrates on the left and right sides of the figure. In the central region, CO₂ molecules are shown forming a cylindrical liquid drop.

ing to the Laplace equation, cylindrical interfaces results in lower solubility than spherical ones but higher than planar interfaces. If the liquid drops were stable and sufficiently large, the dissociation temperature would be shifted towards higher temperatures compared with the planar interface. In this work, as in paper I, drops of CO₂ are only stable for 1 – 20, ns, even less than in the case of the CH₄ bubbles (10 – 25 ns).

In summary, as it happens in the case of the CH₄ hydrate, the presence of liquid drops of CO₂ modifies the prediction of the T_3 of CO₂ hydrates. This is only observed in liquid configurations that have stoichiometric configurations, such as configurations 1 and 4 studied in this work. Hence, we do not recommend the use of stoichiometric configurations in order to get reliable predictions of the T_3 of CO₂ hydrates. Particularly, the recommendation for future studies is to use the configuration 4 ($3 \times 3 \times 3$), which is non-stoichiometric, formed from an appropriate number of water and CO₂ molecules that allow to simulate the system in a reasonable time.

Before finishing this section, it is important to briefly mention how simulation results obtained from stoichiometric configurations 1 and 4 compare with experimental data taken from the literature. As can be seen in Figs. 2a and 2d, simulation predictions exhibit large deviations with respect to exper-

imental data. This is expected since, according to our previous discussion, the use of these kinds of configurations produces an overestimation of the T_3 at all the pressures along the dissociation line of the hydrate. We shall discuss in detail this issue in the last section, where we present the finite-size effect of non-stoichiometric configuration on the T_3 along the dissociation line of the CO₂ hydrate.

B. Effect of the overall size

Once we have analyzed in detail the effect of using two different stoichiometric configurations to determine the dissociation temperature of the CO₂ hydrate in a wide range of pressures, we now turn on to study the finite-size effects on T_3 for non-stoichiometric configurations. In this first section, we concentrate on finite-size effects due to the same increase in the number of molecules in each of the phases involved.

The control configuration used in this section is again configuration 0, with 992 total number of molecules, and an initial configuration formed from 432 (368 + 64) water and CO₂ molecules in the hydrate phase, 368 water molecules in the water phase, and 192 CO₂ molecules in the CO₂-rich liquid phase. Note that the non-stoichiometric composition of the liquid phases is verified since the number of CO₂ molecules in that phase is three times the number corresponding to the stoichiometric composition ($192 = 3 \times 64$). Note that the size of the interfacial area and the length of the simulation box perpendicularly to the interface also increase. We first determine the dissociation temperatures, in the whole range of pressures already considered in the previous section, of configuration 3. According to Table I, this configuration contains a total number of molecules equal to 3348. This means that the system size, in terms of the number of molecules, is multiplied by a factor of 3.375, keeping the non-stoichiometric composition of the liquid phases. As can be seen in Table I, the number of molecules of each species is multiplied by this factor in each of the phases forming the initial simulation box of configuration 3.

We have determined the three-phase coexistence temperature of configuration 3, in the same range of pressures considered previously (from 100 to 6000 bar). Fig. 1c shows the evolution of the potential energy of the system, U , as a function of time, at 2000 bar and temperatures from 285 to 305 K. As in the case of the stoichiometric configurations, we observe the same two behaviors: at the highest temperatures, 298, 300, and 305 K, the potential energy increases very quickly over time, indicating the melting of the CO₂ hydrate. However, at low temperatures, 295, 290, and 285 K, the potential energy shows a decrease, which is more pronounced as the temperature is lower, indicating that the hydrate solid phase is growing. The three-phase coexistence temperature at 2000 bar is estimated at $T_3 = 296(2)$ (at 298 K the potential energy increases and at 295 K, it decreases).

It is interesting to compare the evolution of the potential energy obtained in this configuration and those corresponding to the stoichiometric configurations, 1 and 4. Although system sizes are different, it is clear that the characteristics

sharp decreases observed in Figs. 1a and 1d are not seen in Fig. 1c. The reason, as clearly stated in paper I,⁶¹ is that configuration 3 is not stoichiometric. According to this, the T_3 value corresponding to this configuration is reliable and can be compared with confidence with that obtained by Míguez *et al.*²⁸ for configuration 0. According to Table II, the T_3 at 2000 bar was 292(2) K (configuration 0). In other words, the dissociation temperature of the configuration 3 is 4 K above that of the configuration 0. This result suggests that there is a finite-size effect on T_3 that displaces the dissociation temperature towards higher temperatures, making the hydrate phase more stable at this pressure.

To confirm the stabilization of the hydrate when the number of molecules considered in the simulation box is increased (by a factor of 3.375), we have determined the T_3 at the whole range of pressures, from 100 to 6000 bar. Results are presented in Fig. 2c. As can be seen, the dissociation line corresponding to configuration 3 (maroon diamonds) is systematically displaced with respect to the results obtained from configuration 0 (black crosses). Simulation data obtained from MD-NPT simulations is also included in Table II. Displacement is not homogeneous. At low pressures (below 400 bar), the T_3 increases ~ 2 K, resulting in good agreement with previous results within the error bars. However, as the pressure is increased, displacement increases $\sim 4 - 5$ K at intermediate pressures (1000 and 2000 bar). Finally, at the highest pressures (≥ 3000 bar), the T_3 is shifted 7 - 9 K.

To better understand the finite-size effect on the location of the dissociation line of the CO₂ hydrate, we go further and consider a new larger system. Configuration 6 is formed from a total number of molecules equal to 7936, 3456 corresponding to the (stoichiometric) hydrate phase, and 4480 molecules to the non-stoichiometric liquid phases. Note that now, the total number of molecules in configure 6 is 8 times larger than in configuration 0. Details of the particular number of water and CO₂ molecules can be inspected in Table I. As in the previous cases, we first concentrate on the determination of the T_3 at an intermediate pressure, 2000 bar. As can be seen in Fig. 1f, the system melts at the two highest temperatures considered, 298 and 300 K, and freezes at 295 and 290 K. According to this, the T_3 , at 2000 bar, is 296(2) K. Similarly to what happens with the previous non-stoichiometric configuration 3 (Fig. 1c), the evolution of the potential energy, as a function of time, behaves without the characteristic sharp drop associated with configuration 1 and 4 shown in Figs. 1a and 1d. Note that now, the time required to observe crystallization is much larger than in smaller systems: in configuration 3, we observe crystallization before 100 ns. However, in configuration 6 we see that at temperatures close to the T_3 , i.e., 295 K, crystallization occurs for times higher than 300 ns.

According to the results discussed in the previous paragraph, the T_3 predicted in configurations 3 and 6 are equal, 296(2) K. Taking into account the results obtained for the CH₄ hydrate in paper I,⁶¹ an obvious question arises: is it necessary to simulate systems as large as the configuration 6 to predict T_3 values without finite-size effects? To answer this question, we have obtained the rest of the dissociation temperatures at different pressures. The results are presented in

Fig. 2f. As can be seen, there is a similar shift of the dissociation line of configuration 6 (maroon right triangles) with respect to that of configuration 0 (black crosses) in Figs. 2c and 2f, indicating that we have achieved an asymptotic limit. A careful inspection of Table II confirms this hypothesis: except for the T_3 value at 100 bar, the T_3 values of configurations 3 and 6 are identical (within the error bars). In summary, configurations 3 and 6, formed from the larger unit cells (e.g., $3 \times 3 \times 3$ and $4 \times 4 \times 4$) show convergence of the T_3 values. This suggests that no finite-size effects exist for these system sizes.

C. Effect of size of the interfacial area

In the previous section, we have analyzed the finite-size effects of non-stoichiometric configurations on the dissociation line of the CO₂ hydrate varying the number of molecules in the system isotropically, i.e., increasing the initial hydrate phase in the three directions ($2 \times 2 \times 2$, $3 \times 3 \times 3$, and $4 \times 4 \times 4$). In addition to this, we consider the same number of water molecules in the aqueous phase and with three times more molecules of CO₂ in the CO₂-rich liquid phase to avoid stoichiometric configurations in the liquid phases. We have demonstrated that there exist finite-size effects associated with these changes.

We now investigate if the size of the interfacial area of the simulation box affects the T_3 of the CO₂ hydrate. To this end, we proceed similarly but considering two configurations in which the initial hydrate phases are formed replicating the unit cell as $3 \times 3 \times 2$ (configuration 2) and $4 \times 4 \times 2$ (configuration 5). The corresponding liquid phases are formed from the same number of water molecules in the aqueous phase as in the hydrate and with the number of CO₂ molecules equal to three times those existing in the hydrate. The particular number of molecules in each configuration can be seen in Table I. Note that the total number of molecules in configurations 2 and 5 are 2232 and 3969, respectively.

Figs. 1b and 1e show the evolution of the potential energy of configurations 2 and 5, as functions of time, at 2000 bar, respectively. The same general trend is observed in both figures, similar to those presented by configurations 3 and 6 (Fig. 1c and 1f), i.e., since both configurations are non-stoichiometric they do not show the sharp decrease in potential energy. Following the same procedure used in previous sections, T_3 of both configurations can be easily determined by inspecting the behavior of potential energy at different temperatures. The T_3 values predicted by the molecular models are 291(2) and 289(2) K for configurations 2 and 5, respectively. These results suggest that the T_3 values of both configurations are equal (within the error bars) and also equal to the T_3 value corresponding to the configuration 0 ($2 \times 2 \times 2$) at this pressure. This is a non-expected result according to the behavior observed in the previous section, especially if we take into account that configuration 5 is formed from 3968 molecules, a 20% more molecules than in configuration 3 ($3 \times 3 \times 3$). For this former configuration, T_3 is equal to 296(2) K at the same pressure.

To clarify this point, we have also obtained the T_3 values of configurations 2 and 5 at lower and higher pressures, as we have already done with the configurations previously studied. Fig. 2c and 2f show the pressure-temperature projection of the dissociation line of the CO₂ hydrate using both configurations. The numerical data obtained from MD-NPT simulations are also presented in Table II. As can be seen in the plots, both configurations exhibit very similar T_3 values in the whole range of pressures. Particularly, at low pressures, between 100 and 2000 bar, configurations 0 (black crosses) and 3 (violet down triangles) present the same T_3 values, while configuration 5 (violet left triangles) seem to exhibit slightly lower values of T_3 . However, both results are within the error bars. At higher pressure, above 3000 bar, the dissociation temperatures of configurations 2 and 5 are above those of configuration 0. Nevertheless, the results are again within the error bars and no significant differences are observed between temperatures at the corresponding pressure (see Table II for further details).

In summary, our simulation results indicate that finite-size effects of the T_3 for configurations 2 ($3 \times 3 \times 2$) and 5 ($4 \times 4 \times 2$) are less important or even non-existing compared with the case of configurations 3 ($3 \times 3 \times 3$) and 6 ($4 \times 4 \times 4$). This effect occurs even for configuration 5, which is formed from a larger number of molecules than configuration 2. It is possible that the increase of the number of molecules done anisotropically, i.e., a small increase of the simulation lengths perpendicular to the interface compared with the increase in the interfacial area plane, has less effect of T_3 .

D. Comparison with experimental data

In the previous sections, we discuss the finite-size effects on the T_3 of the CO₂ hydrate in a wide range of pressures. This is performed using 6 different configurations and comparing the results with the configuration 0 obtained several years ago by Míguez *et al.*²⁸ In this Section, we focus on the comparison of the predictions obtained using these configurations with experimental data taken from the literature.

In the original work,²⁸ the authors proposed a modified Berthelot combining rule that allows to predict the dissociation line of the CO₂ hydrate in a wide range of pressures with confidence. Particularly, differences between predictions obtained from configuration 0 and experimental data taken from the literature³ are below 1 K from 100 to 2000 K. At high pressures, at 3000 bar and above, agreement between both results deteriorates. Although the model is able to capture the existence of the reentrant behavior of the dissociation line, the agreement is only qualitative. We recommend the reader to the original paper for a detailed account of this issue.²⁸

Fig. 2 shows the pressure-temperature composition of the dissociation line of the CO₂ hydrate as predicted from the model proposed by Míguez *et al.*²⁸ using the 6 configurations considered in this work. The results obtained using configuration 0 are also presented, as well as the experimental data taken from the literature.³ We only analyze the agreement between experimental data and predictions obtained using non-




FIG. 5: Comparison of the three-phase coexistence temperatures (T_3) for the CO₂ hydrate from the 6 size-dependent configurations analyzed in this work at several pressures. The experimental T_3 values taken from the literature,³ at different pressures, are also included. In all cases, the composition of the initial configuration is analogous, formed by a CO₂ hydrate phase in coexistence with a liquid water phase and a CO₂ liquid phase with the corresponding numbers of molecules in each configuration.

stoichiometric configurations (2, 3, 5, and 6) since stoichiometric systems do not provide the correct T_3 due to the presence of CO₂ liquid drops discussed in Section III. A. Let's concentrate on predictions obtained using configurations 3 and 6. As can be seen, predictions from configuration 3 overestimate the experimental T_3 in the whole range of pressures. Particularly, T_3 is overestimated ~ 5 K at low pressures, below 3000 bar. The agreement between predictions from configuration 6 and experimental data is similar.

The main conclusion is that finite-size effects also affect agreement with experimental data. What is the reason for this? The answer is simple. The unlike interaction parameter associated with the Berthelot rule, ξ , was established by Míguez *et al.*²⁸ using configuration 0, i.e., using a $2 \times 2 \times 2$ configuration and a certain cutoff distance for the dispersive interactions. We will not discuss here the effect of the cutoff distance on the T_3 of hydrates. We recommend the reader paper III of this series in which we analyze the effect of the dispersive interactions on the three-phase equilibria of CH₄ and CO₂ hydrates.⁶³ According to this, the value $\xi = 1.13$ obtained by Míguez *et al.*²⁸ allows to quantitatively predict the dissociation line of the CO₂ hydrate for the particular configuration ($2 \times 2 \times 2$) used to fit experimental data. As we have demonstrated in Section III. B, the use of larger configurations, i.e., configurations 3 and 6, produces a displacement of the dissociation line of the CO₂ towards higher temperatures, making the hydrate phase more stable. In fact, the same effect produces the use of a positive modification of the Berthelot rule ($\xi > 1$). This suggests that the $\xi = 1.13$ value is too high for describing in a quantitative way the dissociation line of the CO₂ hydrate using configurations not affected by finite-size effects, i.e., configurations 3 or 6.

Finally, we have summarized the T_3 values obtained for each of the 6 size-dependent configurations, at all the pressure considered (from 100 to 6000 bar), the in bar graphs of Fig. 5. In addition, we have also included the experimental T_3 for comparison reasons (first column). The order of the columns, from left to right, takes into account the effect of the stoichiometric configurations (1 and 4), of the overall size of the system (3 and 6), and of the size of the interfacial area (2 and 5). The first configurations in each of the three groups, 1, 2, and 3, correspond to the smaller system analyzed, and the second ones, 4, 5 and 6, to the larger systems. See Table I for further details. As can be seen, the main conclusions stated in the previous sections can be observed in Fig. 5 in the whole range of pressures analyzed in this work: (1) overestimation of the T_3 values, with respect to the experimental data taken from the literature, in the case of the stoichiometric configurations (1 and 4); (2) similar T_3 values for configurations 3 ($3 \times 3 \times 3$) and 6 ($3 \times 3 \times 3$); and (3) configurations 2 and 5 exhibit the same T_3 values at all pressures (within the error bars) but below the experimental T_3 values.

IV. CONCLUDING REMARKS

In this second part of a three-paper series dedicated to investigating finite-size effects in methane and carbon dioxide,

as well as the influence of dispersive interactions on both hydrates, our focus is on the CO₂ hydrate. Specifically, we delve into the finite-size effects affecting the determination of the three-phase coexistence temperature (T_3) for CO₂ hydrate, employing molecular dynamics simulations in conjunction with the direct coexistence technique. Adhering to the methodology outlined in the first paper⁶¹, we examine six size-dependent configurations using realistic water and CO₂ models (i.e., TIP4P/Ice and TraPPE) to assess the impact of size and composition on the estimation of T_3 . Given the similarity of our findings to those in the first paper⁶¹, we provide a concise summary of the most pertinent results:

- The simulation results obtained for the CO₂ hydrate confirm the sensitivity of T_3 depending on the size and composition of the system, explaining the discrepancies observed in the original work by Míguez *et al.*²⁸ in 2015. This is in agreement with the findings of paper I,⁶¹ not only at a particular pressure but in a wide range of pressures considered.
- Configurations with stoichiometric composition at temperatures below T_3 evolve into a singular phase of CO₂ hydrate growth, as in the case of the methane hydrate.⁶¹ This is confirmed in the whole dissociation line, from 100 to 6000 bar. In this case, there is no a bubble of methane but a liquid drop of CO₂. The mechanism of growing is via the emergence of a liquid drop of CO₂ within the liquid and the subsequent formation of an oversaturated CO₂ solution in water. Conversely, an excess of molecules in the CO₂-rich liquid phase in the initial configuration leads to the coexistence of CO₂ hydrate and CO₂ liquid phases without the formation of drops.
- Finite-size effects are pronounced in small systems with stoichiometric composition (e.g., configuration 1 with a unit cell of $2 \times 2 \times 2$), resulting in an overestimation of T_3 due to liquid drop formation during hydrate growth, causing a false stability of CO₂ hydrate by increasing CO₂ solubility. Note that this is a common conclusion in both methane and carbon dioxide hydrates.
- Non-stoichiometric configurations with larger unit cells, like $3 \times 3 \times 3$ and $4 \times 4 \times 4$, show convergence of T_3 values, suggesting that finite-size effects for these system sizes, regardless of drop formation, can be safely neglected.
- To study the T_3 of the CO₂ hydrate the best choice is configuration 3, which provides an accurate T_3 value, and affordable simulation times.

The primary outcome of this study, entirely consistent with the findings in the initial paper of this series for CH₄ hydrate⁶¹, can be succinctly summarized as follows: when using the direct coexistence technique to estimate the T_3 of CO₂ hydrate, it is crucial to avoid small stoichiometric configurations such as configuration 1. These configurations tend to

develop drops at the onset of the run, leading to an overestimation of the T_3 value. The recommended and optimal choice is configuration 3, which provides an accurate T_3 value, and the computational resources required for simulating this system are currently feasible.

This study, complementing the initial paper in the series⁶¹, presents valuable insights into the finite-size effects observed in simulations of CO₂ hydrate. The findings highlight the possibility of mitigating finite-size effects in estimating T_3 by thoughtfully selecting system configurations. We anticipate that these results will contribute to a better understanding of finite size effects in the determination of T_3 for methane hydrates, addressing discrepancies in the existing literature and assisting researchers in selecting appropriate system sizes for future investigations. Our future research will delve into examining the potential impact of cutoff values and guest types on T_3 values, exploring how these factors are influenced by finite-size effects.

ACKNOWLEDGMENTS

This work was funded by Ministerio de Ciencia e Innovación (Grant No. PID2019-105898GA-C22, PID2021-125081NB-I00 and PID2022-136919NB-C32), Junta de Andalucía (P20-00363), and Universidad de Huelva (P.O. FEDER UHU-1255522 and FEDER-UHU-202034), all four cofinanced by EU FEDER funds. This work was also funded by Project No. ETSII-UPM20-PU01 from “Ayudas Primeros Proyectos de la ETSII-UPM”. M.M.C. acknowledges CAM and UPM for financial support of this work through the CavI-tieS project No. APOYO-JOVENES-01HQ1S-129-B5E4MM from “Accion financiada por la Comunidad de Madrid en el marco del Convenio Plurianual con la Universidad Politecnica de Madrid en la linea de actuacion estimulo a la investigacion de jovenes doctores” and CAM under the Multiannual Agreement with UPM in the line Excellence Programme for University Professors, in the context of the V PRICIT (Regional Programme of Research and Technological Innovation). S.B. acknowledges Ayuntamiento de Madrid for a Residencia de Estudiantes grant. The authors also gratefully acknowledge the Universidad Politecnica de Madrid (www.upm.es) for providing computing resources on Magerit Supercomputer. We also acknowledge and additional computational resources from Centro de Supercomputación de Galicia (CESGA, Santiago de Compostela, Spain), at which some of the simulations were run.

AUTHORS DECLARATIONS

CONFLICTS OF INTEREST

The authors have no conflicts to disclose.

DATA AVAILABILITY

The data that support the findings of this study are available within the article.

REFERENCES

- E. D. Sloan, “Fundamental principles and applications of natural gas hydrates,” *Science* **426**, 353–359 (2003).
- C. A. Koh, A. K. Sum, and E. D. Sloan, “State of the art: Natural gas hydrates as a natural resource,” *J. Nat. Gas Sci. Eng.* **8**, 132–138 (2012).
- E. D. Sloan and C. Koh, *Clathrate Hydrates of Natural Gases*, 3rd ed. (CRC Press, New York, 2008).
- J. A. Ripmeester and S. Alavi, *Clathrate Hydrates: Molecular Science and Characterization* (Wiley-VCH: Weinheim, Germany, 2022).
- H. Barthélémy, M. Weber, and F. Barbier, “Hydrogen storage: Recent improvements and industrial perspectives,” *International Journal of Hydrogen Energy* **42**, 7254–7262 (2017).
- S. Chen, Y. Wang, X. Lang, S. Fan, and G. Li, “Rapid and high hydrogen storage in epoxy-cyclopentane hydrate at moderate pressure,” *Energy* **268**, 126638 (2023).
- Y. Zhang, G. Bhattacharjee, J. Zheng, and P. Linga, “Hydrogen storage as clathrate hydrates in the presence of 1, 3-dioxolane as a dual-function promoter,” *Chemical Engineering Journal* **427**, 131771 (2022).
- K. A. Kvenvolden, “Methane hydrate - A major reservoir of carbon in the shallow geosphere,” *Chem. Geol.* **71**, 41 (1988).
- G. J. MacDonald, “The future of methane as an energy resource,” *Annu. Rev. Energy* **15**, 53 (1990).
- C. Bourry, J. L. Charlou, J. P. Donval, M. Brunelli, C. Focsa, and B. Chazallon, “X-ray synchrotron diffraction study of natural gas hydrates from african margin,” *Geophys. Res. Lett.* **34**, L22303 (2007).
- H. Lu, Y. Seo, J. Lee, I. Moudrakovski, J. A. Ripmeester, N. R. Chapman, R. B. Coffin, G. Gardner, and J. Pohlman, “Complex gas hydrate from the cascadia margin,” *Nature* **445**, 303 (2007).
- B. Lal and O. Nashed, *Chemical Additives for Gas Hydrates* (Springer, 2020).
- M. M. Ghiasi, A. H. Mohammadi, and S. Zandehboudi, “Modeling stability conditions of methane clathrate hydrate in ionic liquid aqueous solutions,” *J. Mol. Liq.* **325**, 114804 (2021).
- H. Tanaka, T. Yagasaki, and M. Matsumoto, “On the occurrence of clathrate hydrates in extreme conditions: Dissociation pressures and occupancies at cryogenic temperatures with application to planetary systems,” *Planet. Sci. J.* **1**, 80 (2020).
- M. M. Conde, M. Rovere, and P. Gallo, “Spontaneous NaCl-doped ice at seawater conditions: focus on the mechanisms of ion inclusion,” *Phys. Chem. Chem. Phys.* **19**, 9566 (2017).
- M. M. Conde, M. Rovere, and P. Gallo, “Spontaneous nacl-doped ices ih, ic, iii, v and vi. understanding the mechanism of ion inclusion and its dependence on the crystalline structure of ice,” *Phys. Chem. Chem. Phys.* **23**, 22897–22911 (2021).
- O. Prieto-Ballesteros, J. S. Kargel, M. Fernández-Sampedro, F. Selsis, E. S. Martínez, and D. L. Hogenboom, “Evaluation of the possible presence of clathrate hydrates in europa’s icy shell or seafloor,” *Icarus* **177**, 491–505 (2005), europa Icy Shell.
- J. S. Kargel, J. Z. Kaye, J. W. Head, G. M. Marion, R. Sassen, J. K. Crowley, O. P. Ballesteros, S. A. Grant, and D. L. Hogenboom, “Europa’s crust and ocean: Origin, composition, and the prospects for life,” *Icarus* **148**, 226–265 (2000).
- O. Prieto-Ballesteros, J. S. Kargel, A. G. Faireén, D. C. Fernández-Remolar, J. M. Dohm, and R. Amils, “Interglacial clathrate destabilization on Mars: Possible contributing source of its atmospheric methane,” *Geology* **34**, 149–152 (2006).
- E. Pettinelli, B. Cosciotti, F. Di Paolo, S. E. Lauro, E. Mattei, R. Orosei, and G. Vannaroni, “Dielectric properties of jovian satellite ice analogs for subsurface radar exploration: A review,” *Reviews of Geophysics* **53**, 593–641 (2015).

- ²¹S. C. Peter, "Reduction of CO₂ to chemicals and fuels: a solution to global warming and energy crisis," *ACS Energy Letters* **3**, 1557–1561 (2018).
- ²²K. O. Yoro and M. O. Daramola, "CO₂ emission sources, greenhouse gases, and the global warming effect," in *Advances in carbon capture* (Elsevier, 2020) pp. 3–28.
- ²³D. M. D'Alessandro, B. Smit, and J. R. Long, "Carbon dioxide capture: prospects for new materials," *Angewandte Chemie International Edition* **49**, 6058–6082 (2010).
- ²⁴W. Gao, S. Liang, R. Wang, Q. Jiang, Y. Zhang, Q. Zheng, B. Xie, C. Y. Toe, X. Zhu, J. Wang, *et al.*, "Industrial carbon dioxide capture and utilization: state of the art and future challenges," *Chemical Society Reviews* **49**, 8584–8686 (2020).
- ²⁵X. Wang, F. Zhang, and W. Lipiński, "Research progress and challenges in hydrate-based carbon dioxide capture applications," *Applied Energy* **269**, 114928 (2020).
- ²⁶N. N. Nguyen, V. T. La, C. D. Huynh, and A. V. Nguyen, "Technical and economic perspectives of hydrate-based carbon dioxide capture," *Applied Energy* **307**, 118237 (2022).
- ²⁷J. Zheng, Z. R. Chong, M. F. Qureshi, and P. Linga, "Carbon dioxide sequestration via gas hydrates: a potential pathway toward decarbonization," *Energy & Fuels* **34**, 10529–10546 (2020).
- ²⁸J. M. Míguez, M. M. Conde, J.-P. Torré, F. J. Blas, M. M. Piñeiro, and C. Vega, "Molecular dynamics simulation of CO₂ hydrates: Prediction of three phase coexistence line," *J. Chem. Phys.* **142**, 124505 (2015).
- ²⁹Y.-T. Tung, L.-J. Chen, Y.-P. Chen, and S.-T. Lin, "Growth of structure I carbon dioxide hydrate from molecular dynamics simulations," *The Journal of Physical Chemistry C* **115**, 7504–7515 (2011).
- ³⁰H. W. Horn, W. C. Swope, J. W. Pitera, J. D. Madura, T. J. Dick, G. L. Hura, and T. Head-Gordon, "Development of an improved four-site water model for biomolecular simulations: Tip4p-ew," *The Journal of chemical physics* **120**, 9665–9678 (2004).
- ³¹J. G. Harris and K. H. Yung, "Carbon dioxide's liquid-vapor coexistence curve and critical properties as predicted by a simple molecular model," *The Journal of Physical Chemistry* **99**, 12021–12024 (1995).
- ³²J. Costandy, V. K. Michalisa, I. N. Tsimpanogiannis, A. K. Stubos, and I. G. Economou, "The role of intermolecular interactions in the prediction of the phase equilibria of carbon dioxide hydrates," *J. Chem. Phys.* **143**, 094506 (2015).
- ³³M. H. Waage, T. J. H. Vlught, and S. Kjelstrup, "Phase diagram of methane and carbon dioxide hydrates computed by Monte Carlo simulations," *J. Phys. Chem. B* **121**, 7336–7350 (2017).
- ³⁴L. Jiao, Z. Wang, J. Li, P. Zhao, and R. Wan, "Stability and dissociation studies of CO₂ hydrate under different systems using molecular dynamic simulations," *Journal of Molecular Liquids* **338**, 116788 (2021).
- ³⁵X. Hao, C. Li, Q. Meng, J. Sun, L. Huang, Q. Bu, and C. Li, "Molecular dynamics simulation of the three-phase equilibrium line of CO₂ hydrate with opc water model," *ACS omega* **8**, 39847–39854 (2023).
- ³⁶N. Qiu, X. Bai, N. Sun, X. Yu, L. Yang, Y. Li, M. Yang, Q. Huang, and S. Du, "Grand canonical monte carlo simulations on phase equilibria of methane, carbon dioxide, and their mixture hydrates," *The Journal of Physical Chemistry B* **122**, 9724–9737 (2018).
- ³⁷S. El Meragawi, N. I. Diamantonis, I. N. Tsimpanogiannis, and I. G. Economou, "Hydrate–fluid phase equilibria modeling using pc-saft and peng–robinson equations of state," *Fluid Phase Equilibria* **413**, 209–219 (2016).
- ³⁸A. Jäger, V. Vinš, J. Gernert, R. Span, and J. Hrubý, "Phase equilibria with hydrate formation in H₂O+ CO₂ mixtures modeled with reference equations of state," *Fluid Phase Equilibria* **338**, 100–113 (2013).
- ³⁹L. Sun, H. Zhao, S. B. Kiselev, and C. McCabe, "Predicting Mixture Phase Equilibria and Critical Behavior Using the SAFT-VRX Approach," *Fluid Phase Equilib.* **228–229**, 275–282 (2005), **Sun2005b**.
- ⁴⁰S. Blázquez, M. M. Conde, C. Vega, and E. Sanz, "Growth rate of CO₂ and CH₄ hydrates by means of molecular dynamics simulations," *The Journal of Chemical Physics* **159** (2023).
- ⁴¹H. Wang, Y. Lu, X. Zhang, Q. Fan, Q. Li, L. Zhang, J. Zhao, L. Yang, and Y. Song, "Molecular dynamics of carbon sequestration via forming CO₂ hydrate in a marine environment," *Energy & Fuels* (2023).
- ⁴²A. M. Fernández-Fernández, M. Pérez-Rodríguez, A. Comesaña, and M. M. Piñeiro, "Three-phase equilibrium curve shift for methane hydrate in oceanic conditions calculated from molecular dynamics simulations," *J. Mol. Liq.* **274**, 426–433 (2019).
- ⁴³S. Blázquez, C. Vega, and M. M. Conde, "Three phase equilibria of the methane hydrate in NaCl solutions: A simulation study," *J. Mol. Liq.* **383**, 122031 (2023).
- ⁴⁴J. Grabowska, S. Blázquez, E. Sanz, E. G. Noya, I. M. Zerón, J. Algaba, J. M. Míguez, F. J. Blas, and C. Vega, "Homogeneous nucleation rate of methane hydrate formation under experimental conditions from seeding simulations," *J. Chem. Phys.* **158** (2023).
- ⁴⁵J. Algaba, I. M. Zerón, J. M. Míguez, J. Grabowska, S. Blázquez, E. Sanz, C. Vega, and F. J. Blas, "Solubility of carbon dioxide in water: Some useful results for hydrate nucleation," *The Journal of Chemical Physics* **158** (2023).
- ⁴⁶A. Z. Panagiotopoulos, "Molecular simulation of phase coexistence: Finite-size effects and determination of critical parameters for two- and three-dimensional Lennard-Jones fluids," *International journal of thermophysics* **15**, 1057–1072 (1994).
- ⁴⁷P. Orea, J. López-Lemus, and J. Alejandro, "Oscillatory surface tension due to finite-size effects," *The Journal of chemical physics* **123** (2005).
- ⁴⁸K. Binder and M. Müller, "Computer simulation of profiles of interfaces between coexisting phases: Do we understand their finite size effects?" *International Journal of Modern Physics C* **11**, 1093–1113 (2000).
- ⁴⁹H. L. Vörtler, K. Schäfer, and W. R. Smith, "Simulation of chemical potentials and phase equilibria in two- and three-dimensional square-well fluids: finite size effects," *The Journal of Physical Chemistry B* **112**, 4656–4661 (2008).
- ⁵⁰M. M. Conde, M. Rovere, and P. Gallo, "High precision determination of the melting points of water TIP4P/2005 and water TIP4P/Ice models by the direct coexistence technique," *J. Chem. Phys.* **147**, 244506 (2017).
- ⁵¹M. M. Conde and C. Vega, "Determining the three-phase coexistence line in methane hydrates using computer simulations," *J. Chem. Phys.* **133**, 064507 (2010).
- ⁵²V. Buch, P. Sandler, and J. Sadlej, "Simulations of H₂O solid, liquid, and clusters, with an emphasis on ferroelectric ordering transition in hexagonal ice," *J. Phys. Chem. B* **102**, 8641–8653 (1998).
- ⁵³J. D. Bernal and R. H. Fowler, "Simulations of H₂O solid, liquid, and clusters, with an emphasis on ferroelectric ordering transition in hexagonal ice," *J. Chem. Phys.* **1**, 515–548 (1933).
- ⁵⁴D. van der Spoel, E. Lindahl, B. Hess, G. Groenhof, A. E. Mark, and H. J. Berendsen, "Gromacs: Fast, flexible, and free," *J. Comput. Chem.* **26**, 1701–1718 (2005).
- ⁵⁵J. J. Potoff and J. I. Siepmann, "Vapor-liquid equilibria of mixtures containing alkanes, carbon dioxide, and nitrogen," *AIChE Journal* **47**, 1676–1682 (2001).
- ⁵⁶J. L. F. Abascal, E. Sanz, R. G. Fernández, and C. Vega, "A potential model for the study of ices and amorphous water: TIP4P/Ice," *J. Chem. Phys.* **122**, 234511–1–234511–9 (2005).
- ⁵⁷M. A. Cuendet and W. F. V. Gunsteren, "On the calculation of velocity-dependent properties in molecular dynamics simulations using the leapfrog integration algorithm," *J. Chem. Phys.* **127**, 184102/1–9 (2007).
- ⁵⁸S. Nosé, "A molecular dynamics method for simulations in the canonical ensemble," *Mol. Phys.* **52**, 255–268 (1984).
- ⁵⁹M. Parrinello and A. Rahman, "Polymorphic transitions in single crystals: A new molecular dynamics method," *J. Appl. Phys.* **52**, 7182–7190 (1981).
- ⁶⁰U. Essmann, L. Perera, M. L. Berkowitz, T. Darden, H. Lee, and L. G. Pedersen, "A smooth particle mesh Ewald method," *J. Chem. Phys.* **103**, 8577–8593 (1995).
- ⁶¹S. Blázquez, J. Algaba, J. M. Míguez, C. Vega, F. J. Blas, and M. M. Conde, "Three-phase equilibria of hydrates from computer simulation. I. Finite-size effects in the methane hydrate," *The Journal of Chemical Physics*, submitted (2024).
- ⁶²J. Grabowska, S. Blázquez, E. Sanz, I. M. Zerón, J. Algaba, J. M. Míguez, F. J. Blas, and C. Vega, "Solubility of methane in water: some useful results for hydrate nucleation," *J. Phys. Chem. B* **126**, 8553–8570 (2022).
- ⁶³J. Algaba, S. Blázquez, J. M. Míguez, M. M. Conde, and F. J. Blas, "Three-phase equilibria of hydrates from computer simulation. III. Effect of dispersive interactions in methane and carbon dioxide hydrates," *The Journal of Chemical Physics*, submitted (2023).